



Received on 21 October 2017; received in revised form, 14 November 2017; accepted, 18 November 2017; published 01 March 2018

EVALUATION OF *IN-VITRO* ANTIOXIDANT AND ANTI-PROLIFERATIVE EFFECT OF DEFATTED EXTRACT OF *COCOS NUCIFERA* KERNEL ON HUMAN BREAST CANCER CELL LINE

Radhakrishnan Dhanyakrishnan ¹, Mary Chacko Sunitha ¹, Bhaskara Prakashkumar ¹, Sukumaran Sandya ^{2,3} and Kottayath Govindan Nevin ^{*1}

School of Biosciences ¹, Mahatma Gandhi University, PD Hills PO, Kottayam - 686560, Kerala, India. Inorganic and Physical Chemistry ², Spectroscopy / Analytical / Test Facility, Entrepreneurship Centre, Society for Innovation and Development (SID) ³, Indian Institute of Science, Bangalore - 560012, Karnataka, India.

Keywords:

Coconut kernel,
Cytotoxicity, ROS,
MCF-7, Free Radicals

Correspondence to Author:

Dr. Kottayath Govindan Nevin

School of Biosciences,
Mahatma Gandhi University,
PD Hills PO, Kottayam - 686560,
Kerala, India.

E-mail: nevinkg@gmail.com

ABSTRACT: The present study was designed to evaluate the *in-vitro* antioxidant effects and potential benefits of using coconut kernel extract in preventing the proliferation of human breast cancer cell line, MCF-7. Defatted extract from coconut kernel (CK_f) was evaluated for its radical scavenging, prevention of human LDL oxidation, cytotoxic, apoptotic and nuclear effects on MCF-7 cells. Intracellular production of reactive oxygen species (ROS) and superoxide radical was studied. It was showed that CK_f significantly scavenged free radicals and prevented LDL oxidation. MCF-7 cell viability was found to be decreased in a dose-dependent manner following treatment with CK_f. AO-EB and DAPI staining confirmed that treatment with CK_f induced nuclear damage and apoptosis as distinctly visualized in cells. Our observations suggest that nuclear/cellular damage and poor mitochondrial membrane potential (MMP) appears to be an indirect consequence of the excessive generation of reactive oxygen species (ROS) and superoxide radicals.

INTRODUCTION: Purpose and Rationale:

Several studies have been done on the potential anti-proliferative, and apoptotic effects of plant/fruit derived extracts on different cell lines ^{1, 2}. Since, plants have proved to be an excellent source of bioactive molecules, many of these compounds have been developed as drugs or precursors for the design and synthesis of potent drugs.

Modulation in the levels of antioxidant enzymes, generation of reactive oxygen species, differential gene expression and activation of cell death pathways might account for a few of the mechanisms responsible for the anti-proliferative/anti-tumor effects of such extracts ³. Coconut (*Cocos nucifera*) has been used in traditional folklore medicine to treat several disorders and as a functional food in different countries.

Apart from the fatty acids, coconut kernel contains proteins, fibers and several other biologically active minor components ⁴. It is in this light that we have attempted to investigate the effects of minor components containing coconut kernel extract, whose anti-tumor potency remains unreported till date on breast cancer cell lines *in-vitro*.

	<p>DOI: 10.13040/IJPSR.0975-8232.IJP.5(3).186-93</p>
	<p>The article can be accessed online on www.ijpjournals.com</p>
<p>DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5(3).186-93</p>	

MATERIALS AND METHODS:

Defatting and Extraction: Samples of Dwarf x Tall (DxT) variety of coconuts were collected from an authorized nursery (Agrofert Private Nursery, Kottayam, Kerala, India). Kernel from coconut was removed and defatted with petroleum ether (60-80) °C using a Soxhlet apparatus. The residue obtained after defatting was dried, weighed and exhaustively extracted using 80% methanol to obtain the extract (CK_f).

Radical Scavenging Assays: The stable 1, 1-diphenyl-2-picrylhydrazyl radical (DPPH) was used for the determination of free radical- the scavenging activity of CK_f⁵. Different concentrations of the sample were added at an equal volume to the methanolic solution of DPPH (0.135 mM). After 30 min dark incubation at room temperature, the absorbance was recorded at 517 nm. 2, 2'- azino - bis (3 - ethylbenzothiazoline - 6- sulphonic acid) (ABTS) assay was done according to the procedure described by Re *et al.*,⁶ with slight modifications. Test samples of 20 µL were allowed to react with 180 µL of ABTS⁺ solution for 12 min in the dark at room temperature. The free radical inhibitory activity was measured as the percentage scavenging calculated by the following formula:

$$\text{Scavenging activity (\%)} = \left[\frac{(Ac - At)}{Ac} \times 100 \right]$$

Where, Ac is the absorbance value of the control and At is the absorbance value of the test samples.

In-vitro LDL Oxidation Studies: Human LDL was isolated from the plasma of healthy individuals by precipitation method using sodium citrate buffer (0.064M pH 5.04) containing 50,000 U/l heparin developed by Wieland & Seidel. Approval for the study was obtained from the Institutional Human Ethics Committee (IHEC), School of Biosciences, Mahatma Gandhi University and informed consent was given by all volunteers⁷. Concentration of LDL was expressed as LDL protein measured by the method described by Lowry *et al.*,⁸ to evaluate the ability of CK_f to inhibit copper (Cu) mediated LDL oxidation, samples were pre-treated with various concentration of CK_f and LDL oxidation was initiated by adding Copper Sulfate at a final concentration of 10 µM/mL. After 24 h of incubation, oxidation was terminated by the addition of 33mM EDTA. TBARS, in each tube, was estimated in each tubes by the method of

Ohkawa *et al.*,⁹ Electrophoretic mobility of CK_f treated LDL to detect LDL fragmentation was done using SDS PAGE (10% gradient)¹⁰.

Cytotoxicity, Morphology and Nuclear Damage: MCF-7 cells were procured from National Centre for Cell Science (NCCS), Pune India. The cells were grown and maintained in a humidified incubator at 37 °C under 5% CO₂ atmosphere in Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 10% foetal bovine serum (FBS). The effects of varying concentrations of CK_f on cytotoxicity, nuclear integrity and morphology were visualized by staining methods. MCF-7 cells were seeded in 96 well plates at a density of 3 × 10⁴ cells / well, and the cells were treated with CK_f at varying concentrations between 12.5 - 50 mg/ml for 24 h at 37 °C with 5% CO₂. Following incubation, the wells were washed with PBS and treated with MTT for evaluating cytotoxicity. AO/EB and DAPI stains were used for studying morphology, apoptosis, nuclear shape, size, fragmentation. Cells were examined at 20x magnification using an Olympus inverted fluorescence microscope with a FITC filter^{11, 12, 13}.

Intracellular ROS and Superoxide Radical Measurement: Generation of excessive intracellular ROS following treatment of cells with extract was examined using a fluorescent probe, 2', 7'-Dichlorofluorescein Diacetate (DCFH-DA) and superoxide radicals by MitoSOX assay^{14, 15}. Briefly, 96 well plates seeded with MCF-7 cells were allowed to become confluent and incubated with different concentrations of CK_f (35 and 50 mg/ml) for 24 h. Treated cells were then washed with phosphate buffered saline (PBS) and incubated with fresh DCFH-DA (100 µM) and MitoSOX Red (5 µM) in PBS for 30 min at 37 °C in 5% CO₂. Fluorescent images of cells were taken using an inverted microscope (Olympus 1 × 51).

Mitochondrial Membrane Potential Analysis: Rhodamine 123 Staining: To determine the effect of CK_f on the mitochondrial electric potential across the inner membrane, rhodamine 123 (R-123), a lipophilic cationic indicator was used to label mitochondria¹⁶. MCF-7 cells were seeded into 96-well culture plates at 1 × 10⁴ cells/well and incubated for 24 h at 37 °C and 5% CO₂. After 24h, cells were treated with different doses of CK_f (35

and 50 mg/ml) for 24 h. After 24 h, R-123 solution was added into the media containing cells and incubated for 20 min. Subsequently, the cells were washed with PBS twice, and the cellular images were taken at 525 nm using the fluorescence microscope (Olympus 1 × 51).

Statistical Analyses: All the data are expressed as the mean ± standard deviation of three independent experiments. Statistical comparison was performed using SPSS 19 software *via* a one-way analysis of variance (ANOVA) followed by Duncan's multiple range test (DMRT). P values of less than 0.05 (P < 0.05) were considered significant.

RESULTS:

CK_f Shows *in-vitro* Free Radical Scavenging Activity and Prevents LDL Oxidation: Methanolic extract (CK_f) showed significant DPPH radical scavenging effect in a concentration-dependent manner (IC₅₀ 1.26 µg/µL). ABTS assay showed greater antiradical and antioxidant activities of CK_f. The extract also showed significant cation radical scavenging effect in a concentration-dependent manner (IC₅₀ 0.27 µg/µL) **Fig. 1.** Pre-treatment of human LDL with CK_f and subsequent exposure to CuSO₄ showed lower levels

of TBARS, indicative of lesser oxidative damage compared to oxidized LDL control **Fig. 2A.**

Based on the electrophoretic mobility assay, it was found that LDL pre-treated with a higher concentration of CK_f showed a profound protective effect on apoproteins of LDL **Fig. 2B.** Oxidation of LDL by Cu²⁺ is considered a more aggressive system that transmits through Cu²⁺ binding, causing continuous free radical formation with reduction of lipoprotein antioxidants¹⁷.




FIG. 1: IN-VITRO RADICAL SCAVENGING ACTIVITY OF CK_f: DPPH AND ABTS ASSAY. Values are expressed as mean ± SD of three independent experiments.





FIG. 2: EFFECT OF CK_f ON PREVENTING Cu²⁺ INDUCED LDL OXIDATION IN-VITRO. Results are expressed as the mean ± SD of three separate experiments. A: TBARS production in native, oxidized and CK_f pre-treated LDL, B: Electrophoretic mobility of oxidized and treated LDL. *Significant compared to control (p < 0.05), [§]Significant compared to oxidized LDL (P < 0.05).

Cytotoxic Effect of CK_f towards MCF-7: Changes in Nucleus and Morphology: It was observed that cytotoxicity of CK_f towards MCF-7 cells was concentration- dependent. The percentage of cell death was observed to increase at a steady pace till about 25 mg/ml after which the cytotoxicity appears to be relatively absolute in all

concentrations >25 mg/ml **Fig. 3.** **Fig. 4** distinctly shows CK_f treated cells with brightly fluorescing Ethidium bromide in the nucleus and periphery which is indicative of loss of membrane integrity and early apoptosis. **Fig. 5** shows that CK_f treated cells have different size, nuclear fragmentation patterns, and chromatin condensation compared to

non-treated control cells. This indicates that nuclear damage precedes cell death following treatment with CK_f. Plant-derived compounds have been

reported to cause cell death in cancer cell lines by inducing DNA double-stranded breaks as a result of apoptosis¹⁸.




FIG. 3: CYTOTOXIC EFFECT OF CK_f ON BREAST CANCER CELL LINES (MCF-7) *IN-VITRO*. Values are expressed as mean \pm SD of three separate experiments, A: MTT assay to evaluate the cytotoxic effect of different concentrations of CK_f against MCF-7 cells. B: light microscopic images of MCF-7 cells treated with varying concentrations of the extract (24h) (Magnification 20x)

CK_f Reduced the Mitochondrial Membrane Potential with Increased ROS and Superoxide Levels in MCF-7 Cells: The results showed that CK_f treatment induced a dose-dependent decrease **Fig. 5** in the mitochondrial membrane potential compared to the control cells which showed a very

high fluorescence. The levels of intracellular ROS have risen tremendously in cells treated with the CK_f. Meanwhile, the mitochondrial superoxide production in CK_f treated MCF-7 cells were also found to be increased significantly compared to control cells.





FIG. 4: ANALYSIS OF CHANGES IN CELL MORPHOLOGY AND VIABILITY OF MCF-7 CELLS TREATED WITH DIFFERENT CONCENTRATIONS OF CK_f. The images were taken using a fluorescent microscope and are the representation of three independent experiments (Magnification 20x).




FIG. 5: ANALYSIS OF NUCLEAR DAMAGE AND CHROMATIN CHANGES IN MCF-7 CELLS TREATED WITH DIFFERENT CONCENTRATIONS OF CK_f. The images were taken using a fluorescent microscope and are the representation of three independent experiments (Magnification 20x).




FIG. 6: ANALYSIS ON THE EFFECT OF CK_f TREATMENT ON THE ROS, SUPEROXIDE PRODUCTION AND MITOCHONDRIAL MEMBRANE POTENTIAL IN MCF-7 CELLS. The images were taken using a fluorescent microscope and are the representation of three independent experiments (Magnification 20x).

DISCUSSION: The present study evaluated the antioxidant and anti-cancer activity of the defatted coconut kernel extract in a breast cancer cell line *in-vitro*. It was found that the extract showed significant *in-vitro* scavenging activity of free radicals and prevented human LDL oxidation. Previous studies in our laboratory showed that the coconut kernel extract contained polyphenolic and flavonoid class of compounds. (Unpublished data) The free radical scavenging and antioxidant activities of phenolic compounds/flavonoids are dependent upon the arrangement of functional groups about the nuclear structure. Both the number and configuration of H-donating hydroxyl groups are the main structural features influencing the antioxidant capacity of phenolics¹⁹. CK_f was found to increase the production of free radicals

and decreased the mitochondrial membrane potential of MCF-7 cells. This observation indicates that drug-induced- ROS/superoxide generation in cells might contribute to the cytotoxic effect of the extract on breast cancer cell lines *in-vitro*. In our study, we demonstrated the effects of the methanol extract derived from the coconut kernel on the growth and viability of the breast cancer cell line, MCF-7.

These observations led us to hypothesize that the extract induced oxidative stress plays a significant role in decreasing cell viability. It is well known that the basal levels of intracellular ROS are significantly higher in cancer cells as compared to the normal cells which are what drives the growth, progression and other metabolic activities of these

cells. However, the pro-carcinogenic effect of this basal ROS level can propel a cell towards apoptosis once a threshold ROS limit has been exceeded^{20, 21}. From our observations as depicted in **Fig. 4**, the cells treated with varying concentrations of extract displayed a higher threshold fluorescence in contrast to the control cells which hints at the pro-oxidant potential of the CK_f in general. Generally, ROS generated as a part of regular cell metabolism is balanced by the intracellular glutathione and thioredoxin antioxidant systems.

Mitochondria are also a significant contributor to cell ROS through the flow of electrons in the electron transport chain. Considering the prospects of oxidative stress-induced cell death also prompts us to consider the possibility of compromised cellular antioxidant defenses which could be the cause of cytotoxicity²². Our hypothesis is strengthened through the nuclear-stained images as shown in **Fig. 4** which indicates the formation of apoptotic bodies and nuclear fragmentation in concentrations as low as 12.5 mg/mL which is indicative of late-stage apoptosis. The mitochondrial specific ROS probe labeled MCF-7 cells **Fig. 6** showed an increased fluorescence at concentrations >35mg/ml indicating the loss of mitochondrial membrane integrity at higher doses of the extract. Compromised mitochondrial membrane potential caused by excessive stress can initiate the signaling cascade of apoptosis which could possibly account for the CK_f induced cytotoxicity.

CONCLUSION: These results propel us to envision the benefits involved in identifying the bioactive components of CK_f, which exert a cytotoxic effect on breast cancer cell line, MCF-7, *in-vitro*. Our experimental results indicate that increased oxidative stress, compromised mitochondrial membrane integrity, and nuclear damage are major events in CK_f induced cytotoxicity. However, the mechanism of CK_f action *in-vitro* can only be explained after evaluating changes in gene-expression levels, and nuclear damage are major events in CK_f.

ACKNOWLEDGEMENT: This study was supported by the grant from the Coconut Development Board (CDB), Ministry of Agriculture, Government of India (Grant No.1345/

2011). The authors are grateful to School of Biosciences, Mahatma Gandhi University and Department of Biotechnology, Government of India for the excellent research facilities supported through the prestigious DBT - MSUB - IPLS (BUILDER) programme (No.BT/PR4800/INF/22/152/2012).

CONFLICT OF INTEREST: The authors declare no potential conflicts of interest concerning the research and publication of this article.

REFERENCES:

1. Lin HH, Huang HP, Huang CC, Chen JH and Wang CJ: Hibiscus polyphenol-rich extract induces apoptosis in human gastric carcinoma cells *via* p53 phosphorylation and p38 MAPK/FasL cascade pathway. *Molecular Carcinogenesis* 2005; 43(2): 86-99. 10.1002/mc.20103.
2. Seeram NP, Adams LS, Henning SM, Niu Y, Zhang Y, Nair MG and Heber D: *In-vitro* antiproliferative, apoptotic and antioxidant activities of punicalagin, ellagic acid and a total pomegranate tannin extract are enhanced in combination with other polyphenols as found in pomegranate juice. *Journal of Nutritional Biochemistry* 2005; 16(6): 360-367. <https://doi.org/10.1016/j.jnutbio.2005.01.006>.
3. Vadodkar AS, Suman S, Lakshmanaswamy R and Damodaran C: Chemoprevention of breast cancer by dietary compounds. *Anticancer Agents in Medicinal Chemistry* 2012; 12(10): 1185-1202. <https://doi.org/10.2174/187152012803833008>
4. Renjith RS, Chikku, AM and Rajamohan T: Cytoprotective, anti-hyperglycemic and phytochemical properties of *Cocos nucifera* (L.) inflorescence. *Asian Pacific Journal of Tropical Medicine* 2013; 6(10): 804-810. [https://doi.org/10.1016/S1995-7645\(13\)60142-X](https://doi.org/10.1016/S1995-7645(13)60142-X).
5. Koleva II, Van Beek TA, Linssen JPH, de Groot A and Evstatieva LN: Screening of plant extracts for antioxidant activity: A comparative study on three testing methods. *Phytochemical Analysis* 2002; 13(1): 8-17. 10.1002/pca.611.
6. Re R, Pellegrini N, Proteggente A, Pannala A, Yang M and Rice-Evans C: Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biology and Medicine* 1999; 26(9): 1231-1237.
7. Wieland H and Seidel D: A simple specific method for precipitation of low-density lipoproteins. *Journal of Lipid Research* 1983; 24(7): 904-909.
8. Lowry OH, Rosebrough NJ, Farr AL and Randall RJ: Protein measurement with the Folin phenol reagent. *Journal of Biological Chemistry* 1951; 193(1): 265-275.
9. Ohkhawa H, Ohishi N and Yagi K: Assay of lipid peroxides in animal tissues by the thiobarbituric acid reaction. *Analytical Biochemistry* 1979; 95(2): 351-358. [https://doi.org/10.1016/0003-2697\(79\)90738-3](https://doi.org/10.1016/0003-2697(79)90738-3).
10. Sakai J, Hoshino A, Takahashi S, Miura Y, Ishii H, Suzuki H, Kawarabayasi Y and Yamamoto T: Structure, chromosome location, and expression of the human very low-density lipoprotein receptor gene. *Journal of Biological Chemistry* 1994; 269(3): 2173-2182.
11. Marks DC, Belov L, Davey MW, Davey RA and Kidman AD: The MTT cell viability assay for cytotoxicity testing in multidrug-resistant human leukemic cells. *Leukemia*

- Research 1992; 16(12): 1165-1173. [https://doi.org/10.1016/0145-2126\(92\)90114-M](https://doi.org/10.1016/0145-2126(92)90114-M).
12. Baskić D, Popović S, Ristić P and Arsenijević NN: Analysis of cycloheximide-induced apoptosis in human leukocytes: fluorescence microscopy using annexin V/ propidium iodide versus acridine orange / ethidium bromide. *Cell Biology International* 2006; 30(11): 924-932. <https://doi.org/10.1016/j.cellbi.2006.06.016>.
 13. Chazotte B: Mounting live cells onto microscope slides. *Cold Spring Harbour Protocol* 2011; 1: 1-4. doi: 10.1101/pdb.prot5554.
 14. Warleta F, Quesada CS, Campos M, Allouche Y, Beltrán G and Gaforio JJ: Hydroxytyrosol protects against oxidative DNA damage in human breast cells. *Nutrients* 2011; 3(10): 839-857. doi:10.3390/nu3100839
 15. Robinson KM, Janes MS, Pehar M, Monette JS, Ross MF, Hagen TM, Murphy MP and Beckman JS: Selective fluorescent imaging of superoxide *in-vivo* using ethidium-based probes. *Proceedings of National Academy of Sciences USA* 2006; 103(41): 15038-15043. doi: 10.1073/pnas.0601945103.
 16. Nobili S, Lippi D, Witort E, Donnini M, Bausi L, Mini E and Capaccioli S: Natural compounds for cancer treatment and prevention. *Pharmacological Research* 2009; 59(6): 365-378. <https://doi.org/10.1016/j.phrs.2009.01.017>.
 17. Costa-Mugica A, Batista- Gonzalez AE, Mondejar D, Soto-López Y, Brito-Navarro V, Vázquez AM, Brömme D, Zaldívar-Muñoz C, Novoa AV, de Oliveira e Silva AM and Mancini-Filho J: Inhibition of LDL-oxidation and antioxidant properties related to polyphenol content of hydrophilic fractions from seaweed *Halimeda incrassata* (Ellis) Lamourou. *Brazilian Journal of Pharmaceutical Sciences* 2012; 48(1): 31-37. <http://dx.doi.org/10.1590/S1984-82502012000100004>.
 18. Matkar SS, Wrischnik LA and Hellmann-Blumberg U: Sanguinarine causes DNA damage and p53-independent cell death in human colon cancer cell lines. *Chemico-Biological Interactions* 2008; 172(1): 63-71. <https://doi.org/10.1016/j.cbi.2007.12.006>
 19. Hepsibha B, Thendral S, Satiya C, Saravana Babu V, Premalakhmi and Sekar T: *In-vitro* studies on antioxidant and free radical scavenging activities of *Azima tetraacantha* Lam leaf extracts. *Indian Journal of Science and Technology* 2010; 3(5): 571-577.
 20. Martin-Cordero C, Leon-Gonzalez AJ, Calderon-Montano JM, Burgos-Moron E and Lopez-Lazaro M: Pro-oxidant natural products as anticancer agents. *Current Drug Targets* 2012; 13(8): 1006-1028. <https://doi.org/10.2174/138945012802009044>
 21. Schumacker PT: Reactive oxygen species in cancer cells: live by the sword, die by the sword. *Cancer Cell* 2006; 10(3): 175-176. <https://doi.org/10.1016/j.ccr.2006.08.015>.
 22. Pelicano H, Carney D and Huang P: ROS stress in cancer cells and therapeutic implications. *Drug Resistance Updates* 2004; 7(2): 97-110. <https://doi.org/10.1016/j.drup.2004.01.004>.

How to cite this article:

Dhanyakrishnan R, Sunithav MC, Prakashkumar B, Sandya S and Nevin KG: Evaluation of *in-vitro* antioxidant and anti-proliferative effect of defatted extract of *Cocos nucifera* kernel on human breast cancer cell line. *Int J Pharmacognosy* 2018; 5(3): 186-93. doi link: [http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5\(3\).186-93](http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5(3).186-93).

This Journal licensed under a Creative Commons Attribution-Non-commercial-Share Alike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)