



Received on 11 January, 2018; received in revised form, 05 February, 2018; accepted, 13 February, 2018; published 01 June, 2018

BENEFICIAL ROLE OF CURCUMIN IN DISEASES TREATMENT

Seyyed Hossein Hassanpour ^{* 1}, Mohammad Amin Dehghani ², Seyyed Mozaffar Alipour ³, Seyyedeh Zeinab Karami ⁴ and Fatemeh Dehghani ⁵

Young Researchers and Elite Club ¹, Yasooj Branch, Islamic Azad University, Yasooj, Iran.

Department of Toxicology ², School of Pharmacy, Ahvaz Jundishapour University of Medical Sciences, Ahvaz, Iran.

Department of Environmental Health, School of Health ³, Yasouj University of Medical Sciences, Yasouj, Iran.

Department of Biology ⁴, School of Basic Sciences, Yasouj University, Yasouj, Iran.

Department of Genetics ⁵, Faculty of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran.

Keywords:

Curcumin, Cancer, Inflammation, Apoptosis

Correspondence to Author: Seyyed Hossein Hassanpour

Young Researchers and Elite Club,
Yasooj Branch, Islamic Azad
University, Yasooj, Iran.

E-mail: Dr.hossein1366@yahoo.com

ABSTRACT: Curcumin regulates cellular signaling pathway related to cancer growth, inflammation and apoptosis and due to biological activities can be target DNA, RNA and enzymes. In addition, curcumin has inhibitory effect against MARK, P53, β -catenin and AKT. It can be consider as a promising agent to treat many diseases but it is important that should be acceptably carried out cellular targeting with curcumin. Here, we reviewed beneficial effects of curcumin in treatment of diseases. Our review study was confirmed that curcumin has potential effect in diseases treatment but there is a dire need for increase of its bioavailability in order to promotion of its efficacy. We suggest that should be design the methods for increase of cellular targeting by curcumin in further studies.

INTRODUCTION: Curcumin is considered as a phenolic compound with therapeutic properties including anti-inflammation, antioxidant and anticancer ¹. In previous studies, it has been found that it is safe compound for the body following its administration in patients with cancer and healthy individuals. In fact, its therapeutic effects is more than its side effects ^{2, 3}. However, despite having such a beneficial effect on human biological, it has not very high bioavailability in human due to poor absorption and fast metabolism ^{4, 5}.

In animal and human, major amount of curcumin is excreted as urine and feces and is absorbed only a small portion of curcumin that combines with glucuronide and sulfate in order to create its conjugated forms in liver. These events lead to reduction of plasma concentrations of free curcumin ^{2, 5-9}.

In addition, curcumin is an insoluble compound in water at neutral and basic conditions due to production of phenolate that leads to poor availability in aqueous systems ^{10, 11}. Using nanoparticles can increase cellular uptake and stability of curcumin so that solid lipid nanoparticles promote distribution and stability of curcumin in aqueous solutions that lead to improvement of the anticancer effect of curcumin ¹². In addition, silica nanoparticle can be a suitable delivery system for curcumin because it has been

	DOI: 10.13040/IJPSR.0975-8232.IJP.5(6).327-30
	Article can be accessed online on: www.ijppjournal.com
DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5(6).327-330	

demonstrated that silica nanoparticle increase anti-tumor properties of curcumin¹³. In this study, we reviewed the beneficial effect of curcumin in order to treatment of diseases.

2. Review Method: Here, we searched paper related to role of curcumin treatment of diseases with keywords such as curcumin and cancer, curcumin and Parkinson's diseases, curcumin and Alzheimer disease, curcumin and histone acetyl transferase (HAT), curcumin and diabetes in databases include web of science, PubMed and Scopus. Then, the papers were fully read and summarized here.

3. Curcumin Beneficial Effects in Diseases

Treatment: According to a study Yallapu *et al.*, 2014, PLGA or poly (lactic-co-glycolic acid) combined with curcumin leads to a reduction of tumor cells growth (prostate cancer). Poly (lactic-co-glycolic acid) combined with curcumin had pivotal role in inhibition of beta-catenin expression in tumor cells. In addition, it led to inhibition of anti-apoptotic proteins (Mcl-1, Bcl-xL) through suppression of STAT3 and AKT phosphorylation. Down-regulation of miR21 and up-regulation of miR-205 also were showed after treatment with poly (lactic-co-glycolic acid) combined with curcumin. Given that this effects poly (lactic-co-glycolic acid) combined with curcumin can be consider as a promising agent to treat cancer due to good delivery into tumor cells¹⁴. Thiolated chitosan nanoparticle can promote anticancer effect of curcumin because it increases curcumin bioavailability. In addition, treatment of mice with colon cancer by curcumin loaded with thiolated chitosan nanoparticle led to dramatically reduction of cancer cells¹⁵.

Kumar *et al.*, 2014 examined that treatment of ovarian cancer cells (SKOV-3) with curcumin-loaded polymer of 2-hydroxy methyl methacrylate resulted in decrease of cancer cells by arresting of cell cycle in G0/G1 and induction of apoptosis¹⁶. It has been reported that curcumin-combined silica nanoparticle conjugated with hyaluronic acid results in cytotoxic effects on human colon carcinoma (colo-205) that is due to appropriate delivery of curcumin in to cancer cells through silica nanoparticle conjugated with hyaluronic acid¹⁷. Curcumin-loaded PLGA-PEG nanoparticle has

anti-tumor effects against breast cancer (cell line MCF-7) due to obvious reduction of cancer cells followed by treatment with curcumin-loaded PLGA-PEG nanoparticle¹⁸. The effect of polysorbate 80-coated cerasomes in order to deliver curcumin in mice with Parkinson's disease was confirmed promotion of cross of blood-brain barrier and subsequently improvement of mice with Parkinson's disease¹⁹. Curcumin and piperine coated with glycerol monooleate easily pass the blood-brain barrier and prevent formation of alpha synuclein oligomers. In addition, curcumin and piperine coated with glycerol monooleate reduce effect of rotenone as a Parkinson's disease inducer. Curcumin and piperine coated with glycerol monooleate is an important factor in reduction of oxidative stress, apoptosis and autophagy pathway and ultimately protect dopaminergic neurons against damage²⁰.

A research group was found that amine-functionalized mesoporous silica nanoparticle could be a good delivery system for curcumin in order to abrogation of Parkinson's disease complications because administration of curcumin-loaded amine-functionalized mesoporous silica nanoparticle leads to reduction of alpha synuclein accumulation²¹. Mathew *et al.*, 2012 designed a water-soluble nanoparticle (PLGA nanoparticle conjugated with Tet-1 peptide) for transferring of curcumin. Based our their study, curcumin loaded-PLGA nanoparticle conjugated with Tet-1 peptide prevents the accumulation of amyloid beta and had antioxidant effect and not cytotoxic effect therefore it can be very effective in the treatment of Alzheimer's disease²². It has been showed improvement of memory and reduction of amyloid plaques in mice with Alzheimer's disease after treatment with highly stabilized curcumin nanoparticle according to study conducted by Kin Cheng *et al.*, 2013. Meanwhile, the presence of high level of curcumin in plasma indicated enhancement of its pharmacokinetic characteristic using the nanoparticle²³.

Curcumin-loaded PLGA nanoparticle is effective in neural stem cell differentiation at rat's hippocampal and sub ventricular. In fact, injection of this compound into hippocampal leads to neurogenesis induction through expression of genes related to cell growth and proliferation (reelin, nestin and

Pax6) and differentiation of neurons, such as neurogenin, neuro D1, neuregulin, neuroligin and Stat3. These events ultimately result in improvement of learning and memory due to destruction of beta amyloid²⁴. Curcumin-coated magnetic nanoparticle has diagnostic value for Alzheimer's disease because it is a non-invasive diagnostic procedure by MRI and reduces PET disadvantages²⁵. Curcumin - loaded PLGA nanoparticle is a good agent against Alzheimer's disease due to it protects neurons against oxidative stress by up-regulation of expression of genes like APOE, APOJ, TRX, GLRX REST that indicate its high uptake in neurons²⁶. Curcumin inhibits histone acetyl transferase, a main reason of cardiac hypertrophy and heart failure, and prevents hypertrophy^{27, 28}. Furthermore, curcumin protects heart against damage caused by ischemia/reperfusion^{29, 30}.

In cells, curcumin leads to proteasome-dependent degradation of histone acetyl transferase. This event is main reason for inhibition of histone acetyl transferase by curcumin. In fact, curcumin is a selective inhibitor of histone acetyl transferase²⁸. By the mechanism, curcumin prevents excessive increase of ratio heart weight to body weight³¹. Retinopathy occurs in diabetes due to inability of body for elimination of free radicals and pro-inflammatory molecules such as NF- κ B, VEGF, and IL1 β . Curcumin prevents progression of diabetes retinopathy by inhibition of free radicals formation and reduction of inflammation³².

CONCLUSION: The use of herbal medicines is increasing in recent years. Thus, efforts to review and revive herbal medicines are reasonable so that to understand their effect mechanism lead to achieve new strategies to treat many diseases. The curcumin is most important component of turmeric. However, its low bioavailability has limited its healing properties. Today, it has been proposed several mechanisms to increase its bioavailability that can be improve its absorption. For example, the nanoparticles can significantly increase curcumin effectiveness. In this review study, we showed that to apply strategies for promotion of curcumin bioavailability lead to increase of its efficacy in treatment of diseases therefore we suggest that design further studies based on this view.

ACKNOWLEDGEMENT: Us acknowledgement and gratefulness at the beginning and at last is to god who gave us the gift of the mind. The authors thank Young Researchers and Elite Club, Yasooj Branch, Islamic Azad University due to cooperation in this study.

CONFLICT OF INTEREST: The authors declare that there is no conflict of interest regarding this study.

FINANCIAL SUPPORT AND SPONSORSHIP: This study was supported by the authors named in this article alone.

CONTRIBUTION OF AUTHORS: This work was done by the authors named in this article and all liabilities pertaining to claims relating to the content of this article was borne by the authors named in this article.

ETHICAL APPROVAL: This research does not contain any studies with human participants or animals and was performed by the authors alone.

REFERENCES:

1. Vareed SK, Kakarala M, Ruffin MT, Crowell JA, Normolle DP, Djuric Z, *et al.*: Pharmacokinetics of curcumin conjugate metabolites in healthy human subjects. *Cancer Epidemiology and Prevention Biomarkers* 2008; 17(6): 1411-7.
2. Cheng AL, Hsu CH, Lin JK, Hsu MM, Ho YF, Shen TS, *et al.*: Phase I clinical trial of curcumin, a chemopreventive agent, in patients with high-risk or pre-malignant lesions. *Anticancer Res.* 2001; 21(4B): 2895-900.
3. Lao CD, Ruffin MT, Normolle D, Heath DD, Murray SI, Bailey JM, *et al.*: Dose escalation of a curcuminoid formulation. *BMC complementary and alternative medicine.* 2006; 6(1): 10.
4. Gutierrez VO, Campos ML, Arcaro CA, Assis RP, Baldan-Cimatti HM, Peccinini RG, *et al.*: Curcumin pharmacokinetic and pharmacodynamic evidences in streptozotocin-diabetic rats support the antidiabetic activity to be *via* metabolite (s). *Evidence-Based Complementary and Alternative Medicine* 2015.
5. Jäger R, Lowery RP, Calvanese AV, Joy JM, Purpura M and Wilson JM: Comparative absorption of curcumin formulations. *Nutrition journal* 2014; 13(1): 11.
6. Pan MH, Huang TM and Lin JK: Biotransformation of curcumin through reduction and glucuronidation in mice. *Drug metabolism and disposition* 1999; 27(4): 486-94.
7. Ireson C, Orr S, Jones DJ, Verschöyle R, Lim CK, Luo JL, *et al.*: Characterization of metabolites of the chemopreventive agent curcumin in human and rat hepatocytes and in the rat *in vivo*, and evaluation of their ability to inhibit phorbol ester-induced prostaglandin E2 production. *Cancer research* 2001; 61(3): 1058-64.
8. Phase I. Clinical Trial of Oral Curcumin: Biomarkers of Systemic Activity and Compliance Sharma. Ricky A.6847-54.

9. Dulbecco P and Savarino V: Therapeutic potential of curcumin in digestive diseases. *World J Gastroenterol.* 2013; 19(48): 9256-70.
10. Gantait A, Barman T and Mukherjee PK: Validated method for estimation of curcumin in turmeric powder 2011.
11. Modasiya M, Patel A and Patel V: International Journal of Pharmacy and Life Sciences. *Int J of Pharm and Life Sci (IJPLS)* 2012; 3(3):1490-7.
12. Sun J, Bi C, Chan HM, Sun S, Zhang Q and Zheng Y: Curcumin-loaded solid lipid nanoparticles have prolonged *in vitro* antitumour activity, cellular uptake and improved *in vivo* bioavailability. *Colloids and surfaces b: biointerfaces* 2013; 111: 367-75.
13. Gangwar RK, Tomar GB, Dhumale VA, Zinjarde S, Sharma RB and Datar S: Curcumin conjugated silica nanoparticles for improving bioavailability and its anticancer applications. *Journal of agricultural and food chemistry* 2013; 61(40): 9632-7.
14. Yallapu MM, Khan S, Maher DM, Ebeling MC, Sundram V, Chauhan N, et al.: Anti-cancer activity of curcumin loaded nanoparticles in prostate cancer. *Biomaterials* 2014; 35(30): 8635-48.
15. Anitha A, Deepa N, Chennazhi K, Lakshmanan VK and Jayakumar R: Combinatorial anticancer effects of curcumin and 5-fluorouracil loaded thiolated chitosan nanoparticles towards colon cancer treatment. *Biochimica et Biophysica Acta (BBA)-General Subjects* 2014; 1840(9): 2730-43.
16. Kumar SSD, Surianarayanan M, Vijayaraghavan R, Mandal AB and MacFarlane D: Curcumin loaded poly (2-hydroxyethyl methacrylate) nanoparticles from gelled ionic liquid–*In vitro* cytotoxicity and anti-cancer activity in SKOV-3 cells. *European Journal of Pharmaceutical Sciences* 2014; 51: 34-44.
17. Singh SP, Sharma M and Gupta PK: Cytotoxicity of curcumin silica nanoparticle complexes conjugated with hyaluronic acid on colon cancer cells. *International journal of biological macromolecules* 2015; 74: 162-70.
18. Mirakabad TFS, Akbarzadeh A, Milani M, Zarghami N, Taheri-Anganeh M, Zeighamian V, et al.: A Comparison between the cytotoxic effects of pure curcumin and curcumin-loaded PLGA-PEG nanoparticles on the MCF-7 human breast cancer cell line. *Artificial cells, nanomedicine, and biotechnology* 2016; 44(1): 423-30.
19. Nisi Z, Huang X, Zhifei D, Zheng H and Fei Y: Localized delivery of curcumin-loaded nanoparticles to the brain for treatment of Parkinson's disease using focused ultrasound. *Ultrasonics Symposium (IUS), 2016 IEEE International;* 2016.
20. Kundu P, Das M, Tripathy K and Sahoo SK: Delivery of Dual Drug Loaded Lipid Based Nanoparticles across the Blood–Brain Barrier Impart Enhanced Neuroprotection in a Rotenone Induced Mouse Model of Parkinson's Disease. *ACS Chemical Neuroscience* 2016.
21. Taebnia N, Morshedi D, Yaghmaei S, Aliakbari F, Rahimi F and Arpanaei A: Curcumin-Loaded Amine-Functionalized Mesoporous Silica Nanoparticles Inhibit α -Synuclein Fibrillation and Reduce Its Cytotoxicity-Associated Effects. *Langmuir* 2016; 32(50): 13394-402.
22. Mathew A, Fukuda T, Nagaoka Y, Hasumura T, Morimoto H, Yoshida Y, et al.: Curcumin loaded-PLGA nanoparticles conjugated with Tet-1 peptide for potential use in Alzheimer's disease. *PLoS one* 2012; 7(3): e32616.
23. Cheng KK, Yeung CF, Ho SW, Chow SF, Chow AH and Baum L: Highly stabilized curcumin nanoparticles tested in an *in vitro* blood–brain barrier model and in Alzheimer's disease Tg2576 mice. *The AAPS journal* 2013; 15(2): 324-36.
24. Tiwari SK, Agarwal S, Seth B, Yadav A, Nair S, Bhatnagar P, et al.: Curcumin-loaded nanoparticles potentially induce adult neurogenesis and reverse cognitive deficits in Alzheimer's disease model *via* canonical Wnt/ β -catenin pathway. *ACS nano* 2013; 8(1): 76-103.
25. Cheng KK, Chan PS, Fan S, Kwan SM, Yeung KL, Wang Y-XJ, et al.: Curcumin-conjugated magnetic nanoparticles for detecting amyloid plaques in Alzheimer's disease mice using magnetic resonance imaging (MRI). *Biomaterials* 2015; 44: 155-72.
26. Djiokeng Paka G, Doggui S, Zaghmi A, Safar R, Dao L, Reisch A, et al.: Neuronal uptake and neuroprotective properties of curcumin-loaded nanoparticles on SK-N-SH cell line: role of poly (lactide-co-glycolide) polymeric matrix composition. *Molecular pharmaceutics* 2015; 13(2): 391-403.
27. Li HL, Liu C, De Couto G, Ouzounian M, Sun M, Wang A-B, et al.: Curcumin prevents and reverses murine cardiac hypertrophy. *The Journal of clinical investigation* 2008; 118(3): 879-93.
28. Marcu MG, Jung YJ, Lee S, Chung EJ, Lee MJ, Trepel J, et al.: Curcumin is an inhibitor of p300 histone acetyltransferase. *Medicinal chemistry* 2006; 2(2): 169-74.
29. Kapakos G, Youreva V and Srivastava AK: Cardiovascular protection by curcumin: molecular aspects 2012.
30. Yeh CH, Chen TP, Wu YC, Lin YM and Lin PJ: Inhibition of NF κ B Activation with Curcumin Attenuates Plasma Inflammatory Cytokines Surge and Cardiomyocytic Apoptosis Following Cardiac Ischemia / Reperfusion. *Journal of Surgical Research* 2005; 125(1): 109-16.
31. Mito S, Watanabe K, Harima M, Thandavarayan RA, Veeraveedu PT, Sukumaran V, et al.: Curcumin ameliorates cardiac inflammation in rats with autoimmune myocarditis. *Biological and Pharmaceutical Bulletin* 2011; 34(7): 974-9.
32. Kowluru RA and Kanwar M: Effects of curcumin on retinal oxidative stress and inflammation in diabetes. *Nutrition and metabolism* 2007; 4(1): 8.

How to cite this article:

Hassanpour SH, Dehghani MA, Alipour SM, Karami SZ and Dehghani F: Beneficial role of curcumin in diseases treatment. *Int J Pharmacognosy* 2018; 5(6): 327-30. doi link: [http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5\(6\).327-30](http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5(6).327-30).

This Journal licensed under a Creative Commons Attribution-Non-commercial-Share Alike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)