



Received on 05 February, 2018; received in revised form, 21 March, 2018; accepted, 30 March, 2018; published 01 June, 2018

PHARMCOLOGICAL ACTIVITY OF *THUJA ORIENTALIS* LINN.

Sandip Kshirsagar, Sunanda Malode* and Santosh Bansode

Department of Pharmacognosy, Kasturi Shikshan Sanstha's College of Pharmacy, Shikrapur, Pune - 412208, Maharashtra, India.

Keywords:

Thuja orientalis, Morpankhi, Thujone, Pharmacological activity

Correspondence to Author:

Sunanda Malode

Department of Pharmacognosy, Kasturi Shikshan Sanstha's College of Pharmacy, Shikrapur, Pune - 412208, Maharashtra, India.

Email: sunandamalode93@gmail.com

ABSTRACT: *Thuja orientalis* (Commonly- Morpankhi, Family- Cupressaceae) is an evergreen, monoecious trees or shrubs used in various forms of traditional medicines and homeopathy in various ways. In traditional practices, Thuja is used for treatment of bronchial catarrh, enuresis, cystitis, psoriasis, uterine carcinomas, amenorrhea and rheumatism. Recent re-researches in different parts of the world have shown that *Thuja orientalis* and its active component thujone have the great potential against a various health problems. *Thuja orientalis* preparations can be efficiently used against microbial/worm infection. It can be used as antioxidant, anticancer and anti-inflammatory agent. Instead of these effects, it can be also used as insecticidal, molluscicidal and nematicidal activity against different pests. The present review highlights the some important bio-logical properties of *Thuja orientalis*.

INTRODUCTION: *Thuja orientalis* is a common ornamental evergreen tree that is originally native to Northwest China belonging to family Cupressaceae. It is highly aromatic and resinous shrub that widely cultivated in gardens located in temperate and semi-temperate areas. *Thuja orientalis* is hardy, large evergreen shrub or small to medium sized-tree rarely exceeding 20 m in nature. It has a dense, pyramidal shape, but often exhibits a more open and spreading form. It prefers moist, well-drained soil and full sun. The bark is gray with brown highlights and has thin but deep furrows. The bark has a rugged charm about it, especially on large mature specimens.

Younger bark is a reddish-brown colour and exfoliates in long, thin strips. It is endemic to north-western China. It is also now naturalized as an introduced species elsewhere in Asia: eastward to Korea and Japan; southward to northern India; and westward to northern Iran. The common name 'arbor-vitae' is from Latin, 'Tree of life', and is based on its association with long life and vitality in Buddhist thought in China. This is probably based on the tree's unchanging evergreen nature in the cold dry climate of northwest China and its longevity; some of the larger specimens planted around Buddhist temples in China are said to be in excess of 1,000 years old.

Although generally accepted as the only member of its genus, it has been included by botanists in the different classification. In older texts, Platycladus was often included in Thuja, but it is only distantly related to that genus. Differences from Thuja include distinct cones, wingless seeds, and almost scentless foliage.

	QUICK RESPONSE CODE DOI: 10.13040/IJPSR.0975-8232.IJP.5(6).331-336
	Article can be accessed online on: www.ijpjournal.com
DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5(6).331-336	

Later, genus name has been changed as *Biota orientalis* and latest as *Platyclusus orientalis*. But, it has been still known and sold as *Thuja orientalis* in market.

Plant Profile:

Taxonomical Classification:

Kingdom: Plantae
 Division: Coniferophyta
 Class: Pinopsida
 Order: Pinales
 Family: Cupressaceae
 Genus: Platyclusus

Vernacular Names:

Hindi: Morpankhi.
 Marathi: Thuja, Morphanakhi
 Chinese: Baishu, Xiangbai, Cebai, Bianbai
 Italian: Tuiaorientale
 Japanese: Konotegashiwa
 Spanish: Uya De La China
 Trade Name: Chinese arborvitae
 English: Thuja orientalis, Biota, Tree of life, Book leaf pine.
 French: Thuya oriental, ThuyaD'orient, Thuya De Chine
 German: Morgelaendischer Lebensbaum,
 Chinesische Thuja, Lebensbaum,

Species:

Thuja koraiensis
Thuja occidentalis L.
Thuja plicata
Thuja standishii
Thuja sutchuenensis

Synonyms:

Biota orientalis,
Cupressus pendula Thunb.
Platyclusus stricta Spach
Thuja acuta Moench
Thuja decora Salisb.
Thuja orientalis L.

Morphology Plant:

Plant Type: Shrub or small
 Origin: Northwest China
 Height: 15 to 20 feet
 Spread: 10 to 15 feet
 Plant habit: Spreading

Leaves: Leaves light green or yellow-green, becoming brown after 3 or 4, seasons, persistent, scale-like, in opposite-decussate pairs.

Flowers: Not ornamentally important, Monoecious

Fruit: Fleshy, glaucous blue-green colour when developing, Woody horn-like cone when mature, Medium-brown colour.

Bark: Shredding, Brown to gray-brown colour.



FIG. 1: TREE, SEED AND BARK

Chemical Constituents: The main constituents of essential oils mono- and ses-quiterpenes including carbohydrates, phenols, alcohols, ethers, aldehydes and ketones are responsible for the bio-logical activity of aromatic and medicinal plants.

Thuja orientalis leaves contain rhodoxanthin, amentoflavone, hinokiflavone, quercetin, myricetin,

carotene, xan-thophylls and ascorbic acid. The seed yields fatty oil having the following composition palmitic 5.28, stearic, 7.3; C₁₈ unsaturated acids, 1829 (Linolenic, 44.6%); and C₂₀ unsaturated acids, 6.10%.

The Heartwood contains aroma- dendrin, taxifolin, widdrene, cedrol, thujopsadiene, dehydro - α -

curcumene, β -isobiotol and curcumenether. The fruit oil contained α -Pinene (52.4%), 3-Carene (14.2%), α -Cedrol (6.5%) and phellandrene (5.1%), the leaf oil contained α -pinene (21.9%), α -cedrol

(20.3%), 3-carene (10.5%) and limonene (7.2%) as the main components. Thujone is a ketone and a monoterpene that occurs naturally in two diastereomeric forms: α -Thujone and β -Thujone.




FIG. 2: CHEMICAL CONSTITUENTS OF *THUJA ORIENTALIS*

Uses: It has been used in the treatment of various diseases as a folk medicine such as rheumatism, gout, dermatitis, and diarrhea. The plant has been exhibited extensively biological activities including anti-epileptic, anti-inflammatory, hair growth-promoting, antiviral, anti-allergic, anti-bacterial, antioxidant, and antifungal activities. Traditionally it is used in the treatment of cough. It is also used as a medicinal plant in various forms of traditional medicines like folk medicine, homeopathy, etc. for treatment of bronchial catarrh, enuresis, cystitis, psoriasis, uterine carcinomas, amenorrhea and rheumatism. The root bark is used in the treatment of burns and scalds. The stems are used in the treatment of coughs, colds, dysentery and parasitic skin-diseases.

The leaves are antibacterial, antipyretic, antitussive, astringent, diuretic, refrigerant and stomachic. It can be also used as Insecticidal, molluscicidal and

nematicidal activity against different pests. *Thuja orientalis* is used internally in the treatment of coughs, haemorrhages, excessive menstruation, bronchitis, asthma, skin infections, mumps, bacterial dysentery, arthritic pains and premature blandness. The leaves are antipyretic, astringent, diuretic, emmenagogue, emollient, expectorant, refrigerant and stomachic. Their use is said to improve the growth of hair.

The seed is aperients, lenitive and sedative. It is used internally in the treatment of palpitations, insomnia, nervous disorder and constipation in the elderly. The stems are used in the treatment of coughs, cold, dysentery, rheumatism and parasitic skin diseases. Thujone was a weak inhibitor of acyl-CoA: lysophosphatidylcholine acyl-transferase activity in mouse brain synaptosomes compared to psycho-active cannabinoids.

Biological Activity:

Antipyretic Activity: In the present study the alcoholic extract of the leaf of *Thuja orientalis* were studied for their antipyretic activity by Brewer's yeast-induced pyrexia in rats. It was observed that the alcoholic extract produced significant antipyretic activity ($p < 0.05$). The extract showed marked antipyretic activity in a dose dependent manner. The antipyretic activity of the alcoholic extract was evaluated based on Brewer's yeast-induced pyrexia in rats.

Pyrexia was induced by subcutaneous injection of 10 ml/kg of 15% w/v Brewer's yeast suspension below the nape of the neck. The rectal temperature of each rat was measured at time, 0 hr, using a telethermometer and before injection of the yeast, at 18 hr following yeast injection, the different groups were treated with alcoholic extract (200 and 400 mg/kg), and standard drug, paracetamol (150 mg/kg). Tween 80 (1% v/v) was used as suspending agent. The rectal temperature was then recorded over a period of 6 h².

Anthelmintic Activity: The activity of drugs that expel parasitic worms (helminths) and other internal parasites from the body by either stunning or killing them and without causing significant damage to the host is called anthelmintic activity. The aerial parts of *Thuja orientalis* showed anthelmintic activity. The anthelmintic activity was determined for the methanolic extract of a real part of *Thuja orientalis*, which displayed significant anthelmintic properties at higher concentration. The extract revealed anthelmintic activity in a dose dependent manner which is effective against parasitic infections of human¹¹.

Hair Growth-promoting Activity: *Thuja orientalis* leaves are traditionally used to promote hair growth. Studies proved that the leaves contained strong 5-reductase inhibitors, which may be associated with hair growth promoting activity. Another study showed the decoction of the leaves can promote hair growth by inducing anagen in telogenic C57BL/6 N mice. In decoction-treated mice, an increase in the number and size of hair follicles can be observed, which served as a piece of evidence for the induction of anagen phases.

Using the immunohistochemical analysis, an earlier induction of β -catenin and Sonic hedgehog (Shh)

proteins in the decoction-treated group also can be seen, compared to the control or 1% minoxidil-treated group. Therefore, the hair growth-promoting activity can be explained by the inhibition of 5-reductase and induction of β -catenin and Shhproteins⁴.

Antimicrobial Potential of Various Fractions of *Thuja orientalis*: The present study was conducted to evaluate the antimicrobial activity of various extracts (*viz.* methanol, acetone and ethyl acetate) of *Thuja orientalis* and its preliminary phytochemical screening. *In-vitro* antimicrobial activity was tested by agar well diffusion assay against human pathogenic microbes which are Gram positive (*viz.* *S. aureus* and *B. subtilis*), Gram negative bacteria (*viz.* *P. aeruginosa*, *A. faecalis* and *K. pneumoniae*) and fungi (*A. flavus* and *A. niger*). Methanol and acetone extract of leaf exhibited maximum activity against *B. subtilis* with zones of inhibition of 20.33 mm and 17.83 mm, respectively.

All the extracts of stem were found to be most effective against *P. aeruginosa*. *S. aureus* was found to be sensitive to leaf extracts prepared in methanol, acetone and ethyl acetate with 13.66, 14.03 and 15.00 mm zone of inhibition, respectively. *A. flavus* and *A. niger*, both were found to be sensitive towards the methanol extract of leaf with inhibition zones of 15.50 and 16.00 mm, respectively compare to fungicides (*viz.* ketoconazole and fluconazole). Methanol extract of leaf and ethyl acetate extract of stem exhibit marked antimicrobial activity against *B. subtilis* and *P. aeruginosa*, which was comparable to standard antibiotics (that is, penicillin, ampicillin, tetracycline and streptomycin). It was also observed that leaf extracts were more effective as compared to stem extracts of the plant. Thus, *Thuja orientalis* could be considered as potential source of natural antimicrobials used for the treatment of bacterial and fungal infections⁷.

Anti-Inflammatory and Analgesic Activity: This study was conducted for evaluating a natural source to treat inflammation and pain, to avoid the severe side effects of currently used agents for these ailments. *Thuja orientalis* (More Pankh) is commonly used for the treatment of pain and inflammatory disorders in traditional medicine.

Carrageenan induced inflammatory model, acetic acid induced writhing test and hot plate methods were used to evaluate anti-inflammatory, peripheral and central analgesic properties of aqueous methanolic extract of *Thuja orientalis* fruit (To-Cr) in albino rats. Completely randomized design (CRD) was constructed for the study and one way ANOVA was applied to compare means. The results showed that TO-Cr has significant anti-inflammatory and analgesic properties⁹.

Antibacterial and Antioxidant Activity: In the present study, leaves of *Thuja orientalis* were powdered and extracted by soxhlet extractor in two solvent systems that is, (E1) ethyl acetate: chloroform: ethanol (40: 30: 30) and (E2) methanol: distilled water (70:30). This study conferred the screening of phytochemical constituents, antioxidant activity and antibacterial activity of crude E1 and E2 extract and its fractions. Antioxidant activity was carried out by 2,2-diphenyl-1-picrylhydrazyl(DPPH) assay.

The results indicate that E2 extract (70% methanolic extract) had the highest antioxidant effect (85.25% inhibition) at 100 µg/ml concentration and the crude extracts (E1 and E2 extract) showed significant ($P \leq 0.05$) inhibitory activity against both gram positive and gram negative organisms. It was active against *Staphylococcus aureus*, *Bacillus subtilis*, *Escherichia coli* and *Agrobacterium tumefaciens*.

The minimum inhibitory concentrations (MICs) of E1 extract ranged from 0.40 to 0.85 mg/ml and E2 extract 0.55 to 1.15 mg/ml. The highest antibacterial potentiality was exhibited by E2 extract. The fractions also exhibited antimicrobial activity against all the selected microorganisms. The study revealed that *Thuja orientalis* is a promising phytomedicine for antioxidant and antibacterial activity¹⁰.

Anticancer Activity: Strong 5 α -reductase inhibitor are extracted and fractionated from *T. orientalis* and purified as diterpenes in isolated form⁷. The inhibitors are used either on their own or as active ingredients of therapeutics in the treatment of diseases caused by the over activity of 5 α -reductase or the hyper-secretion of androgens, such as male baldness, androgenetic alopecia,

hirsutism, acne, prostatomegaly and cancer of the prostate reported that the hepato-protective activities and antioxidant activity of *Thuja occidentalis* Linn.^{7,8}

Anti-proliferative and apoptosis - inducing properties of thujone-rich fraction (TRF) separated from *Thuja occidentalis*. Their possible anti-cancer potentials have been noted in the malignant melanoma cellline A375⁶. Concluded that apolysaccharide, or long-chain sugar molecule, derived from *Thuja* leaves extract decreased the inflammation caused by cancer. It also prevented the cancer from metastasizing, or spreading throughout the body¹¹.

Antiviral Activity: *In-vitro*, the antiviral activities of the plant extracts were investigated using the cell-based screening. Plant extracts of *Thuja orientalis* were shown to induce a high cell viability rate after the infection with the influenza A/PR/8/34 virus. The antiviral activity of the plant extracts also increased as a function of the concentration of the extracts and these extracts significantly reduced the visible cytopathic effect caused by virus infections. Furthermore, the treatment with *Thuja orientalis* was shown to have a stronger inhibitory effect than that with *A. spathulifolius* or *P. thunbergii*. These results may suggest that *Thuja orientalis* has anti-influenza A/PR/8/34 activity⁵.

CONCLUSION: I concluded from my review that *Thuja orientalis* Linn. traditionally widely used medicinal plant. *Thuja orientalis* is a common ornamental evergreen tree. It shows a wide range of biological activities that helps to utilize the medicinal benefits of this plant. However, here remains an immense scope for further exploration of this plant and needs the attention of scientists to exploit the full potential activities of this plant.

ACKNOWLEDGEMENT: Nil

CONFLICT OF INTEREST: Nil

REFERENCES:

1. Sharma PC, Yelne MB and Dennis TJ: Database on medicinal Plants used in Ayurveda. Central Council for Research in Ayurveda and Siddha, Department of ISM and H, Ministry of Health and Family Welfare (Govt. of India), New Delhi, Vol. 2.

2. Jaiswal A, Kumar A, Mishra D and Kasula M: Review pharmacological activity of *Platycladus orientalis*. International Research Journal of Pharmacy ISSN 2230 8407.
3. Shan MQ, Shang J and Ding AW: *Platycladus orientalis* Leaves: A systemic review on botany, phytochemistry and pharmacology. The American Journal of Chinese Medicine 42(3), 523-542.
4. Singh JD, Surekha SP and Kumar A: Antimicrobial potential of various fractions of *Thuja orientalis*. African Journal of Microbiology Research 2013; 7(25): 3179-3186.
5. Won, Ji-Na, Lee SY, Song D and Poo H: Antiviral activity of the plant extracts from *Thuja orientalis*, *Aster spathulifolius* and *Pinus thunbergii* against influenza virus A/PR/8/34, J. Microbiol. Biotechnol 2013; 23(1): 125-130.
6. US Patent 5773005 1998. Purified flavonoid and diterpene 5 α -reductase inhibitor from *Thuja orientalis* for androgen-related diseases.
7. Muhammad I, Mossa JS, Al Yahya MA, Ramadan AS and El-Seraly FL: Further antibacterial diterpenes from the bark and leaves of *Juniperus procera*. Hochst. ex. Endl. Phytotherapy Research, 1995; 9: 584-588.
8. Sunila ES, Hamsa TP and Kuttan G: Effect of *Thuja occidentalis* and its polysaccharide on cell-mediated immune responses and cytokine levels of metastatic tumor-bearing animals. Pharm Biol 2011; 49(10): 1965-1073.
9. Tanveer MZ, Javeed A and Ashraf M: Evaluation of Anti-inflammatory and analgesic potential of aqueous methanolic extract of *Thuja orientalis* in albino rats. The Journal of Animal and Plant Sciences 2015; 25(4): 1183-1186.
10. Jasuja ND, Sharma SK, Saxena R and Choudhary J: Antibacterial, antioxidant and phytochemical investigation of *Thuja orientalis* leaves. Journal of Medicinal Plants Research 2013; 7(25): 1886-1893.
11. Srivastavapriya, Kumar P and Singh KD: Biological properties of *Thuja orientalis* Linn. Advances in Life Sciences 2012; 2(2): 17-20.
12. Singh A, Varshneya C and Telang SR: *In-vitro* screening of *Thuja orientalis* leaf extract for anthelmintic activity against haemonchus contortus. Indian Journal of Small Ruminants 2005; 11(1): 98-100.

How to cite this article:

Kshirsagar S, Malode S and Bansode S: Pharmacological activity of *Thuja orientalis* Linn. Int J Pharmacognosy 2018; 5(6): 331-36. doi link: [http://dx.doi.org/10.13040/IJP.SR.0975-8232.IJP.5\(6\).331-36](http://dx.doi.org/10.13040/IJP.SR.0975-8232.IJP.5(6).331-36).

This Journal licensed under a Creative Commons Attribution-Non-commercial-Share Alike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)