



Received on 16 February, 2018; received in revised form, 25 March, 2018; accepted, 30 March, 2018; published 01 June, 2018

## PHARMACOGNOSTIC AND PHYTOCHEMICAL EVALUATION OF *NYCTANTHES ARBOR-TRISTIS* STEM

Bhuwan Chandra Joshi <sup>\*1</sup>, Nilesh Chauhan <sup>1</sup>, Sukanya <sup>2</sup> and Sushmita Uniyal <sup>3</sup>

Department of Pharmacognosy <sup>1</sup>, Sardar Bhagwan Singh Post Graduate Institute of Biomedical Sciences and Research, Balawala, Dehradun - 248001, Uttarakhand, India.

Department of Pharmacy <sup>2</sup>, Central University of Rajasthan Bandarsindri, Kishangarh - 305817, Rajasthan, India.

Gyani Inder Singh Institute of Professional Studies <sup>3</sup>, Dehradun - 248003, Uttarakhand, India.

### Keywords:

*Nyctanthes arbor-tristis*, Pharmacognosy, Fluorescence analysis

### Correspondence to Author:

**Bhuwan Chandra Joshi**

Assistant Professor,  
School of Pharmaceutical Sciences,  
Sardar Bhagwan Singh Post Graduate  
Institute of Biomedical Sciences and  
Research, Balawala, Dehradun -  
248001, Uttarakhand, India.


**E-mail:** bhuwan.joshi000@gmail.com

**ABSTRACT: Objective:** To study detailed pharmacognostic characters of the stem of *Nyctanthes arbor-tristis* (Oleaceae), along with their physico-chemical parameters, fluorescence analysis and phytochemical screening. **Methods:** The pharmacognostic characters were determined in terms of macroscopy, microscopy, powder microscopy, physicochemical analysis, fluorescence analysis and preliminary phytochemical investigation of plant stem. **Results:** The microscopic study shows the general characteristic of stem. Physico-chemical investigation shows the total ash, acid insoluble ash, water soluble ash were  $8.69 \pm 0.17\%$  w/w,  $0.21 \pm 0.11\%$  w/w, and  $3.92 \pm 0.05\%$  w/w respectively. Phytochemical analysis revealed the presence of various phytochemical groups like alkaloids, glycosides, steroids, phenolic, tannins constituents. **Conclusion:** It can be concluded that the established pharmacognostic profile of *Nyctanthes arbor-tristis* stem will be helpful in developing pharmacopoeial standards for correct identification and quality control

**INTRODUCTION:** India has a rich heritage of traditional medicine and traditional health care systems that have been flourishing for many centuries. Now a days, prevalent use of traditional medicines in the developed countries is trending and it has become more popular throughout the world. Traditionally, important large shrub of tropical and subtropical regions of the world have been used to counteract disease <sup>1</sup>.

*Nyctanthes arbour - tristis* Linn. (Oleaceae) commonly known as Parijat or Harsinghar, a hardy large shrub or small tree up to 5 - 10 m in height, widely occurring in outer Himalayan ranges from Kashmir to Nepal and throughout India up to 1000 - 1500 m altitude <sup>2</sup>.

The name 'Nyctanthes' has been coined from two Greek words 'Nykhta' means night and 'anthos' means flower. It is also planted in gardens due to its highly fragrant flowers <sup>3</sup>. It is shrub or small tree, with drooping branches and quadrangular branchlets. Leaves are opposite, ovate, acute or acuminate, entire slight cuneate. Flowers are small, 3 - 7 in head, arranged in trichotomous cymes, delightfully fragrant, sessile, slender, and hairy; corolla glabrous, orange colored and lobes are



white<sup>4, 5</sup>. Fruits are a capsules of 1 - 2 m in diameter, long and broad, compressed, 2 celled separating into 2 flat one seeded carpels, reticular veined and glabrous<sup>6</sup>. Different parts of this plant are used in folk-medicines<sup>7, 8</sup>. The leaves are bitter, useful in chronic fever<sup>9</sup>, malarial fever<sup>10</sup>, obstinate sciatica, constipation, haemorrhoids<sup>11</sup> and eczema<sup>12</sup>. The flowers are astringent, stomachic, and useful in dyspepsia, greyness of hair and baldness<sup>11</sup>. The plant elaborates different classes of organic compounds of medicinal importance including alkaloids, terpenes, steroids,  $\beta$ -sitosterol, glycosides, iridoid glycosides, arbortristiside-A, B, C, D, E<sup>13, 14, 15, 16, 17, 18</sup>. Different parts of this plant are used in Indian systems of medicine for various pharmacological actions like as anti-leishmaniasis, anti-viral, anti-fungal, anti bacterial, anti-pyretic, antihistaminic, anti-malarial, anti-oxidant, hepatoprotective, and anti-inflammatory activities<sup>19</sup>.

The literature survey and scientific data revealed that no systematic pharmacognostical parameter had been carried out on the stem of *Nyctanthes arbor-tristis* Linn. till date. Hence, the objective of present study is to evaluate various pharmacognostic parameters such as macroscopy, microscopy, physicochemical and phytochemical evaluations of the *Nyctanthes arbor-tristis* Linn. (stem).

## MATERIALS AND METHODS:

**Chemicals and Instruments:** Phloroglucinol, glycerin, hydrochloric acid, potassium hydroxide and all other chemicals used in the study were of analytical grade.

**Plant Material:** The stem of *Nyctanthes arbortristis* was collected from Dehradun, Uttarakhand, India in the month of August 2016 and authenticated by Dr. S. K. Srivastava, Botanical Survey of India, Northern regional centre, Dehradun, where a voucher specimen (specimen no. 116216) has been deposited.

**Macroscopic and Microscopic Analysis:** Macroscopic studies were done using simple microscope. The color, shape, size, taste and odour of stem were determined. Microscopic study of fresh stem was carried out by preparing thin transverse section and staining it with concentrated

hydrochloric acid: Phloroglucinol (1:1). Photographs sections were carefully taken. The dried stem were powdered and treated with 5% KOH solution followed by staining with concentrated hydrochloric acid - Phloroglucinol (1:1) for 5 minutes and mounted in 50% glycerine solution<sup>20, 21, 22</sup>.

**Physicochemical Analysis:** Physicochemical parameter such as ash values (total ash, acid insoluble ash) and extractive values (water soluble, alcohol soluble extractives) were determined using powdered drug. The moisture content was detected by loss on drying method<sup>23, 24, 25</sup>.

**Fluorescence Analysis:** For the fluorescence analysis of stem powder it was treated with various chemicals and was observed exclusively to different wavelengths of ultra violet (254 nm and 365 nm) and visible light for observing characteristic colour presentation<sup>22, 26, 27</sup>.

**Preliminary Phytochemical Screening:** Preliminary phytochemical screening was qualitatively tested for the presence of phytochemicals as per described standard methods<sup>22, 28, 29, 30</sup>.

## RESULTS:

**Macroscopic Characteristics:** Organoleptic characters of stem depicted that the stem was woody in nature having light grey to greenish in colour with characteristic odour and taste. The stem was 1 - 10 m tall, erect and branched.

### Microscopic Characteristics:

**Stem Microscopy:** Section of stem appears quadrangular and revealed the following tissues: The epidermis was single, layered consisting of rectangular cells with a thick continuous cuticle on the epidermis along with many multicellular hairs. Cortex was several cells deep below the four protruded comers while only few layers deep at the other places just beneath the epidermis. It was differentiated into collenchyma and parenchyma. Many intercellular spaces were present and the region extended upto the vascular tissue. Vascular bundles were present in the cortex, each protrudes bulb containing one. The pointed xylem end was faced towards outer side in each of the conical bundle. In other words, the vascular bundle was conjoint, collateral, open and exarch.

The microscopy revealed that the endodermis was not well developed. The pericycle was observed in the form of sclerenchymatous patches. Vascular system was composed of primary phloem, secondary phloem, cambium, primary xylem and secondary xylem. Crushed primary phloem was irregularly present in patches below the pericycle. The secondary phloem consisted of sieve tubes, companion cells, phloem parenchyma and was present in the form of a continuous ring. Cambium

was present as one to three cells thick continuous layer in between phloem and xylem. Secondary xylem was present just inner to the cambial ring and consisted of mainly thick walled woody parenchyma and fibres. Trachieds and vessels were also observed. Primary xylem was situated just near the pith in a way facing its protoxylem towards the centre. Pith was found to be thin walled and parenchymatous as shown in Fig. 1.




FIG. 1: TRANSVERSE SECTION OF STEM OF *NYCTANTHES ARBOR-TRISTIS* VIEWED AT 10x; A- WITHOUT STAINING REAGENT, B- WITH STAINING REAGENT

**Stem Powder Microscopy:** Microscopic observation of *Nyctanthes arbor-tristis* stem indicated the presence of parenchyma cells,

collenchyma cells, fiber, xylem vessels and calcium oxalate crystal as shown in Fig. 2.




FIG. 2: POWDERED CHARACTERISTICS OF THE STEM PARTS OF *NYCTANTHES ARBOR-TRISTIS* SHOWING A: COLLENCHYMA CELL (CC) B: PARENCHYMA CELLS (PC) C: CALCIUM OXALATE CRYSTALS (CO) D: XYLEM VESSELS (XV) E: FIBER

**Physicochemical Parameters:** Ash value of the drug gives idea about earthy matter or inorganic composition and other impurities present along with the drug. Various physicochemical parameters such as total ash, acid insoluble ash and water soluble ash of *Nyctanthes arbor-tristis* stem were found to be  $8.69 \pm 0.17$ ,  $0.21 \pm 0.11$  and  $3.92 \pm 0.05\%$  w/w, respectively. However,  $15.93 \pm 0.46\%$  w/w alcohol soluble and  $18.31 \pm 0.46\%$  w/w water soluble extractives were observed. The moisture content of stem powder was nearly  $5.49 \pm 0.02\%$  w/w **Table 1**.

**TABLE 1: PHYSICOCHEMICAL CONSTANTS FOR NYCTANTHES ARBOR-TRISTIS STEM**

S. no.	Physicochemical parameter	Values (% w/w)
1	Moisture content	$5.49 \pm 0.02$
2	Total ash	$8.69 \pm 0.17$
3	Acid insoluble ash	$0.21 \pm 0.11$
4	Water soluble ash	$3.92 \pm 0.05$
5	Alcohol soluble extractive	$15.93 \pm 0.46$
6	Water soluble extractives	$18.31 \pm 0.46$

**Fluorescence Analysis:** Fluorescence analysis of stem powder was carried out after treating it with several solvents and chemicals. Fluorescence was observed at 254 and 365 nm comparing its change of colour in visible light. The observations are presented in **Table 2**.

**TABLE 2: FLUORESCENCE ANALYSIS OF NYCTANTHES ARBOR-TRISTIS STEM**

Treatment	Visible light	Under UV light	
		Short wavelength (254 nm)	Long wavelength (365 nm)
Powder	Brown	Light brown	Green
Powder + Methanol	Brown	Light brown	Yellowish green
Powder + 70% ethanol	Brown	Light brown	Green
Powder + Pet. ether	Light brown	Light green	Green
Powder + 50% H <sub>2</sub> SO <sub>4</sub>	Brown	Greenish brown	Brownish
Powder + 50% HCl	Dark Brown	Green	Green black
Powder + 1N NaOH (aq.)	Light brown	Dark brown	Brownish black
Powder + 1N NaOH (alc.)	Light brown	Dark green	Greenish black
Powder + 50% HNO <sub>3</sub>	Brown	Light brown	Light green
Powder + 5% KOH	Brown	Purplish green	Dark purplish green
Powder + Ammonia	Brown	Green	Black
Powder + Picric acid	Yellowish brown	Green	Dark brown

**Preliminary Phytochemical Screening:** Preliminary phytochemical screening of *Nyctanthes arbor-tristis* is shown in **Table 3**.

**TABLE 3: PHYTOCHEMICAL SCREENING NYCTANTHES ARBOR-TRISTIS STEM**

S. no.	Class of constituents	PEE	CE	EAE	EE	AE
1	Amino acids	-	-	-	-	-
2	Proteins	-	-	-	-	-
3	Carbohydrates	-	-	-	+	+
4	Steroids	+	+	-	-	-
5	Triterpenoids	+	+	-	-	-
6	Alkaloids	-	+	-	+	+
7	Glycosides	-	+	-	+	+
8	Saponins	-	-	-	-	-
9	Flavonoids	-	-	+	+	-
10	Tannins	-	-	-	+	+
11	Phenolic compounds	-	+	+	+	+

PEE- Pet. ether extract, CE- Chloroform extract, EAE- Ethyl acetate extract, EE- Ethanol extract, AE- Aqueous extract, (-) Negative, (+) Positive

**DISCUSSION:** The wide use of herbal drugs in traditional medicines and herbal formulations, standardization that has been made to an important measure for ensuring and justifying quality, purity and authenticity of the crude drugs<sup>31</sup>. Standardization purpose, morphological and microscopic analysis is one of the simplest and cheapest methods to start with establishing the correct identification of the source materials<sup>32, 33</sup>. As there is no pharmacognostic work available on this medicinally potent plant, the present work was undertaken to lay down the standards which could be useful for establishing its authenticity. Organoleptic and microscopic studies are useful identifying parameters for authentication of the drug<sup>34, 35</sup>.

Physicochemical studies acts as a reliable method for detecting adulteration. Physicochemical constants like total ash, acid insoluble ash, and water soluble ash can serve as a valuable source of information which is usually helpful in evaluation of purity and quality of a crude drug. The earthy matter or inorganic composition and other impurities which are present along with the drug are determined by the ash values. Acid-insoluble ash usually indicates the contamination with silicon material like earth and sand. Water-soluble ash was used for the estimation of the amount of inorganic elements<sup>36</sup>. The extractive values give an idea about the chemical constitution of the drug<sup>33</sup>.

Fluorescence analysis is an alternative rapid useful method for identification of authentic samples and recognizing adulterants. In this analysis, the crude drugs may be examined as such, in solution or as extracts and in powdered form<sup>37</sup>. Fluorescence characteristics enable the identification and differentiation of plant materials from their adulterants, when physical and chemical methods are scarce. Various chemical constituents present in the plant material exhibit fluorescence on absorbing light. Fluorescence is shown by some of the constituents even in the visible range in daylight. The ultraviolet light produces fluorescence in many natural products viz. alkaloids like berberine, which does not show fluorescence in the daylight. With the aid of different reagents, the non fluorescent substances can easily be transformed into their fluorescent derivatives or decomposition products<sup>33</sup>.

Phytochemical evaluation and chemo-profiling are useful for the quality assessment of plant materials. Phytochemical compounds in the plant are known to have various therapeutic importances. For instance saponins, terpenoids, flavonoids, tannins, steroids and alkaloids have anti-inflammatory effects. Glycosides, flavonoids, tannins and alkaloids have hypoglycemic activities<sup>38, 39</sup>. Flavonoids possess the hepatoprotective, and antioxidant activities<sup>40</sup>. The saponins have hypocholesterolemic and antidiabetic properties<sup>41</sup>. In the animal studies terpenoids tend to decrease blood sugar level. Steroids as well as triterpenoids exhibit the analgesic properties<sup>42</sup>. The steroids and saponins are responsible for CNS activities<sup>43</sup>.

**CONCLUSION:** The present study was focused on establishing pharmacognostic standards for the identification and authentication of the *Nyctanthes arbor-tristis*. Therefore, the outcomes of the above findings will serve as a promising source for laying down pharmacopoeial standards for the future studies and research.

**ACKNOWLEDGEMENT:** We express our sincere thanks to Shri S. P. Singh Honorable Chairman, Sardar Bhagwan Singh Post Graduate Institute of Biomedical Sciences and Research, Balawala India for providing the facilities.

**CONFLICT OF INTEREST:** We declare that we have no conflict of interest.

## REFERENCES:

1. Mukherjee PK: Quality Control of Herbal Drugs- An Approach to Evaluation of Botanicals Business Horizons, New Delhi, 1st Edition 2002.
2. Nadkarni KM: Indian Materia Medica, Bombay: Bombay Popular Prakashan Pvt. Ltd., 1982; 3: 1.
3. Kirtikar KR and Basu BD: Indian Medicinal Plants-Vol. IV, Sri Satguru Publications, Delhi, 2000
4. Saxena RS, Gupta B and Lata S: Tranquillizing, antihistaminic and purgative activity of *Nyctanthes arbor-tristis* leaf extract, J Ethanopharmacol 2002; 81: 321-325.
5. Ratnasooriya WD, Jayakody JRAC and Hettiarachchi ADI: Sedative effects of hot flower infusion of *Nyctanthes arbor-tristis* on rats, Pharm Biol 2005; 43: 140-146.
6. Bhosale AV, Abhyankar MM, Pawar SJ, Shoeb K and Patil N: *Nyctanthes Arbor-tristis*. A Pharmacognostic Review, Research J. Pharmacognosy and Phytochemistry 2009; 1(2): 91-97.
7. Gacche R and Dhole N: Antioxidant and possible anti-inflammatory potential of selected medicinal plants prescribed in the Indian traditional system of medicine. Pharmaceutical biology. 2006; 44(5): 389-95.
8. Khare CP: Indian medicinal plants: an illustrated dictionary: Springer Science & Business Media, 2008.
9. Gogoi R and Borthakur SK: Notes on herbal recipes of Bodo tribe in Kamrup district, Assam. Ethnobotany 2001; 13: 15-23.
10. Changkija S: Applied Ethnobotany: A Case Study among the Kharias of Central India. Asian Folklore Studies. 1999; 57(2):397-9.
11. Warriar PK and Nambiar V: Indian medicinal plants: a compendium of 500 species: Orient Blackswan; 1993.
12. Bhatt DC, Mitaliya KD and Mehta SK: Observations on ethnoveterinary herbal practices in Gujarat, Ethnobotany 2001, 13: 91-95.
13. Kapoor LD, Survey of Indian plants for saponins, alkaloids and flavonoids, II, Lloydia 1971; 34: 94.
14. Agrawal J and Pal A. *Nyctanthes arbor-tristis* Linn-A critical ethnopharmacological review. Journal of ethnopharmacology. 2013; 146(3):645-58.
15. Khatune NA, Islam ME, Abdur Rahman MA, Mosaddik MA and Haque ME. *In-vivo* cytotoxic evaluation of a new benzofuran derivative isolated from *Nyctanthes arbor-tristis* L. on ehrlich ascite carcinoma cells (EAC) in mice, J Med Sci. 2003; 3(2): 169-73.
16. Mathuram V and Kundu AB: A reinvestigation of the structures of arbortristiosides A and B from *Nyctanthes arbor-tristis*, J *Nyctanthes arbor-tristis* Prod 1991; 54(1): 257-60.
17. Stuppner H, Muller EP, Mathuram V and Kundu AB: Iridoid Glycosides from *Nyctanthes arbor-tristis*, Phytochemistry 1993; 32(2): 375-78.
18. Gupta P, Bajpai SK, Chandra K, Singh KL and Tandon JS: Antiviral profile of *Nyctanthes arbor-tristis* L. against encephalitis causing viruses, Indian J Exp Biol 2005; 43(12): 1156-1160.
19. Jain PK and Pandey A: The wonder of Ayurvedic medicine - *Nyctanthes arbor-tristis*. Int J Herb Med 2016; 4(4): 9-17.
20. Shah BN and Seth AK: Pharmacognostic studies of the *Lagenaria siceraria* (Molina) Standley. International Journal of PharmTech Research. 2010; 2(1):121-4.
21. Kokate CK: Practical Pharmacognosy, Vallabh Prakashan, New Delhi, 4<sup>th</sup> Edition 1994.
22. Khandelwal KR: Practical Pharmacognosy, Nirali publication, Pune, 18<sup>th</sup> Edition, 2007.

23. Government of India. The Ayurvedic pharmacopoeia of India. New Delhi: Ministry of Health and Family Welfare, Department of Indian Systems of Medicines and Homeopathy, 1<sup>st</sup> Edition 1996.
24. WHO: Quality control for medicinal plant material. New Delhi: AITBS Publishers; 1998.
25. WHO: Quality control methods for medicinal plant material. Geneva: WHO, 1992.
26. Kokoski CJ, Kokoski RJ and Slama FJ: Fluorescence of Powdered Vegetable Drugs under Ultraviolet Radiation, J. Amer. Pharma. Asso 1958; 10: 715-717.
27. Joshi BC and Uniyal S: Establishment of quality control protocols and antioxidant activity of *Urtica dioica* L. Journal of Conventional Knowledge and Holistic Health 2017; 1 (1): 1-6.
28. Farnsworth NR: Biological and phytochemical screening of plants, J. Pharm. Sci. 1966; 55: 225-276.
29. Harborne A: Phytochemical methods a guide to modern techniques of plant analysis: springer science & business media, 3rd Edition 1998.
30. Trease GE and Evans WC: Pharmacognosy. Harcourt brace & Co. Asia, Pvt. Ltd., W.B. Saunders Company Ltd., 15<sup>th</sup> Edition 2002.
31. Verma S and Singh SP: Current and future status of herbal medicines, Vet World 2008; 1(11): 347-350.
32. Gopalkrishnan B, Chiranjeev R. Quality Standardization of Flowers of *Nyctanthes arbor-tristis* Linn. International Journal of Pharmacognosy and Phytochemical Research 2017; 9(10): 1314-1317.
33. Sheikh N, Desai T and Patel R: Pharmacognostic Evaluation of *Epilobium hirsutum* Linn. Pharmacognosy Journal 2016; 8 (3): 226-29.
34. Zhao Z, Liang Z and Guo P: Macroscopic identification of Chinese medicinal materials: traditional experiences and modern understanding. J Ethnopharmacol 2011; 131: 556-61.
35. Vilash V, Suja SR, Latha PG and Rajasekharan S: Physicochemical Evaluation and Pharmacognostical Standardization of *Pellionia heyneana* Wedd. Leaf Pharmacognosy Journal. 2016; 8(6). 551-56.
36. Bello H, Mohammed Z and Katsayal U: Pharmacognostic evaluation of the root *Cassia sieberiana* DC: A promising ethnomedicinal plant. Journal of Pharmacognosy and Phytochemistry 2016; 5(3):270-75.
37. Hudson N, Baker A and Reynolds D: Fluorescence analysis of dissolved organic matter in natural, waste and polluted waters-a review. River Research and Applications 2007; 23 (6): 631-49.
38. Sharma B, Balomajumder C and Roy P: Hypoglycemic and hypolipidemic effects of flavonoid rich extract from *Eugenia jambolana* seeds on streptozotocin induced diabetic rats, Food Chem Toxicol 2008; 46(7): 2376-2383.
39. Orhan I, Kupeli E, Sener B and Yesilada E: Appraisal of anti-inflammatory potential of the clubmoss, *Lycopodium clavatum* L., J. Ethnopharmacol 2007; 109: 146-150.
40. Kumar RS, Manivannan R and Balasubramaniam A: Antioxidant and hepatoprotective activity of ethanol extract of *Indigofera trita* Linn. on CCl<sub>4</sub> induced hepatotoxicity in rats, Journal of Pharmacology and Toxicology 2008; 3: 344-350.
41. Rupasinghe HP, Jackson CJ, Poysa V, Di Berado C, Bewley JD and Jenkinson J: Soyasapogenol A and B distribution in Soybean (*Glycine max* L. Merr) in relation to seed physiology, genetic variability and growing location, J Agric Food Chem, 2003; 51: 5888-5894.
42. Srivastava M, Kumar A and Pal M: Phytochemical investigation on *Jatropha curcas* seed cake, Int J Pharm Life Sci, 2010; 1(6): 357-362.
43. Salna KP, Sreejith K, Uthiralingam M, Prince MA, John Milton MC and Fleming AT: Comparative study of phytochemicals investigation of *Andrographis paniculata* and *Murraya koenigii*, IJPPS 2011; 3(3): 291-292.

**How to cite this article:**

Joshi BC, Chauhan N, Sukanya and Uniyal S: Pharmacognostic and phytochemical evaluation of *Nyctanthes arbor-tristis* stem. Int J Pharmacognosy 2018; 5(6): 376-81. doi link: [http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5\(6\).376-81](http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5(6).376-81).

This Journal licensed under a Creative Commons Attribution-Non-commercial-Share Alike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)