



Received on 20 April, 2018; received in revised form, 08 June, 2018; accepted, 13 June, 2018; published 01 August, 2018

PHYTOCHEMICAL STUDY AND EVALUATION OF WOUND HEALING POTENTIAL OF CRUDE LEAVE EXTRACTS OF *OROXYLUM INDICUM* (L.) VENT IN WISTAR RATS

Bhabani Kumari Samarath and Sangram K. Panda *

Jeypore College of Pharmacy, Rondapalli, Jeypore - 764002, Koraput, Odisha, India.

Keywords:

Oroxylum indicum,
Incision wound, Excision wound
and Wound contraction

Correspondence to Author: Sangram K. Panda

Jeypore College of Pharmacy,
Rondapalli, Jeypore - 764002,
Koraput, Odisha, India.


E-mail: sangrampanda04@gmail.com

ABSTRACT: The present study was carried out to evaluate the wound healing potential of different extracts of *Oroxylum indicum* leave in two different types of wound models in albino rats viz., incision and excision, and compare the effects with povidine iodine ointment which is used as standard drug. Among all the extract, the ethanolic extract at a dose of (250 mg/kg) possesses significant increase in wound contraction and formation of scar in excision wound model. The extract showed significant increase in the breaking strength of resutured incision wound as compared to control group ($p < 0.05$). The result of the present study indicate that ethanolic extract of *O. indicum* has more significant wound healing property than the other two extracts in excision and incision wound model.

INTRODUCTION: Herbal medicine has become an integral part of standard healthcare. The World health organization estimated that 80% of people worldwide rely on herbal medicines for some aspect of their primary healthcare¹. Wound healing can be defined as a complex dynamic process that results in the results in the restoration of anatomic continuity and function. It is a finely orchestrated and overlapping sequence of events involving vascular response phase/hemostasis, inflammation, proliferation, maturation and remodeling^{2,3}. Many Ayurvedic herbal plants have a very important role in the treatment of wound. Plants are more potent healers because they promote the repair mechanisms in the natural way. So the present study undergoes to evaluate the wound healing activity of *Oroxylum indicum*.

The tribal areas of Baipariguda, Koraput (District) of Eastern Orissa, due to its unique varieties geographical and climatic factors has had a rich variety of medicinal plant. *Oroxylum indicum* (family: Bignoniaceae) also known as Phana Phana (Oriya) is frequently distributed. And extensively used by the tribal people. *Oroxylum indicum* (L.) Vent is a small or medium sized deciduous tree that grows upto a height of 12 m, with soft light brown or greyish brown bark with corky lenticels⁴. The leaves are very large, 90 - 180 cm long, 2 - 3 pinnate with 5 or more pairs of primary pinnae, rachis very stout, cylindrical, swollen at the junction of branches, leaflets 2 - 4 pairs ovate or elliptic, acuminate, glabrous. The large leaf stalks wither and fall off the tree and collect near the base of the trunk, appearing to look like a pile of broken limb bones.

The flowers are reddish purple outside and pale, pinkish-yellow within, numerous, in large erect racemes⁵. Fruits are flat capsules, 0.33 - 1 m long and 5 - 10 cm broad and sword shaped. Seeds are numerous, 6 cm long, winged all round except at the base.

	QUICK RESPONSE CODE DOI: 10.13040/IJPSR.0975-8232.IJP.5(8).532-35
	Article can be accessed online on: www.ijpjournal.com
DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5(8).532-35	

In India, it is distributed in the Himalayan foothills, Eastern and Western Ghats and North East India^{6,7}. Many other compounds namely oroxyloside methyl ester and chrysin-7-O-methyl glucoside in leaves and stem bark and an anthraquinone, aloemodin in leaves⁸ were also reported from this plant. Analysis of phyto constituents on various extracts of different parts of the plant revealed the presence of flavonoids, alkaloids, saponins, tannins, glycosides, sterols, fats and oils in high, moderate and low concentrations^{9,10}.

Leaves are used as stomachic, carminative and flatulent. Leaf decoction is given in treating rheumatic pain, enlarged spleen^{11,12}, ulcer, cough and bronchitis. The fruits are acid, sweet, antihelminthic, effective in diseases of the throat and heart, piles, bronchitis, used as expectorant, improves appetite and is used in leucoderma¹³.

MATERIALS AND METHODS:

Collection and Authentication of Plant

Material: The leaves of *Oroxylum indicum* were collected from the tribal belts of the local area of Baipariguda of Koraput District, India in the month of January 2018. The plant was identified, confirmed and authenticated by the Biju Patnaik Medicinal Plants Garden and Research Centre, Dr.

M. S. Swami Nathan Research Foundation, Jeypore, Koraput (District), Orissa (Letter No. MJ/SS/P-605/18, dated (7.2.2018)). After authentication leaves were collected in bulk and washed under running tap water to remove adhering dirt. Then the leaves were shade dried. The dried materials were made into coarse powder by grinding in mechanical grinder and stored in a closed air tight container for further use.

Preparation of Extracts: The coarse powder was taken in Soxhlet apparatus and extracted successively with ethanol, ethyl acetate, *n*-butanol and petroleum ether as solvent. A total amount of 250 g coarse powder was extracted with 500 ml of each solvent. For each solvent, 10 cycles were run to obtain thick slurry. Each slurry was then concentrated under reduced pressure to obtain crude extract. All crude extracts were kept in closed air tight containers under cool and dark place for further study^{14,15}.

Phytochemical Investigation: The crude pet. ether chloroform and ethanol extracts of the leaf of *Oroxylum indicum* were subjected to preliminary phytochemical analysis in order to detect the presence of various groups of phytoconstituents by carrying out the chemical analysis^{15,16}.

TABLE 1: PHYTOCHEMICAL SCREENING FOR THE DIFFERENT SOLVENT EXTRACTS OF *O. INDICUM* LEAVES

Extracts	Phytochemicals						
	Alkaloids	Flavonoids	Glycosides	Saponins	Tannins	Phenolic compounds	Carbohydrate
Pet. ether	+++	+++	++	++	+	++	---
Chloroform	++	++	+	--	---	+	---
Ethanol	++	+++	+	++	++	++	---

+++ , Strong; ++, moderately; +, poor presence

Animal: Healthy adult wistar strain of albino rats weighing approximately 180 to 250 g were used. They were housed in standard conditions of temperature (25 ± 2 °C), 12 h light per day cycle, relative humidity of 45 - 55% in animal house of Jeypore College of Pharmacy. They were fed with standard pellets of food and water. Animals were kept and all operation on animals was done in aseptic condition.

Experimental Protocol: Animals were selected, weighed (25 - 30 g) and divided in to five groups (n = 6), namely control, standard drug and three groups belonging to four different extract of *O. indicum*. All the studies conducted were approved

by the Institutional Ethical Committee (1906/PO/Re/S/16/CPCSEA), Jeypore College of Pharmacy, Jeypore, Odisha according to prescribed guide-lines of the Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA), Government of India.

Wound Healing Activity:

Wound Model:

Excision Wound: For the excision wound study, animals were divided into five groups of six rats in each group. Group-I served as control and received only saline 2 ml/kg orally, Group-II served as Standard drug (povidone iodine ointment) applied topically. The pet. ether, chloroform and ethanol

extracts (250 mg/kg) were given orally to Group-III, Group-IV and Group-V of *O. indicum* leave respectively. An impression was made on the dorsal thoracic central region 5 mm away from the ears, by using a round seal of 2.5 cm diameter as described by Morton and Malone. The skin of the impressed area was excised to the full thickness to obtain an area of about 500 mm² under light ether anesthesia in aseptic condition the animals were housed individually. The animals were treated daily with drugs as mentioned above under experimental design from 0th day to 18 days starting from the day of wounding. The percentage wound closure was observed on 4th, 8th, 12th, 16th, 18th post wounding day. Epithelization time (in days) and size of the scar area was noted¹⁷.

Incision Wound: In the Incision wound model, the animals were divided into five groups of six rats in each group, and kept in separate cage. Group-I served as control, received only saline 2 ml/kg orally, Group-II served as standard drug (povidone iodine ointment) applied topically, The pet. ether, chloroform and ethanol extracts (250 mg/kg) were given orally to Group- III, Group-IV and Group-V of *O. indicum* leave respectively. Under light ether anesthesia, the animals were secured to operation

table in its natural position. Two paravertebral straight incisions of 6 cm each were made through the entire thickness of the skin, on either side of the vertebral column with help of sharp blade. After complete homeostasis, the wound was closed by means of interrupted sutures placed at equidistance points about 1 cm apart. Animals were treated daily with drugs as mentioned above under experimental design from 0th day to 10th post-wounding day. The wound breaking strength was determined on 10th day by continuous constant water flow technique of Lee^{18,19}.

TABLE 2: EFFECT OF EXTRACTS OF *O. INDICUM* LEAVE ON THE BREAKING STRENGTH IN INCISION WOUND

S. no.	Group	Breaking strength
1	Control	247.11±27.03
2	petroleum ether	263.16±9.33
3	chloroform	237.21±8.41
4	ethanol	337.17±14.32*

Values are mean ± SE (n - 6)* p<0.05 vs. control

Statistical Analysis: The results are reported as Mean ± SE. Statistical analysis was done using Anova (Tukey - Multiple Comparison Test). When probability (p) was less than 0.05 was considered as significant¹⁹.




FIG. 1: EFFECT OF DIFFERENT EXTRACTS OF *OROXYLLUM INDICUM* ON EXCISION WOUND MODEL




FIG. 2: EFFECT OF DIFFERENT EXTRACTS OF *OROXYLLUM INDICUM* ON INCISION WOUND MODEL

RESULT AND DISCUSSION: The preliminary phytochemical screening showed that the different solvent extracts of *O. indicum* contains alkaloids, flavonoids, glycoside, phenolic compound, saponins and tannins in all the solvent extracts and carbohydrates absent in all the extracts. In the study using excision wound model, animals treated with ethanol extract of *O. indicum* leave showed significant decrease in epithelization period as evidenced by shorter period for fall of eschar as compared to control group (p<0.05) **Fig. 1**.

The extract also facilitated the increase in rate of wound contraction than control group. The petroleum ether extract treated animal (Group-III) showed wound contraction by 43.22%. The chloroform extract treated animal (Group-IV) showed wound contraction by 51.29%. The ethanol extract treated animal (Group-V) showed wound contraction by 81.33% as compared with the control (Group-I) by 67.24% in all the extract **Fig. 2**. The result of present study reveals that ethanolic leave extracts of *O. indicum* possess a prominent

prohealing activity in incision wound model. This was demonstrated by significant increase in the skin tensile strength in methanol extract treated groups ($p < 0.05$) on 10th post wounding day are presented in **Table 2**.

CONCLUSION: In the present study, wound healing activity of *O. indicum* was studied and the results of the present study suggest that local application and systemic administration of ethanol extract of the leaf has shown more significant wound healing activity in excision and incision wound models and support the popular use of plant to open wound in folk medicine. The wound healing property of *O. indicum* has been attributed to its antimicrobial effects. The presence of phytoconstituents like flavonoids, saponins, phenols and tannins either individually or combined together may exhibit the synergistic effect towards healing of wounds. However, further investigation employing isolation of constituents and screening models are needed for further confirmation of wound healing potential of *O. indicum* leave. Thus the traditional use has been pharmacologically validated.

ACKNOWLEDGEMENT: Authors wish to thank to local people of south eastern Odisha for providing valuable information about the plant and the author wish to express their gratitude to Jeypore College of Pharmacy, Rondapalli, Jeypore, Koraput, Odisha to co-operate throughout the project.

CONFLICT OF INTEREST: Nil

REFERENCES:

1. Jacqueline A, Hart R and Shumake L: Systems to rate the strength of scientific evidence University of Maryland Medical Center, Glenwood 2002; 40-45.

2. Chithra P, Sajithal GB and Chandra KG: Indian. J. Exp. Bio. 1998; 36: 896.
3. Jaswanth A, Akilandewari LV, Manimaran S and Ruckmani: Indian J. Pharm. Sci. 2001; 63: 41
4. Dalal NV and Rai VR: *In-vitro* propagation of *Oroxylum indicum* Vent. a medicinally important forest tree, J. For. Res. 2004; 9: 61-65.
5. Gokhale M and Bansal YK: An avowal of importance of endangered tree *Oroxylum indicum* (Linn.) Vent, Nat. Product. Radiance 2006; 5(2): 112-114.
6. Kirtikar KR and Basu DD: Indian medicinal plants Oriental enterprises, Dehradun 2001; 4: 1105-1107.
7. Jayaram K and Prasad MN: Genetic diversity in *Oroxylum indicum* (L.) Vent (Bignoniaceae), a vulnerable medicinal plant by random amplified polymorphic DNA marker, Afr. J. Biotech 2008; 7: 254-262.
8. Lawania RD, Mishra A and Gupta R: *Oroxylum indicum*: a Review, Pharmacol. J. 2010; 2(9): 304-310.
9. Ramaswamy N, Samatha T, Srinivas P and Chary RS: Phytochemical screening and TLC studies of leaves and petioles of *Oroxylum indicum* (L.) Kurz an endangered ethno medicinal tree, Int. J. Pharm. Life Sci. 2014; 4(1): 2306-2313.
10. A. Bisht, K. Zaman, M. Singh, R. Gupta and V. Singh., Pharmacological studies on *Oroxylum indicum* (Linn.) Vent. stem bark, Indian J. Nat. Prod. Resour. 2(4) (2011) 472-478.
11. Prakash P: Indian medicinal plants, Chaukhamba Sanskrit Pratishtan, Delhi 2005; 248-249.
12. Khare CP: *Oroxylum indicum*. in: Khare C.P. editor. Indian Herbal Remedies Rational western therapy, Ayurvedic and other traditional usage, Botany, New York: Springer-Verlag Berlin Heidelberg, 2004; 4: 340-341.
13. Drury CH: Ayurvedic useful plants of India, Asiatic Publishing House, Delhi 2006; 360.
14. Kokate CK, Purohit AP and Gokhale SB: Pharmacognosy, Nirali Prakashan, Pune 2007; 122-135.
15. Harbone JB: Phytochemical methods, London: Chapman and Hall, Ltd., 1973; 49-188.
16. Trease GE and Evans WC: Trease and Evans. Pharmacognosy, A Physician's Guide to Herbal Medicine, Bailliere Tindall, London, 1989; 13: 912.
17. Morton JP and Malon MH: Arch Int Pharmacodyn Ther, 1972; 196: 117.
18. Ehrlich HP and Hunt TK: J Ann of Surg 1969; 170: 203-205.
19. Lee KH: J Pharmacol Sci 1968; 57: 1042.
20. Bolton S: Analysis of variance In Pharmaceutical statistics-practical and clinical application. Marcel Dekker NY 1997; 2(1): 1123.

How to cite this article:

Samarath BK and Panda SK: Phytochemical study and evaluation of wound healing potential of crude leaf extracts of *Oroxylum indicum* (L.) vent in Wistar rats. Int J Pharmacognosy 2018; 5(8): 532-35. doi link: [http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5\(8\).532-35](http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5(8).532-35).

This Journal licensed under a Creative Commons Attribution-Non-commercial-Share Alike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)