



Received on 31 October 2018; received in revised form, 22 November 2018; accepted, 27 November 2018; published 01 December 2018

## CHEMICAL COMPOSITION AND ANTIFUNGAL ACTIVITY OF ESSENTIAL OILS FROM AERIAL PARTS OF *MENTHA PIPERITA* AND *MENTHA ARVENSIS*

Harpreet Kaur<sup>1</sup>, Ritu Tandon<sup>\*1</sup>, Anu Kalia<sup>2</sup> and Charu Maini<sup>1</sup>

Department of Chemistry<sup>1</sup>, Department of Soil Science<sup>2</sup>, Punjab Agricultural University, Ludhiana - 141004, Punjab, India.

### Keywords:

Essential oil, *Mentha piperita*, *Mentha arvensis*, GC-MS, FT-IR, Antifungal activity

### Correspondence to Author:

**Ritu Tandon**

Assistant Chemist,  
Department of Chemistry,  
Punjab Agricultural University,  
Ludhiana - 141004, Punjab, India.

**E-mail:** ritutandon@pau.edu

**ABSTRACT:** The present study deals with chemistry and antifungal activity of essential oil (EO) of two species of *Mentha viz. Mentha piperita* and *Mentha arvensis* which was extracted using Clevenger-type apparatus. Percent yield of EO of *M. arvensis* (2.8%) was found higher than *M. piperita* (2.1%). Compositional analysis of *M. piperita* and *M. arvensis* EO oil by GC-MS revealed presence of 51 and 35 compounds, respectively. Vibration spectroscopy performed in the 500-4000 cm<sup>-1</sup> wave number region depicted variations in the presence of functional groups among the two EOs. The *M. piperita* EO exhibited better antifungal potential against *Penicillium digitatum* in comparison to *M. arvensis* EO. The results were found at par with the standard Carbendazim 50 WP at 250 µg/mL. Optical research microscopy study depicted profound morphological changes in hyphae and spore bearing organs of EO treated fungus thereby supporting better antifungal activity of *M. piperita* EO.

**INTRODUCTION:** Postharvest damages in fruits have been predicted to range between 10 to 40%<sup>1</sup>. Kinnow has very diminutive storage life at ambient conditions and economically stored at low temperature by sellers for marketing in off period. Among various pathogens, attack by fungal pathogens is the major cause for the postharvest losses. The main fungal pathogens of kinnow during storage period are *P. digitatum*, *Penicillium italicum* and *Aspergillus niger*. Out of these, *P. digitatum*, the green mold, causes the major loss and is most dominant mesophilic fungus of division Ascomycota.

It is found in soils of citrus producing areas and is responsible for prevalent post-harvest diseases in citrus fruit causing huge losses due to decay caused by *P. digitatum*<sup>2</sup>. Essential oils are emerging as possible alternatives for preservatives in food industry due to their diverse antimicrobial properties. Essential oils are natural compounds produced as secondary metabolites in the plant system. These are aromatic oily volatile colourless liquids characterized by distinct odour and usually are less denser than water<sup>3</sup>.

These are rich in secondary compounds such as flavonoids, polyphenols and terpenoids and are generally synthesized in specific cell or tissue types in leaves and plant stems. The main volatile components of EOs include monoterpenes, sesquiterpenes and their oxygenated derivatives such as aldehyde, ketones, alcohols, esters and acids<sup>4</sup>. Many EOs and their primary compounds exhibit diverse biological activities such as antibacterial,

	<p><b>DOI:</b> 10.13040/IJPSR.0975-8232.IJP.5(12).767-73</p>
	<p>Article can be accessed online on: <a href="http://www.ijournal.com">www.ijournal.com</a></p>
<p><b>DOI link:</b> <a href="http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5(12).767-73">http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5(12).767-73</a></p>	

antifungal, antioxidant, antimutagenic and antimicrobial activities. Being natural and rich in antioxidants, these are considered safe for human consumption<sup>5, 6</sup>. There are 17,500 medicinal plant species among higher plants and approximately 3,000 EOs exist among which 300 are commercially important having applications in perfume, pharmaceuticals and cosmetics industries<sup>7, 8</sup>. Among various plants, the genus *Mentha* (mint), an imperative member of the Lamiaceae family, is very diverse<sup>9</sup>. It is signified by approximately 19 species and 13 natural hybrids, mainly persistent herbs growing enthusiastically in damp or wet places. Mints can grow under extensive range of agro-climatic conditions and are invasive and fast growing<sup>10</sup>.

Many species and hybrids of *Mentha* also have considerable economic importance such as *Mentha spicata*, *Mentha aquatica*, *Mentha arvensis*, *Mentha canadensis*, *Mentha pulegium* and *Mentha piperita*<sup>11</sup>. Different species of *Mentha* are utilized across the globe for their medicinal and culinary properties. Mints are extensively cultivated for their oils and terpenoid components such as menthol, carvone, linalyl acetate and linalool, for use in cosmetic, food, pharmaceutical, flavour, beverage and allied industries<sup>12</sup>. *Mentha species* essential oils have high antibacterial and antifungal activity and hence can be used as natural fungicides<sup>13</sup>. Commercially, the most sustainable specie is peppermint (*M. piperita*) which is a hybrid of cornmint (*M. canadensis*) and spearmint (*M. spicata*)<sup>5</sup>. Major bioactive compounds of *Mentha* are menthol, menthone and pulegone<sup>14</sup>.

A lot of studies related to antimicrobial activities of EOs have been performed but the anti-mycotic potential of *Mentha* EOs against postharvest pathogen, *P. digitatum* has not been explored as yet. The present investigation has been carried out on the compositional analysis of *M. arvensis* and *M. piperita* leaves EOs focusing on the morphological variations of *P. digitatum* caused by the essential oils.

## MATERIALS AND METHODS:

**Plant Material:** At full flowering stage during May 2017, the mature fresh leaves of two *Mentha species* namely *M. arvensis* and *M. piperita* were collected from Herbal Garden, Department of

Agronomy, Punjab Agricultural University, Ludhiana, India. The species were identified and authenticated by a plant agronomist, Dr. Amanpreet Singh from Department of Agronomy, Punjab Agricultural University, Ludhiana.

**Extraction of Essential Oil:** The fresh leaves were washed twice with double distilled water to remove dust, at room temperature. The fresh leaves of *M. arvensis* and *M. piperita* were coarsely grounded prior to the operation and after that subjected to hydro-distillation in Clevenger-type apparatus for 5h at 60 °C (until no more oil was obtained). Extracted essential oil was refluxed with diethyl ether and dried over anhydrous sodium sulphate, filtered and stored at 4 °C in dark amber colored vials till further analysis. The yield of essential oils was determined by using formula:

$$\text{Yield (\%)} = \frac{\text{Volume of essential oil extracted (ml)}}{\text{Weight of sample taken (g)}} \times 100$$

## Chemical Characterization of Essential Oils:

The essential oils were analyzed using GC-MS (QP2010 Plus, Shimadzu, Japan), equipped with Rtx-5 MS capillary column (30.0 m × 0.25 mm i.d., 0.25 µm film thickness) for the separation of the components of essential oil. The injector was maintained at 260 °C and operated in split injection mode with split ratio 120.0 the split valve closed for 1 min.

Helium gas was used as the carrier gas at a constant pressure of 69 kPa. The column oven was initially maintained at 50 °C for 2 min, raised to 210 °C at 3°C per min for 2 min and then to 250 °C at 6 °C per min for 6 min. The interface temperature was 270 °C and the sample was run in EI mode. Data acquisition was started 4 min after injection. The MS operating parameters used were; ionization voltage (EI) 70 eV, peak width 2 s, scan speed 1250 m/z; scan range 40-650 m/z and detector voltage 1.5 V.

**Essential Oils Identification:** The compounds of essential oils were identified based on the comparison of their relative retention times and mass spectra with literature data including Flavors and Fragrances of Natural and Synthetic Compounds 2 (FFNSC2), WILEY 8 and National Institute of Standards and Technology 14 (NIST14) libraries.

**FT-IR Analysis:** The vibrational spectroscopy of both *Mentha species* essential oil was performed on a Thermo Nicolet model 6700 FT-IR spectrometer at room temperature using ATR assembly at a resolution of  $3.86\text{ cm}^{-1}$  in the  $500\text{-}4000\text{ cm}^{-1}$  domains.

#### Determination of Antifungal Activity:

**Procurement of Test Fungus:** The culture of the test fungus *P. digitatum* isolated from kinnow fruit surface was procured from IARI (Indian Agriculture Research Institute), New Delhi with ITCC No. 7190.

**Poisoned Food Technique:** The antifungal activity of the *Mentha* essential oils against *P. digitatum* culture was evaluated by poisoned food technique<sup>15</sup>. All tests were performed in triplicates with different working concentrations ( $\mu\text{g/mL}$ ) of the oil i.e. 1000, 500, 400, 300, 200 and 100  $\mu\text{g/mL}$  of *M. piperita* and *M. arvensis* essential oils along with standard Carbendazim 50 WP (250  $\mu\text{g/mL}$ ). For this purpose, each essential oil was serially diluted using 0.5% tween 20 to prepare test solutions of 1000, 500, 400, 300, 200 and 100  $\mu\text{g/mL}$ . One ml of each concentration was incorporated into 15 ml molten potato dextrose agar medium and mixed well in petridishes. After solidification of medium, 100  $\mu\text{l}$  of prepared spore suspension containing  $10^4$  colony forming units (CFU  $\text{ml}^{-1}$ ) of fungal spores were spread over medium. The petridishes were tightly sealed with parafilm and incubated at  $25\text{ }^\circ\text{C}$  in incubator. The colony count was recorded until 7<sup>th</sup> day after inoculation (DAI). The percentage inhibition was calculated from the mean value of colony numbers of treated and untreated fungus.

#### Optical Research Microscopy of Fungal Growth:

The morphological changes in fungus after treatment were studied by optical microscopy. The fungal samples were stained with lactophenol cotton-blue dye and observed under Optical research microscope (make Leica DM 5000B) at 200 and 400 X magnifications to examine any morphological changes. Untreated samples were also stained and examined. Optical micrographs were taken with the help of CCD camera attachment (Leica DFC 420C camera).

**Statistical Analysis:** The experimental results were expressed as mean of three replicates. The data was

analysed using the SAS software 9.0. Differences between means were tested using least significant difference and treatment means were compared with Duncan's multiple range test ( $P < 0.05$ ) and t-grouping and P-value of  $P < 0.05$  was considered as significant.

## RESULTS AND DISCUSSION:

**Yield and Chemical Composition of Essential Oils:** Essential oils isolated from fresh leaves of both *Mentha species* were pale yellow in colour. The percentage yield of *M. arvensis* essential oil (2.8%) was found to be higher than *M. piperita* essential oil (2.1%). The major compounds identified by GC-MS analysis in *M. piperita* and *M. arvensis* essential oils are shown in **Table 1**. A total of 51 and 30 components were identified in *M. piperita* and *M. arvensis* essential oil, respectively.

**TABLE 1: MAJOR CONSTITUENTS (AREA %) IN MENTHA PIPERITA AND MENTHA ARVENISIS**

Name	Area (%)	
	<i>M. piperita</i>	<i>M. arvensis</i>
Menthol	58.80	84.63
Pulegone	6.62	-
Isomenthone	6.42	-
L-menthol	-	4.58
Menthyl acetate	3.94	1.11
Menthofuran	3.11	-
Cyclohexanone	-	2.97
Neomenthone	2.64	-
(E)-Caryophyllene	1.95	-
Limonene	1.62	1.54
Neoisomenthol	-	1.46
Germacrene D	1.43	-
2-Cyclohexen-1-one	1.41	-
2-Isopropyl-5-methyl cyclohexanol	1.35	-
Eucalyptol	0.90	0.07
$\alpha$ -Pinene	0.71	0.10
(E)- $\beta$ -Famesene	0.58	-
2-Isopropenyl-5-methylcyclohexanol	0.57	-
Viridiflorol	0.55	-

**Table 2** showed comparative existence of some compounds in *M. piperita* and *M. arvensis*. Menthol, the major component of both essential oils was found in higher concentration in *M. arvensis* (84.63%) than in *M. piperita* (58.80%).



However, other compounds such as pulegone (6.62%), isomenthone (6.42%), menthofuran (3.11%), neomenthone (2.64%), caryophyllene (1.94%), germacrene (1.43%), 2-cyclohexen-1-one (1.41%), 2-isopropyl-5-methyl cyclohexanol (1.35%) etc. were found to be absent in essential oil of *M. arvensis*.

**TABLE 2: COMPARISON OF MINOR CONSTITUENTS OF MENTHA PIPERITA AND MENTHA ARVENSIS**

S. no.	Name	<i>M. piperita</i>	<i>M. arvensis</i>
1	Ethyl hexanol	+	-
2	Bicyclogermacrene	+	-
3	Caryophyllene oxide	+	-
4	Phytol acetate	+	-
5	Lavandulyl acetate	+	-
6	Neomenthol	+	-
7	$\alpha$ -Cadinol	+	-
8	$\alpha$ -Bourbonene	+	-
9	Neomenthyl acetate	+	-
10	$\delta$ -Cadinene	+	-
11	Bicyclo[3.1.0]hexane-6-methanol	+	-
12	1,6-Octadien-3-ol	+	-
13	Spathulenol	+	-
14	Trans-Nerolidol	+	-
15	2,5-diethyl-Tetrahydrofuran	+	-
16	Isomenthyl acetate	+	-
17	Eugenol	+	-
18	$\alpha$ -Gurjunene	+	-
19	(+)-4-Carene	+	-
20	3-Cyclohexene-1-methanol	+	-
21	BHT	+	-
22	1-Octen-3-ol	+	-
23	4,8- $\beta$ -epoxy-Caryophyllane	+	-
24	P-Menthane-1,2,3-triol	+	-
25	Ispulegyl acetate	+	-


**FT-IR Analysis:** The FT-IR spectra of essential oil of both *Mentha species* revealed distinct variation for occurrence of several peaks indicating variation in compounds possessing different functional groups. The FT-IR spectrum of *M. piperita* showed a broad band at  $3367.1\text{ cm}^{-1}$  due to bonded O-H stretching, a band at  $1706\text{ cm}^{-1}$  due to C=O stretching and a sharp band at  $2888.7\text{ cm}^{-1}$  due to C-H stretching whereas in *M. arvensis* the similar stretching bands were observed at  $3346\text{ cm}^{-1}$ ,  $1656\text{ cm}^{-1}$  and  $2884.8\text{ cm}^{-1}$  respectively **Fig. 1A** and **Fig. 1B**. However, a sharp bonded O-H stretch band and small overtone due to C=O band was observed in FT-IR spectrum of *M. arvensis* in comparison to *M. piperita* essential oil.

**Antifungal Activity Assay:** The antifungal activities of *Mentha species* essential oils were assessed against *P. digitatum* w.r.t. absolute control and standard Carbendazim 50 WP. Results of the antifungal activity on seventh day after inoculation (DAI) showed 100% inhibition at  $1000\text{ }\mu\text{g/mL}$  in essential oils obtained from both species whereas at  $100\text{-}500\text{ }\mu\text{g/mL}$  *M. piperita* essential oil showed better percent inhibition compared to *M. arvensis* essential oil **Fig. 2**.

**FIG. 1: FT-IR SPECTRA OF ESSENTIAL OILS EXTRACTED FROM MENTHA PIPERATA (1A) AND MENTHA ARVENSIS (1B) USING ATTENUATED TOTAL REFLECTANCE MODE****FIG. 2: ANTIFUNGAL ACTIVITY OF THE ESSENTIAL OIL OF M. PIPERITA AND M. ARVENSIS AGAINST P. DIGITATUM**

The maximum inhibition was observed at 500 µg/mL for both essential oils i.e. 71% and 55% *M. piperita* and *M. arvensis* essential oils respectively. Standard Carbendazim 50 WP showed 99.7% inhibition which was found higher than both the


essential oils **Fig. 3B, 3C and 3D**. All the interactions i.e. between different concentrations of both essential oils with respect to control (i.e. without treatment) **Fig. 3A** was significant with coefficient of variance value of 0.765.




**FIG. 3: ANTIFUNGAL ACTIVITY OF *M. PIPERITA* AND *M. ARVENSIS* ESSENTIAL OIL AT 500 µg/mL 7<sup>th</sup> DAY AFTER INOCULATION (3A) CONTROL (3B) CARBENDAZIM 50 WP (3C) *M. PIPERITA* AND (3D) *M. ARVENSIS***

**Optical Research Microscopy of Essential Oil - *P. Digitatum* Studies:** Optical research micrographs of control *P. digitatum* showed presence of

spores in groups and normal morphology of hyphae possessing dense cytoplasm **Fig. 4A and 4B**.



**FIG. 4: OPTICAL RESEARCH MICROSCOPY SHOWING HYPHAE AND SPORE MORPHOLOGY OF CONTROL 4(A) & 4(B) FUNGUS TREATED WITH 500 µg/mL *M. ARVENSIS* EO 4(C) & 4(D) FUNGUS TREATED WITH *M. PIPERITA* EO 4(E) & 4(F)**



**FIG. 5: HYPHAL DIAMETER (µm) OF CONTROL (A) AND FUNGAL MYCELIA TREATED WITH *M. PIPERITA* (B) AND *M. ARVENSIS*(C)**

Whereas, on treatment with *M. piperita* **Fig. 4C, 4D** and *M. arvensis* **4E and 4F** essential oil, scattered spores in the form of smaller groups and as short spore chains respectively. The optical micrographs also exhibited modified hyphal morphology indicated by decrease in diameter and number of hyphal branches than control. Moreover, cytoplasm was less dense exhibiting shrinkage which led to fragmentation of fungal hyphae, loss of linearity including irregular hyphae and formation of warty surfaces (cell wall pitting phenomena) in fungus treated with *M. piperita*

essential oil. The hyphal diameter was observed to be greater in untreated fungus followed by fungus treated with *M. arvensis* and *M. piperita* essential oil **Fig. 5**.

**DISCUSSION:** Percent yield of essential oils of both the species are well in correlation with the available literature with minute variations<sup>13,16</sup>. The GC-MS of essential oil of both the *Mentha species* showed a broad spectrum of compounds with distinct variations. These variations are well explained due to different climatic conditions in which the crop was reared. Further, time, season of sample collection and soil composition of the sampling area also contributed to the composition of essential oil.

Almost all chemical compounds reported in essential oil from fresh leaves of *M. piperita* exhibited one or the other biological activity<sup>17,18</sup>. Differences in compositional analysis by GC-MS correlate very well with the FT-IR spectra of both species essential oils since more predominant bands were observed for *M. piperita* essential oil than *M. arvensis* essential oil. The FT-IR spectra of *M. piperita* depicted more interactions between various compounds may be because of greater number of compounds in it as shown by its GC-MS. The position of the absorption band due to valence vibration of O-H bonds can be used to quantify hydrogen bonding. When hydrogen bonding is stronger, the O-H bond length increases and bond force constant decreases, so the valence vibration is identified at lower frequency values compared to the values determined in the absence of association with hydrogen bonds. The association through hydrogen bonds led to wide band at  $3367.1\text{ cm}^{-1}$  in *M. piperita* essential oil and sharp band at  $3346\text{ cm}^{-1}$  in *M. arvensis* essential oil and a red shift occurred due to O-H stretching by  $11.1\text{ cm}^{-1}$  of *M. arvensis* essential oil in comparison to *M. piperita* essential oil.

Samfira et al.,<sup>19</sup> reported O-H stretch at  $2950\text{ cm}^{-1}$  and C=O peak at  $1700\text{ cm}^{-1}$  for *M. piperita* essential oil extracted from leaves collected from western part of Romania. Prakash and Yunus<sup>20</sup> found O-H stretch at  $3355.6\text{ cm}^{-1}$  and C=O peak at  $1702.5\text{ cm}^{-1}$  for *M. arvensis* essential oil extracted from leaves collected from Lucknow city. Shift in frequency bands of functional groups was observed

due to climatic changes. The results from poisoned food technique followed by percentage inhibition calculation showed more per cent inhibition with less number of colonies in *M. piperita* at all tested concentrations than in *M. arvensis* at similar concentrations. At the concentration of  $1000\mu\text{g/mL}$  both species essential oils showed antifungal activity at par with standard Carbendazim 50 WP.

Results observed in our studies confirmed enhanced efficacy of *M. piperita* essential oil (71%) compared to *M. arvensis* essential oil (55%) at  $500\mu\text{g/ml}$  in controlling the dreaded post-harvest fungus *P. digitatum*. Essential oil of *M. piperita* leaves possess significant antifungal activity may be due to presence of pulegone, isomenthone, menthofuran and neomenthone which were identified absent in *M. arvensis* essential oil.

Optical research microscopy of fungus treated with essential oils revealed fragmentation of fungal hyphae, loss of linearity including irregular hyphae and formation of warty surfaces. In fungus treated with *M. piperita* essential oil malformed conidiophore structures were recorded which culminated to formation of impaired conidia.

All these findings indicate that the antifungal mode of action of essential oil on *P. digitatum* is a result of damage to the cell plasma membrane integrity. Owing to our results, similar phenomenon of the lipophilicity of essential oils causing membrane expansion, augmented membrane permeability and fluidity, inhibition of respiration and variation in ion transport processes of fungi which results into induced the outflow of other cellular components was observed in earlier studies<sup>21,22,23</sup>.

**CONCLUSION:** The per cent yields of *M. piperita* and *M. arvensis* essential oil obtained from leaves was calculated. Comparison showed that *M. arvensis* leaf essential oil yield (2.8%) was higher as compared to *M. piperita* (2.1%). Gas chromatography- Mass Spectrometry (GC-MS) of *M. piperita* and *M. arvensis* essential oil revealed the presence of 51 and 30 compounds respectively, menthol being the major compound in both essential oils. The FT-IR spectra of essential oil of both *Mentha species* showed prominent variation in compounds indicated by different shape and position of IR bands.

Screening of antifungal activity revealed that *M. piperita* essential oil (71%) was found to be more effective than *M. arvensis* (55%) against *P. digitatum*. Optical research microscopy further revealed that the essential oils showed their efficacy by disrupting the cell plasma membrane, thereby causing the fragmentation of fungal hyphae.

**ACKNOWLEDGEMENT:** The authors wish to thank the Director, Punjab Horticultural Post-Harvest Technology Centre, Punjab Agricultural University, Ludhiana, Punjab and the Director, EMNL, for providing the necessary infrastructural facilities and Head, Department of Chemistry, College of Basic Sciences and Humanities, Punjab Agricultural University, Ludhiana, Punjab, India for providing the contingency for purchase of consumables for this study.

**CONFLICT OF INTEREST:** The author declare no conflict of interest.

#### REFERENCES:

1. Kaur N, Bala A and Kumar A: Prevent kinnow produce from post harvest disorders and diseases. *Rashtriya Krishi* 2014; 9(1): 45-46.
2. Mandal G: Effect of lac-wax, citrashine and individual shrink wrapping of fruits on storage life of late harvested kinnow under ambient condition. *IJBEAS* 2015; 1(3): 84-89.
3. Miguel MG: Antioxidant and anti-inflammatory activities of essential oil: A short review. *Molecules* 2010; 15: 9252-9287.
4. Sutar I, Akkol EK, Tosun A and Keles H: Comparative pharmacological and phytochemical investigation on the wound-healing effects of the frequently used essential oils. *J Essent Oil Res* 2014; 26(1): 41-49.
5. Bozovic M, Pirolli A and Ragno R: *Mentha suaveolensehrh* (Lamiaceae) essential oil and its main constituent piperitenone oxide: Biological activities and chemistry. *Molecules* 2015; 20: 8605-8633.
6. Dadalioglu I and Evrendilek GA: Chemical compositions and antibacterial effects of essential oils of turkish oregano (*Origanum minutiflorum*), bay laurel (*Laurus nobilis*), spanish lavender (*Lavandula stoechas* L.) and fennel (*Foeniculum vulgare*) on common food borne pathogens. *J Agric Food Chem* 2004; 52: 8255-8260.
7. Bruneton J: Pharmacognosy, Phytochemistry, Medicinal plants: Essential oils. Lavoisier Publishing, Paris. Edition 2, 1999: 461-780.
8. Bakkali F, Averbeck S, Averbeck D and Idaomar M: Biological effects of essential oils-A review. *Food Chem Toxicol* 2008; 46: 446-475.
9. Elansary HO and Ashmawy NA: Essential oils of mint between benefits and hazards. *J Essent Oil Bear PI* 2013; 16(4): 429-438.
10. Kumar P, Mishra S, Malik A and Satya S: Insecticidal properties of *Mentha species*. *Ind Crops Prod* 2011; 34: 802-817.
11. Mirzaie-Nodoushan H, Rezale MB and Jaimand K: Path analysis of the essential oil-related characters in *Mentha spp*. *Flavour Frag J* 2001; 16(6): 340-343.
12. Chaddha KL and Gupta R: Japanese mint. *Adv Horticult* 1995; 11: 687-725.
13. Moghtader M: *In-vitro* antifungal effects of the essential oil of *Mentha piperita* L. and its comparison with synthetic menthol on *Aspergillus niger*. *Afr J Plant Sci* 2013; 7(11): 521-527.
14. Luca A and Ciucanu I: Solid-phase microextraction of menthol from peppermint. *Ser Chem* 2011; 20(1): 71-80.
15. Balouiri M, Sadiki M and Ibsouda SK: Methods for *in-vitro* evaluating antimicrobial activity: A review. *J Pharma Anal* 2016; 6: 71-79.
16. Gende LB, Mendiara S, Fernandez NJ, Baren CV, Lira AD, Bandoni A, Fritz R, Floris I and Eguaras M: Essentials oils of some *Mentha spp.* and their relation with antimicrobial activity against *Paenibacillus larvae*, the causative agent of American foulbrood in honey bees, by using the bioautography technique. *Bulletin of Insectology* 2014; 67(1): 13-20.
17. Satoyanova A and Georgiev E: Chemical composition and antimicrobial activity of Bulgarian peppermint oils. *Bul Ess Oil Perfum Cosm (Bulgaria)* 2003; 43-47.
18. Georgiev S and Topalova V: Chemical composition and antimicrobial activity of Bulgarian peppermint oils. *Plant Sci* 1989; 26: 33-37.
19. Samfira I, Rodino S, Petrache P, Cristina RT, Butu M and Butnariu M: Characterization and identity confirmation of essential oils by mid infrared absorption spectrophotometry. *Digest Journal of Nanomaterials and Biostructures* 2015; 10(2): 557-566.
20. Prakash N and Yunus M: Fourier transform infrared spectroscopy analysis of oil of *Mentha arvensis* grown at sites varying with vehicular traffic loads in Lucknow city, India. *Int J Environ* 2013; 2(1): 16-25.
21. Wu Y, Yang O and Tao N: Plasma membrane damage contributes to antifungal activity of citronellal against *Penicillium digitatum*. *J Food Sci Technol* 2016; 53(10): 3853-3858.
22. Tyagi AK and Malik A: Liquid and vapour-phase antifungal activities of selected essential oils against *Candida albicans*: microscopic observations and chemical characterization of *C. citratus*. *BMC Complement Altern Med* 2010; 10: 1-65.
23. Zhou HE, Tao NG and Jia L: Antifungal activity of citraloctanal and  $\alpha$ -terpineol against *Geotrichumcitri aurantii*. *Food Control* 2014; 37: 277-283.

#### How to cite this article:

Kaur H, Tandon R, Kalia A and Maini C: Chemical composition and antifungal activity of essential oils from aerial parts of *Mentha piperita* and *Mentha arvensis*. *Int J Pharmacognosy* 2018; 5(12): 767-73. doi link: [http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5\(12\).767-73](http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5(12).767-73).

This Journal licensed under a Creative Commons Attribution-Non-commercial-Share Alike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)