



Received on 15 January, 2017; received in revised form, 22 March, 2017; accepted, 25 March, 2017; published 01 April, 2017

PHYTOCHEMICAL, *IN VITRO* ANTIOXIDANT AND ANTIBACTERIAL ACTIVITIES OF *ARGYREIA PILOSA* WIGHT & ARN. (WHOLE PLANT)

D. S. N. B. K. Prasanth ^{*1}, Atla Srinivasa Rao ² and Rajendra Prasad Yejella ³

Department of Pharmacy ¹, JNTUK, Kakinada- 533003, Andhra Pradesh, India.

Shri Vishnu College of Pharmacy ², Bhimavaram, Andhra Pradesh, India.

University College of Pharmaceutical Sciences ³, Visakhapatnam, Andhra Pradesh, India.

Keywords:

Argyrea pilosa, Antioxidant, Tannins, Flavonoids, Antibacterial and Phytochemical analysis

Correspondence to Author:

D. S. N. B. K. Prasanth

Department of Pharmacy,
JNTUK, Kakinada- 533003,
Andhra Pradesh, India.

E-mail: dsnbkprasanth@gmail.com

ABSTRACT: Background: The whole plant of *Argyrea pilosa* Wight & Arn. (Convolvulaceae) continuously to be utilized in Indian indigenous systems as a remedy for cough, quinsy and applied externally in case of itch, eczema and other skin troubles, antidiabetic, antiphlogistic, rheumatism and reduce burning sensation. Still, there were constrained phytochemical or biological studies on the whole plant of *A. pilosa*, as well as no studies which align using its conventional medicinal uses. **Aim:** The aim of this study was to determine the total tannin and flavonoids contents, antimicrobial and antioxidant activity of two extracts of the whole plant of *A. pilosa* and compare them to find out the better extract for upcoming studies. **Materials and Methods:** The whole plant material was subjected to cold maceration with ethyl acetate and methanol to obtain ethyl acetate extract (APEE) and methanolic extract (APME). Phytochemical constituents of these extracts were determined as per standard procedure. Total Phenolics had been estimated by the Folin– Ciocalteu colorimetric method using tannic acid as standard. Total Flavonoids had been estimated by aluminum chloride assay using quercetin as standard and the antioxidant capacity was determined by the 2,2-diphenyl-1-picrylhydrazyl radical scavenging assay, hydrogen peroxide assay and reducing capacity. The antimicrobial activity was studied with the agar diffusion method and minimal inhibitory concentrations (MIC) of these extracts were determined. **Results:** APEE has better yield than APME. The phytochemical constituents present in both APEE and APME have revealed the presence of flavonoids, alkaloids, phenols, tannins, glycosides, amino acids and proteins except for steroids and acid compounds which are only present in APME. Total tannin content and total flavonoids content of APEE is more than in APME. Both REE and RME have antioxidant and antibacterial potency. Statistically there is no significant difference between the antioxidant potency of APEE and APME. But APEE and APME are statistically different from each other in terms of their antibacterial strength, APEE being better than APME in this case.

INTRODUCTION: Phytochemicals tend to be widely available at diverse ranges in several medicinal plants. Most of the plants utilized in conventional drugs are efficient for different disorders triggered by oxidative stress and microbial infections.

Studies have revealed that medicinal plants show antioxidant along with antimicrobial activity ¹. Oxygen, the most crucial element for the endurance of living organisms is extremely reactive and may do severe harm to healthy cells of the body as free radical. Oxidation generates free radicals that begins sequence reaction within the cell that triggers harm to and also fatality of cell ². An antioxidant ends these type of radical responses by eliminating free radical intermediates, that causes aging and disorders such as atherosclerosis, cancer, cardiovascular diseases and CNS disorders ³.

	QUICK RESPONSE CODE
	DOI: 10.13040/IJPSR.0975-8232.IJP.4(4).109-17
Article can be accessed online on: www.ijpjournal.com	
DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.4(4).109-17	

Therefore, play a important role in prohibiting or lagging the oxidation at the cellular level ⁴. Man made antioxidants *i.e.*, butylated hydroxytoluene, butylated hydroxyanisole and tert-butyl hydroquinone being examined all over the world for their possible health risks and have been found to promote hepatic damage and leading to cancer to laboratory animals ⁵. In contrast, the herbs prove to have less or no side effects, ³ thus channelizing the interest towards the nature for antioxidant potency.

As a result of developing resistance to antibiotics of several microbes, plant extracts and plant derived compounds are of current interest as antiseptics as well as antimicrobial agents in medication. The effectiveness of the plants in treating various disorders is certainly well established along with a significant level of study has been carried out within this area by investigators in India as well as abroad. The worldwide breakthrough of multi-drug resistant bacterial strains is progressively restricting the potency of existing medicines and considerably leads to remedy failing of infections. For example include methicillin-resistant staphylococci, pneumococci resistant to penicillin and macrolides, vancomycin-resistant enterococci and multi-drug resistant Gram-negative organisms ¹.

A. pilosa is an ornamental, in addition to a medicinal plant. All parts of this plant are widely used as a folklore medicine for the treatment of various ailments by the Indian traditional healer. Its root is utilized to cure a various illness like sexually transmitted diseases *viz.*, gonorrhoea and syphilis, blood diseases. Traditionally, the paste of the leaves is applied to the neck region for cough, quinsy and applied externally in case of itch, eczema and other skin troubles, antidiabetic, antiphlogistic, rheumatism, reduce burning sensation and antidiabetic ⁶. Young wines are mixed together with rhizome of ginger are spread all around the body to relieve from fever ⁷. The decoction of its root used to treat diarrhea and cathartic ^{8,9}.

A vast range of phytochemical constituents has been separated from the genus *Argyreia i.e.*, glycosides, alkaloids, amino acids, proteins, flavonoids, triterpene and steroids ¹⁰. The genus *Argyreia* has been reported various biological activities including nootropic, aphrodisiac,

antioxidant, antiulcer, immune-modulatory, hepatoprotective, anti-inflammatory, anti-hyperglycemic, antidiarrheal, antimicrobial, antiviral, nematocidal, anticonvulsant, analgesic, anti-inflammatory, wound healing and central nervous depressant activities ¹⁰⁻¹⁴. Even though the drug has many uses, it's pharmacological and phytochemistry are very poorly explored ¹⁵.

Although the plant has been extensively used for its traditional value, but antibacterial and antioxidant activities remains unexplored. Therefore the current investigation had been carried out to study the antimicrobial and antioxidant potency of this medicinal plant.

MATERIALS AND METHODS:

Chemicals: All analytical grade chemicals were utilized in this study were procured from E. Merck, Germany. 2, 2-diphenyl-1-picrylhydrazyl (DPPH), tannic acid, ascorbic acid, quercetin, Folin-Ciocalteu's reagent, aluminum chloride, hydrogen peroxide, ferric chloride, potassium ferricyanide, Dimethyl sulfoxide, agar powder, beef extract and peptone were obtained from Hi-Media, Mumbai; petroleum ether, toluene, acetone, ethyl acetate and methanol.

Plant Material: The plant material was obtained from Tirupati, chittoor district of Andhra Pradesh, India during the month of March 2016 and authenticated by Dr. K. Madhava chetty, Taxonomist, Sri venkateswara University tirupati, India. Voucher specimen No. 1922 was deposited at the herbarium for future reference. Then it was shade dried, powdered and sieved through 20 mesh and kept in an air tight container for future use ^{16, 17}.

Extraction Method: The whole plant of *A. pilosa* were collected, washed in tap water to get rid of any contaminants and then dried to remove water from the surface of the plant. The plant was cut into small pieces and dried in shade in room temperature for 7 days. The dried plant material was grinded to coarse powder and defatted using petroleum ether. Equal quantity of defatted dried material of *Argyreia pilosa* was subjected to maceration with ethyl acetate and methanol for 72 hr each. It was then filtered and concentrated (rotary-evaporator) to obtain dried ethyl acetate

extract (APEE) and methanolic extract (APME). Percentage yield was calculated using the following Eqn. 1.

$$\text{Percentage yield (\%)} = \frac{\text{Weight of Extract (g)}}{\text{Weight of defatted dried plant material (g)}} \times 100$$

Qualitative Phytochemical analysis: The extracts had been analyzed for the presence of alkaloid, glycoside, steroids, triterpenoids, flavonoids, proteins, amino acids, carbohydrates, reducing sugar, tannin and saponin using standard methods^{15, 16, 17}.

Quantitative Phytochemical analysis:

Determination of total flavonoids content: Total flavonoids contents of two extracts had been estimated by aluminum chloride colorimetric assay depending on the development of a complex flavonoid-aluminium, having a maximum absorbance at 510 nm¹⁸. Quercetin utilized to make the calibration curve. One milliliter of sample or Quercetin standard solution was added into a 10 mL volumetric flask, mixed with 4 mL of 60% ethanol and 0.3 mL of 5% NaNO₂ for 6 min, After that, 0.3 mL of 10% AlCl₃ was added to react for another 6 min. The reaction was stopped by 4 mL of 4% NaOH, and the total volume was topped up to 10 mL with 60% ethanol. The absorbance was measured after 15 min at 510 nm. Total flavonoids contents were expressed as mg Quercetin equivalents per gram dry weight of each extract¹⁹. All samples were analyzed in triplicate.

Determination of total tannin content: Total tannin content (TTC) was estimated using Folin-Ciocalteu reagent (FCR) taking tannic acid as standard²⁰. FCR being sensitive to reducing compounds including polyphenols produce blue colour upon reaction. The extracts utilized in this analysis should be in the concentration of 1 mg/ml. The reaction mixture was made by combining 0.5 ml of extract, 2.5 ml of 10% Folin-Ciocalteu's reagent mixed with water, in addition to add 2.5 ml of 7.5% Na₂CO₃. Blank was prepared by same procedure without sample. The samples had been incubated in a thermostat at 45 C for 45 min. The absorbance was determined by making use of spectrophotometer at $\lambda_{\text{max}} = 765 \text{ nm}$. The samples have been prepared in triplicate for each analysis and the mean value of absorbance had been

acquired. The same method was repeated for the standard solution of tannic acid and the calibration line was construed. Based on the measured absorbance, the concentration of phenolics was calculated (mg/ml) from the calibration line. All the determinations were done in triplicate. The total phenolic content is expressed in mg tannic acid equivalent (TAE) per gram.

Antioxidant Assays:

DPPH free radical scavenging assay: DPPH scavenging capacity of each extract was determined according to the method of Md. Nur Alam *et. al.* A stock solution made up of 40 µg/mL of DPPH (in anhydrous ethanol, w/v) had been prepared. 1 ml of sample was mixed with 4 ml of ethanol DPPH solution. The mixture had been shaken vigorously as well as incubated at room temperature for 30 min in the dark. The absorbance was determined at 517 nm. A control was prepared without sample. Ascorbic acid was used as positive controls. The DPPH radical scavenging ability was calculated according to the following equation:

$$\text{DPPH scavenging rate (\%)} = \frac{[A_0 - A_1]}{A_0} \times 100$$

Where A₀ is the absorbance of control, and A₁ is the absorbance of sample. Each sample was analyzed in triplicate²¹.

Hydrogen peroxide scavenging activity assay: A solution of hydrogen peroxide (40 mM) is prepared in phosphate buffer (50 mM pH 7.4). The concentration of hydrogen peroxide is determined by absorption at 230 nm using a spectrophotometer. Extract (20–60 lg/mL) in distilled water is added to hydrogen peroxide and absorbance at 230 nm is determined after 10 min against a blank solution containing phosphate buffer without hydrogen peroxide. The percentage of hydrogen peroxide scavenging is calculated as follows:

$$\% \text{ scavenged (H}_2\text{O}_2) = \frac{[A_i - A_t]}{A_i} \times 100$$

Where A_i is the absorbance of control and A_t is the absorbance of test^{21, 22}.

Reducing power method: In the method, 2.5 ml of 0.2 M phosphate buffer with pH 6.6 and 2.5 ml of K₃Fe (CN)₆ (1% w/v) are combined to 1.0 ml of sample mixed in distilled water. The resulting mixture is incubated at 50 °C for 20 min, followed

by the addition of 2.5 mL of Trichloro acetic acid (10% w/v). The mixture is centrifuged at 3000 rpm for 10 min to collect the upper layer of the solution (2.5 mL), mixed with distilled water (2.5 mL) and 0.5 mL of FeCl₃ (0.1%, w/v). The absorbance is then measured at 700 nm against blank sample^{21, 23}.

Antibacterial activity:

Test microorganisms: The antibacterial activity of the plant extracts had been examined against some Gram positive and Gram negative bacteria *i.e.*, *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *streptococcus epidermidis* and *Escherichia coli*, which are obtained from Laboratory of microbiology, NICU, SUM Hospital, Bhubaneswar.

Well diffusion method for determination of zone of inhibition:

The antibacterial study of APCE, APEE and APME had been carried out through well diffusion method^{24, 25}. The inoculation of the microorganism had been prepared from bacterial culture. 15 ml of nutrient agar medium was put in clean sterilized petri-plates and allowed to cool and solidify. 100µl of broth of bacterial stain was pipetted out and spread over the medium evenly by a spreading rod till it is dried properly. Wells of 6 mm in diameter are bored using sterile cork borers. Solutions of both the extracts (10 mg/ml) in DMSO (dimethyl sulphoxide) were prepared. 100 µl of plant extract solutions was added to the wells. The petri-plates were incubated at 37 °C for 24 h. Ciprofloxacin (1 mg/ml) was used as positive control and DMSO was taken as negative control.

Antibacterial activity was evaluated by measuring the zone of inhibition diameters. All the determinations were performed in triplicate.

RESULTS AND DISCUSSION: Extraction of defatted plant material of *A. pilosa* had been performed utilizing chloroform, ethyl acetate and methanol as liquid extractor to yield dried APCE (Buff powder), APEE (Blackish green powder) and APME (Dark Green Powder). Defatting of plant material is a pre-extraction process that eliminates unwanted fatty and waxy substances from the plant material. On determining percentage yield (%) using Eqn.1, % yield of APME (5.38%) and APEE (2.25%) are shown in **Fig. 1**.

Percentage (%) yield= Weight of extract (g) / Weight of plant material (g) X 100




FIG. 1: PERCENTAGE (%) YIELD OF EXTRACTS

The results of qualitative phytochemical analysis of crude powder of *A. pilosa* are tabulated in **Table 1**.

TABLE 1: PHYTOCHEMICAL ANALYSIS OF VARIOUS EXTRACTS OF WHOLE PLANT OF ARGYREIA PILOSA WIGHT & ARN.

Phytoconstituents	Method	Ethylacetate Extract	Methanolic Extract
Flavonoids	Shinoda Test	+	+
	Zn. Hydrochloride test	+	+
	Lead acetate Test	+	+
Volatile oil	Stain test	-	-
	Alkaloids	Wagner Test	+
Tannins & Phenols	Hager's Test	+	+
	FeCl ₃ Test	+	+
	Potassium dichromate test	+	+
Saponins	Foaming Test	-	-
Steroids	Salkowski test	-	+
Carbohydrates	Molish test	-	-
Acid compounds	Litmus test	-	+
	Glycoside	Keller-Killani Test	+
Amino acids	Ninhydrin test	+	+
Proteins	Biuret	+	+

The outcomes of quantitative estimation of total tannin content (TTC) and total flavonoid content (TFC) are being represented in **Fig. 2 and 3**

respectively. It is definitely observable through the results that TTC and TFC in REE > RME.




FIG. 2: STANDARD CURVE OF TANNIC ACID AND COMPARISON BY GRAPHICAL REPRESENTATION OF TOTAL TANNIN CONTENT




FIG. 3: STANDARD CURVE OF QUERCETIN AND COMPARISON BY GRAPHICAL REPRESENTATION OF TOTAL FLAVONOIDS CONTENT

Antioxidant activity of any crude extract might be because of several mechanisms. Binding of transition metal ion, radical scavenging, decomposition of peroxide, reducing capacity, prevention of continued hydrogen abstraction and prevention of chain initiation are some of these mechanisms. DPPH is a stable free radical having absorption at 515 nm. When reduced by an antioxidant to form DPPH, the natural deep violet colour of DPPH changes to pale yellow²⁶. The change in the colour is proportional to the strength of the antioxidants and a significant decrease in the absorbance of the reaction mixture indicates significant free radical activity of the test material. Hydrogen peroxide (H₂O₂) scavenging activity of

the plant extracts are assessed spectrophotometrically with the disappearance of H₂O₂ at 230 nm²⁷. H₂O₂ generates hydroxyl radical within the tissues and is harmful to them. Therefore, removal of H₂O₂ is extremely important²⁸. Results of Hydrogen peroxide (H₂O₂) scavenging activity is presented in **Fig. 4**.

Based on the reducing power of the screening substance, the yellow colour of the Fe(III) becomes to Fe(II), Perl's Prussian blue which may be measured spectrophotometrically at 700 nm²¹. The reductive capacity of APEE and APME is compared to that of the Ascorbic acid in **Fig. 5**.

It is noticed that there exists positive correlation between the antioxidant activity and reducing power of APEE and APME. Like the antioxidant activity study, reducing power of REE and RME also increases with the increase in concentration.

Reductones, which exhibit the antioxidant activity by breaking the free radical chain and donating a hydrogen atom, are also responsible for the reducing properties²⁹.




FIG. 4: ANTIOXIDANT ACTIVITY STUDY OF ASCORBIC ACID, APME AND APEE




FIG. 5: REDUCING POWER ASSAY OF ASCORBIC ACID, APEE AND APME

The IC_{50} values of APEE and APME are portrayed in **Table 2**. The IC_{50} value of APEE is less than APME, which states the fact that APEE shows better antioxidant property than APME. The antioxidant activity has positive correlation with total phenolic and flavonoids content¹, which act as scavengers to various oxidising species³⁰. For that reason, it was quite sensible to determine the TPC and TFC in the plant extract. TPC and TFC of REE is more than that of RME as evident from **Fig. 3B** and **Fig. 3F**, respectively reasoning the result obtained that IC_{50} of REE is less than that of RME.

TABLE 2: IC_{50} VALUES OF APEE AND APME

Testing sample	IC_{50} (µg/ml)	
	DPPH radical scavenging activity	Hydrogen peroxide assay
Ascorbic acid	34.14	32.67
APEE	140.23	132.32
APME	147.34	137.53

For the current study, the potency of APEE and APME is screened against gram positive and gram negative bacteria. The well diffusion method of screening antibacterial activity is considered to provide better results and is associated with the carrier solvent, DMSO which is likely to diffuse highly across the medium³¹.

In this study, DMSO is employed as a solvent for solubilizing APEE and APME for antibacterial study. DMSO is a polar aprotic solvent which has the capability to dissolve both polar and nonpolar compounds. DMSO can be used as cryoprotectant for long term low temperature conservation of various biomaterials³² and thus considered to have no impact on the growth or inhibition of growth on microorganisms. Although the literature facilitates the use of DMSO as solvent with regard to antibacterial studies, a negative control (*i.e.*, only DMSO) is also studied in each plate to be extra sure.

In the present circumstances of increasing resistance of microbes towards the worldwide used Antibacterial ³³, it is crucial that the actual ingredients having antibacterial potential needs to be extracted. The current study is useful to find the extract that exhibits better antibacterial activity. In this study, the plant material was extracted with ethyl acetate and methanol by maceration. This study also compares the antibacterial activity of ethyl acetate and methanolic extracts of *A. pilosa*, APEE and APME, respectively. Results depicted in **Fig. 6** and **7** shows that both REE and RME at different concentrations 1, 10, 100, 250, 500 and 1000 µg/ml (100 µl incorporated in each well) possess antibacterial activity towards both gram-positive and gram-negative bacteria. DMSO shows no zone of inhibition (ZI).

All the results are mean±SD (n=3). The antibacterial activity of APEE and APME may be attributed to TTC and TFC in APEE and RME. Flavonoids and phenolic compounds are recognized for their role in protecting plants against microbial attack ³⁰. Tannins also act as astringents and have the ability to bind and precipitate and shrink proteins ³⁴. This ability of tannins directly damages the bacterial cell wall and precipitate proteins killing the bacteria ^{35, 36}. From the results of ZI depicted in **Table 3** and from the **Fig. 5** which compares the ZI formed by REE and RME against different bacteria, it is deemed fit to state that REE has better antibacterial activity than RME.

TABLE 3: ZONE OF INHIBITION (ZI) OF THE EXTRACTS (APEE AND APME) AGAINST DIFFERENT GRAM NEGATIVE AND GRAM POSITIVE BACTERIA

Test Organisms	APEE(µg/ml) ZI±S.D (mm)						APME(µg/ml) ZI±S.D (mm)					
	Standard (5µg/ml)	Concentration (µg/ml)					Standard (5µg/ml)	Concentration (µg/ml)				
		10	100	250	500	1000		10	100	250	500	1000
<i>E. coli</i>	18±0.577	5±0.232	8±0.179	12±0.545	15±0.318	17±0.253	18±0.225	4±0.179	6±0.325	12±0.291	14±0.381	16±0.485
<i>P.aeruginosa</i>	19±0.623	4±1.332	6±0.265	8±0.21	9±0.229	12±0.789	19±0.623	4±0.21	5±0.212	6±0.296	8±0.288	10±0.568
<i>S. aureus</i>	17±0.232	5±0.323	9±0.364	13±0.598	14±0.558	15±0.553	17±0.364	3±0.359	8±0.206	12±0.402	13±0.577	14±1.015
<i>B. subtilis</i>	16±1.23	4±0.623	6±0.485	7±0.307	8±0.615	10±0.458	16±0.485	4±0.489	5±0.223	7±0.525	8±0.750	10±0.5




FIG. 6: ANTIBACTERIAL ACTIVITY OF APEE AND APME AGAINST ESCHERICHIA COLI, PSEUDOMONAS AERUGINOSA, STREPTOCOCCUS AUREUS AND STAPHYLOCOCCUS EPIDERMIDIS




FIG. 7: ZONE OF INHIBITION OF VARIOUS BACTERIA FOR STANDARD, APEE AND APME

CONCLUSION: Whole plant of *A. pilosa* was extracted with ethyl acetate and methanol to obtain APEE and APME. Both APEE and APME was subjected to phytochemical screening and it was found that both contain flavonoids, alkaloids, tannins, glycosides, proteins and amino acids except for steroids and acid compounds which is only present in APME. TTC and TFC is more in APEE than APME. Both APEE and APME show antioxidant and antibacterial activity, APEE is claimed to have more antibacterial and antioxidant potency than APME.

REFERENCES:

- Mital Kaneria and Sumitra Chanda. Evaluation of antioxidant and antimicrobial properties of *Manilkara zapota* L. (chiku) leaves by sequential soxhlet extraction method. *Asian Pac J Trop Biomed* 2012; 2: S1526-33.
- Aruoma OI. Free radicals, oxidative stress, and antioxidants in human health and disease. *J Amer Oil Chem Soc* 1998; 75: 199-212.
- Jayaprakasha GK, Rao LJ, Sakariah KK. Antioxidant activities of flavin in different *in vitro* model systems. *Bioorg Med Chem* 2004; 12: 5141-6.
- Muniandy P, Shori AB, Baba AS. Influence of green, white and black tea addition on the probiotic yogurt during refrigerated storage. *Food Packaging and Shelf Life* 2016; 8:1-8.
- Vhangani LN, Wyk JV. Antioxidant activity of Maillard reaction products (MRPs) in a lipid rich model system. *Food Chem* 2016; 208: 301-308.
- Marles R. J. and Farnsworth N. R. Antidiabetic plants and their active constituents. *Phytomedicine* 1995; 2: 137-89.
- Bhandary M.J, Chandrashekar K.R and Kaveriappa K.M. Medical ethnobotany of the Siddis of Uttara Kannada district, India, *J Ethnopharmacol.* 1995; 47: 149-58.
- Abdolbaset Ghorbani, Gerhard Langenberger, Liu Feng and Joachim Sauerborn. Ethnobotanical study of medicinal plants utilised by Hani ethnicity in Naban River Watershed National Nature Reserve, Yunnan, China. *J Ethnopharmacol* 2011; 134: 651-67.
- Ameenah Gurih-Fakim, Mala Sewraj, Joseph Gueho and Ehsan Dulloo. *Medicinal ethnobotany of some weeds of Mauritius and Rodrigues*, *J Ethnopharmacol* 1993; 39: 175- 85.
- Galani V. J, Patel B. G, and Patel N. B. *Argyrea speciosa* (Linn. f.) sweet: A comprehensive review. *Pharmacogn Rev* 2010; 4:172-8.
- Lalan BK, Hiray RS and Ghongane BB. Evaluation of Analgesic and Anti-Inflammatory Activity of Extract of *Holoptelea Integrifolia* and *Argyrea Speciosa* in Animal Models. *J Clin Diagn Res* 2015; 9: FF01-4.
- Yadav KS, Yadav NP, Rawat B, Rai VK, Shanker K and Venkateswara Rao C. An assessment of wound healing potential of *Argyrea speciosa* leaves. *Scientific World J.* 2014; 1-6.
- Galani VJ & Patel BG. Psychotropic activity of *Argyrea speciosa* roots in experimental animals. *Ayu* 2011; 32: 380-4.
- Motawi TK, Hamed MA, Hashem RM, Shabana MH and Ahmed YR. Protective and therapeutic effects of *Argyrea speciosa* against ethanol-induced gastric ulcer in rats. *Z Naturforsch C* 2012; 67: 47-57.
- Sandeep kataria, Shaival Kamalaksha rao, Anil Bhandari and Dilsher Kaur. Pharmacognostical Standardization of *Corchorus depressus* (L.) Stocks (Tiliaceae)- A promising ethnomedicinal plant. *Ind J Tradit Knowl* 2013; 12: 489-97.
- Namrata Choudhary, MB Siddiqui and Sayyada Khatoun. Pharmacognostic evaluation of *Tinospora cardifolia* (Willd.) Miers and identification Markers. *Ind J Tradit Knowl* 2014; 13: 543-50.
- Milan S. Stanković. Total Phenolic Content, Flavonoid Concentration and Antioxidant Activity of *Marrubium Peregrinum* L. Extracts. *Kragujevac J. Sci.* 2011; 33:63-72.
- Akrouf A, Gonzalez LA, El Jani H, Madrid PC. Antioxidant and antitumor activities of *Artemisia campestris* and *Thymelaea hirsuta* from southern Tunisia. *Food Chem Toxicol.* 2011; 49: 342-7.
- Marinova. D, Ribarova. F, Atanassova. M. Total phenolics and total flavonoids in bulgarian fruits and vegetables. *J. Chem. Technol. Metall.* 2005; 40: 255-60.
- Rahate KP, Padma R, Parvathy NG, Renjith V. Quantitative estimation of tannins, phenols and antioxidant activity of methanolic extract of *Imperata cylindrical*. *Ind.J Pharm.Edu. Res.* 2013; 4: 73-7.
- Md. Nur Alam, Nusrat Jahan Bristi, Md. Rafiqzaman. Review on *in vivo* and *in vitro* methods evaluation of antioxidant activity. *Saudi Pharm J.* 2013; 21: 143-52.

22. Sur TK, Hazra A, Hazra AK, Bhattacharyya D. Antioxidant and hepatoprotective properties of Indian Sunderban mangrove *Bruguiera gymnorhiza* L. leave. J Basic Clin Pharma 2016; 7: 75-9.
23. Harpreet Walia, Subodh Kumar and Saroj Arora. Comparative analysis of antioxidant and phenolic content of chloroform extract/fraction of *Terminalia chebula*. J Basic Clin Pharma 2011; 2: 115-24.
24. Chitravadivu C, Manian S, Kalaichelvi K. Antimicrobial studies on selected medicinal plants, Erode region, Tamilnadu, India. Middle East J Sci Res 2009; 4: 147-52.
25. Nashwa T, Hassan HM, AbdelGhani SMM, Radwan IA, Hammouda O. Comparative chemical and antimicrobial study of nine essential oils obtained from medicinal plants growing in Egypt. Beni-Seuf Univ. J. Appl. Sci. 2014; 3: 149-56.
26. Shekhar TC, Anju G. Antioxidant activity by DPPH Radical Scavenging method of *Ageratum conyzoides* Linn. Leaves. Am J Ethnomed 2014; 1: 244-9.
27. Fernando CD, Soysa P. Optimized enzymatic colorimetric assay for determination of Hydrogen peroxide (H₂O₂) Scavenging activity of plant extracts. Methods X 2015; 2: 283-91.
28. Keser S, Celik S, Turkoglu S, Yilmaz O, Turkoglu I. Hydrogen peroxide radical scavenging and total Antioxidant Activity of Hawthorn. Chemistry Journal 2012; 2: 9-12.
29. Kumaran A, Karunakaran RJ. *In vitro* antioxidant activities of methanol extracts of five *Phyllanthus* species from India; LWT- Food, Science and Technology 2007; 40: 344-52.
30. Harborne JB, Williams CA. Advances in flavonoid research since 1992. Phytochemistry 2000; 55: 481-504.
31. Valgas C, Machado de Souza S, Smania EFA, Smania A. Screening methods to determine antibacterial activity of natural products. Braz J Microbiol 2007; 38: 369-80.
32. Huang P, Dong A, Caughey WS. Effects of Dimethyl sulfoxide, Glycerol and Ethylene glycol on secondary structures of Cytochrome C and Lysozyme as observed by Infrared spectroscopy. J Pharm Sci. 1995; 84(4): 387-92.
33. Zheng B, Li A, Jiang X, Hu X, Yao J, Zhao L, et al. Genome sequencing and genomic characterization of a tigeicycline-resistant *Klebsiella pneumoniae* strain isolated from the bile samples of a cholangiocarcinoma patient. Gut Pathogens 2014; 6: 1-7.
34. Ashok PK, Upadhaya K. Tannins are astringent. J Pharmacog Phytochem 2012; 1: 45-50.
35. Baskaran C, Rathabai V, Kanimozi D. Screening of antimicrobial activity and phytochemical analysis of various leaf extract of *Murraya koenigii*. Int J Res Ayurveda Pharm 2011; 2: 1807-10.
36. Taveria de Jesus NZ, Falcao HDS, Gomes IF, Leite TJDA, Lima GRDM, Babosa-Filho JM, et al. Tannins, peptic ulcer and related mechanisms. Int J Mol Sci 2012; 13: 3203-28.

How to cite this article:

Prasanth DSNBK, Rao AS and Yejella RP: Phytochemical, *in vitro* antioxidant and antibacterial activities of *Argyrea pilosa* wight & arn. (whole plant). *Int J Pharmacognosy* 2017; 4(4): 109-17. doi link: [http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.4\(4\).109-17](http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.4(4).109-17).

This Journal licensed under a Creative Commons Attribution-Non-commercial-Share Alike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)