



Received on 03 December 2016; received in revised form, 18 January 2017; accepted, 22 February 2017; published 01 March 2017

ANTIDIABETIC POTENTIAL OF IVY GOURD (*COCCINIA GRANDIS*, FAMILY: CUCURBITACEAE) GROWN IN SRI LANKA: A REVIEW

A. P. Attanayake*¹, K. A. P. W. Jayatilaka¹ and L. K. B. Mudduwa²

Department of Biochemistry¹, Department of Pathology², Faculty of Medicine, University of Ruhuna, Sri Lanka.

Keywords:

Antidiabetic potential,
Coccinia grandis, Standardization,
Toxicity studies

Correspondence to Author:

Dr. A. P. Attanayake

Department of Biochemistry,
Faculty of Medicine, University of
Ruhuna, Sri Lanka.

E-mail: anoja715@yahoo.com

ABSTRACT: Leaves of ivy gourd; *Coccinia grandis* (L.) Voigt (Family: Cucurbitaceae) is one of the well-known leafy vegetables, which has been consumed by Sri Lankans for centuries. The therapeutic potential of ivy gourd against diabetes is brought mainly by its antihyperglycemic, β -cell regenerative, antihyperlipidemic and antioxidant properties. Also, the optimum effective therapeutic dose of the aqueous *C. grandis* extract was found to be safe in terms of hepatotoxicity, renotoxicity, hematotoxicity *in-vivo*. The scientific investigations confirmed the effectiveness of this leafy vegetable for the development of potential nutraceuticals against diabetes mellitus.

INTRODUCTION: Diabetes mellitus has become a major global epidemic over the past few decades. The prevalence of diabetes has been increased worldwide and has reached alarming levels in many countries around the world. It is estimated that 415 million people (6.4% of the adult population worldwide) were affected by diabetes in 2015, and the number is projected to increase to 642 million in 2040. Accordingly, it is predicted to become the 7th leading cause of death in the world by the year 2040¹. In the past couple of decades, evidence from prospective observational studies and clinical trials has converged to support the importance of individual nutrients, foods, and dietary patterns in the prevention and management of diabetes mellitus.

Medicinal plants have long been used as medications or/and as simple dietary adjuncts to the existing therapies since antiquity. Before the introduction of insulin in 1922, the treatments for diabetes mellitus relied mainly on the recommendations of herbal preparations of edible medicinal plants². However, from the past to the present, the dietary recommendations for diabetes have widened, with the appreciation that fiber-rich food such as leafy vegetables would be able to enhance the glycemic control of patients with diabetes and thereby prevent the progression of the disease into diabetic complications^{3,4,5}. However, in this sense, the investigation of antidiabetic activity of edible, medicinal plants is imperative, as with proven *in-vivo* efficacy. These would be recommended as dietary adjuncts to the conventional therapies and more importantly can attribute for the development of potential sources of nutraceuticals.

A multitude of herbs, spices, and other plant materials have been described in the management of diabetes mellitus in Sri Lanka⁶.

	QUICK RESPONSE CODE DOI: 10.13040/IJPSR.0975-8232.IJP.4(3).79-83
	The article can be accessed online on www.ijpjournal.com
DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.4(3).79-83	

However, several edible, medicinal plants have been attributed to a special value in the diet of Sri Lankans. Ivy gourd; *Coccinia grandis* (L.) Voigt (Family: Cucurbitaceae) is one of the well-known leafy vegetables, which has been consumed by Sri Lankans for centuries. It is also known as a scarlet gourd in English; Kowakka, Kobowakka, Kemwell in Sinhala; Kovai, Kwai in Tamil. The plant is widely distributed in Southern, Western and North Central regions in the country. It is a perennial climber with single tendrils and glabrous leaves. Generally, the leaves have five lobes and are 6.5–8.5 cm long and 7–8 cm wide⁷. Even though the reputed effects of leaf extract of *C. grandis* have not been evaluated critically, it is widely used for the glycemic control by the local population. Accordingly, the results of a cross-sectional survey using an interviewer based questionnaire clearly showed that the leaves of *C. grandis* had been commonly used as a complementary and alternative medicine by patients with diabetes mellitus in Sri Lanka⁷.

Several scientific reviews are published on pharmacological properties of *C. grandis* grown in different geographical locations in the world; however, to date a comprehensive report on the antidiabetic potential of *C. grandis* of Sri Lankan origin has not been published. The present review aims to provide a comprehensive summary of the investigations carried out on the antidiabetic and toxicological effects of the leaves of Ivy gourd grown in Sri Lanka to corroborate its proposed antidiabetic potential in patients with diabetes mellitus.

Antihyperglycemic Activity: Hyperglycemia is the fundamental biochemical derangement in diabetes mellitus, causing oxidative and nitrosative stress, activation of inflammatory pathways and endothelial dysfunction, leading to vascular complications^{8, 9, 10}. However, effective blood glucose control is primarily targeted in the management of diabetes mellitus. Investigations carried out by Attanayake and coworkers^{11, 12} demonstrated that the leaf extract of *C. grandis* of Sri Lankan origin exerted both acute and long term antihyperglycemic effects in healthy, alloxan induced and streptozotocin-induced diabetic rats. The optimum therapeutic dose of the aqueous leaf extract of *C. grandis* upon a glucose load was

0.75g/kg in healthy and diabetic rats. Leaf extract of *C. grandis* at the optimum effective dose lowered the blood glucose concentration gradually during four hours in diabetic rats. The highest reduction (33%) in the blood glucose concentration was observed in the second hour after the administration of *C. grandis* extracts in streptozotocin-induced diabetic rats. Also, Munasinghe and co-workers¹³ reported a phase 1 clinical trial of an ingestion of a meal containing 20 g of leaves of *C. grandis* mixed with a measured amount of scraped coconut and table salt were able to show acute hypoglycemic effects in a selected group of healthy subjects.

The acute antihyperglycemic activity of *C. Grandis* may be due to a reduction in the absorption of glucose in the small intestine, stimulation of glucose-induced secretion of insulin from the pancreatic β -cells, enhancement of uptake of glucose by peripheral tissues, etc.¹⁴ The long term antihyperglycemic activity is also evident in *C. grandis* treated diabetic rats. The diabetic rats treated with the optimum therapeutic dose of the aqueous extract of *C. grandis* exhibited a remarkable glycemic control as evident by a reduction in the percentage of HbA_{1c} in streptozotocin-induced diabetic rats¹⁵. Accordingly, the concentration of fasting serum fructosamine, insulin and C-peptide were decreased significantly in diabetic rats. The results reported by Attanayake et al.,¹⁵ corroborate with the findings on antihyperglycemic efficacy of *C. grandis* grown in India^{16, 17, 18}.

β -cell Regenerative Potency: The optimum pancreatic β -cell function is essential for the regulation of intracellular glucose homeostasis. Several studies have provided evidence that loss of functional β -cell mass through apoptosis and impaired proliferation consequent to hyperglycemia is considered as a hallmark of diabetes mellitus¹⁹. Regulation of functional β -cell mass has been considered as a critical therapeutic challenge in patients with the disease²⁰. However, islet cell regeneration has gained much interest and has been considered as a strategy to restore the loss of β -cell mass in diabetes mellitus. Stimulation of β -cell proliferation/regeneration may have an impact on the insulin-producing cells in the pancreas²¹. One such approach to foster restoration and regeneration

of β -cells is from putative exogenous sources as edible medicinal plants²². Accordingly, the extract of *C. grandis* of Sri Lankan origin showed β -cell regeneration as evident through an increase in the number of insulin secreting β -cells in streptozotocin-induced diabetic rats. An increase in the diameter of small, medium and large islets was found in *C. grandis* treated diabetic rats, and it was confirmed that the functional mass of the islet and the entire regenerative capacity of the pancreas increased with the treatment of *C. grandis* extract in diabetic rats¹⁵. In contrast, none of the species of *C. grandis* except the variety found in Sri Lanka reported the β cell regenerative potency *in vivo* to date.

Antihyperlipidemic Activity: Lipid abnormalities are postulated to be major causes of complications associated with diabetes mellitus. Indeed, dyslipidemia in diabetes mellitus is generally comprised of low levels of high-density lipoprotein cholesterol (HDL-C), high levels of low-density lipoprotein cholesterol (LDL-C) and triglycerides (TG)²³. Several studies have revealed that a positive correlation between dyslipidemia and development of premature atherosclerosis, coronary insufficiency and myocardial infarction in diabetic subjects^{9, 22}. Thus, the effect of natural products in the management of diabetic complications has recently received considerable attention, highlighting the importance of edible leafy vegetables as regulators of metabolism of carbohydrate and lipids²⁴.

Recent studies by Attanayake and coworkers have demonstrated that a significant reduction in the concentration of serum TC, TG, LDL-C and an elevation in the concentration of HDL-C in streptozotocin-induced diabetic rats on the 30th day of study¹⁵. This is similarly mentioned by other authors who observed potential antihyperlipidemic effects in serum lipid parameters in diabetic rats with the leaf extract of *C. grandis* of Indian origin^{25, 26}. The favorable effects of aqueous leaf extract of *C. grandis* on serum lipid parameters in addition to the glycemic control would be useful in the development of nutraceuticals targeting diverse disease pathologies of diabetes mellitus. The increase in the concentration of HDL-C by *C. grandis* is of the therapeutic advantage as it improves the glycemic control as well as reduces

the cardiovascular risk associated with diabetes mellitus. The strong antihyperlipidemic effect by the administration of *C. grandis* in diabetic rats could be through its control on hyperglycemia or/and direct actions on the absorption of intestinal cholesterol and biosynthesis of cholesterol as suggested by many authors^{26, 27}.

Antioxidant Activity: Oxidative stress is one of the major factors underlying the resistance for insulin in disease pathogenesis of diabetes mellitus²⁸. Elevated free fatty acids, glycation of proteins, other circulating factors, leptin and the simultaneous decline of cellular antioxidant defense mechanisms in patients with diabetes contribute to causing an overproduction of reactive oxygen species⁹.

Edible, medicinal plant extracts with strong antioxidative properties have been the recent focus in diminishing cellular oxidative stress in diabetes mellitus. The increase in reactive oxygen species may lead to disruption of cellular functions, oxidative damage to membranes and lipid peroxidation that consequently lead to the occurrence of diabetic complications. Also, it has been shown that dietary supplementation with natural antioxidants such as Vitamins C and E, melatonin, and flavonoids attenuates the oxidative stress in diabetic subjects²⁹.

The overall experimental data reported by Attanayake et al.,³⁰ indicated that the administration of leaf extract of *C. grandis* was able to improve the activities of antioxidant enzymes such as glutathione reductase (EC 1.6.4.2), glutathione peroxidase (EC 1.11.1.9), glutathione S-transferase (EC 2.5.1.18), increase the concentration of reduced glutathione and decrease the concentration of malonaldehyde as a product in the lipid peroxidation in diabetic rats.

Also, the *in-vitro* glycation induced protein cross-linking inhibitory effects of the leaf extract of *C. grandis* was also proven³¹. The *in-vivo* antioxidant activity is further corroborated by the relatively high *in-vitro* antioxidant activity of *C. grandis* compared determined using three *in-vitro* assays of 2,2-diphenyl-2-picrylhydrazyl hydrate (DPPH) radical scavenging activity, ferric reducing power (FRAP) and nitric oxide (NO) inhibition³².

The *in-vivo* antioxidative activity of *C. grandis* may be due to the presence of a wide range of phytochemicals principally due to the high content of polyphenolic and flavonoid compounds as suggested by Attanayake et al.³²

Safety Profile:

Chemical Standardization: Standardization of crude plant extracts involves the process of prescribing a set of standards or inherited characteristics, constant parameters, definitive quality and quantitative values that carry an assurance of quality, safety and reproducibility³³. The *C. grandis* grown in Sri Lanka has a very low amount of acid insoluble ash content, and a moderate level of total ash content indicates the purity of the leaves of *C. grandis* leaves that could be useful in human consumption and development of nutraceutical products. According to the recent report by Attanayake et al.,³⁴ heavy metals were not detected in *C. grandis* leaves of Sri Lankan origin.

***In-vivo* Toxicity Studies:** The use of edible, medicinal plants for therapeutic purposes has been increasingly popular as they are believed as beneficial and free of side effects^{35, 36}. However, the rationale for the utilization of plant food has rested largely on long-term clinical experience with little or no scientific data on safety. Attanayake and coworkers reported that the administration of *C. grandis* at a range of doses (0.25 g/kg-2.00 g/kg) in healthy Wistar was well tolerated by test animals, suggesting it's *in-vivo* safety. Further, the long term administration of the extract to Wistar rats for 28 days was not associated with adverse effects reflected in the general condition, growth, relative weight of organs, clinical biochemical, hematological values and more importantly did not cause abnormalities in the histopathology of body tissues³⁷. However, a proper clinical evaluation may be required to define a safe dosage in humans precisely.

CONCLUSION: This is the first scientific review that describes the potential antidiabetic activities and toxicological effects of *C. grandis* grown in Sri Lanka. The fact that the aqueous leaf extract of *C. Grandis* exerts antihyperglycemic, antihyperlipidemic, antioxidant activities *in-vivo* merits its use in the development of potential nutraceuticals

against diabetes mellitus. However, there is an urgent need for the determination of its clinical efficacy on diabetic subjects through well-defined and adequately powered randomized controlled clinical trials before the recommendation of ivy gourd as an effective dietary supplement for patients with diabetes mellitus.

ACKNOWLEDGEMENT: Nil

CONFLICT OF INTEREST: Nil

REFERENCES:

1. Zimmet PZ, Magliano DJ, Herman WH and Shaw JE: Diabetes: A 21st-century challenge. *Lancet Diabetes Endocrinology* 2014; 2(1): 56-64.
2. Bhattacharya M: A historical exploration of Indian diets and a possible link to insulin resistance syndrome. *Appetite* 2015; 95: 421-454.
3. Cho SS, Qi L, Fahey GCr and Klurfeld DM: Consumption of cereal fiber, mixtures of whole grains and bran, and whole grains and risk reduction in type 2 diabetes, obesity, and cardiovascular disease. *American Journal of Clinical Nutrition* 2013. <http://dx.doi.org/10.3945/ajcn.113.067629>.
4. Post RE, Mainous AG, King DE and Simpson KN: Dietary fiber for the treatment of type 2 diabetes mellitus: A meta-analysis. *Journal of American Board of Family Medicine* 2012; 25 (1): 16-23.
5. Ley SH, Hamdy O, Mohan V and Hu FB: Prevention and management of type 2 diabetes: Dietary components and nutritional strategies. *The Lancet* 2014; 383(9933): 1999-2007.
6. Ediriweera ERHSS and Ratnasooriya WD: A review on herbs used in the treatment of diabetes mellitus by Sri Lankan Ayurvedic and Traditional Physicians. *Ayu* 2009; 30: 373-391.
7. Medagama AB, Bandara R, Abeysekera RA, Imbulpitaya B and Pushpakumari T: Use of complementary and alternative medicines (CAMs) among type 2 diabetes patients in Sri Lanka: A cross-sectional survey. *BMC Complementary and Alternative Medicine* 2014. <http://doi.org/10.1186/1472-6882-14-374>.
8. Jayaweera DMA: Medicinal Plants (indigenous and exotic) used in Ceylon. 2nd ed. Sri Lanka, Colombo 1982.
9. Forbes JM, Cooper ME: Mechanisms of diabetic complications. *Physiological Reviews* 2011; 93(1): 137-188.
10. ADA (American Diabetes Association). Clinical practice recommendations *Diabetes Care* 2016; 20(30): S16.
11. Attanayake AP, Jayatilaka KAPW, Pathirana C and Mudduwa LKB: Study of antihyperglycaemic activity of some Sri Lankan medicinal plants in alloxan-induced diabetic rats. *Ancient Science of life* 2013; 32(4): 193-8.
12. Attanayake AP, Jayatilaka KAPW, Pathirana C and Mudduwa LKB: Acute hypoglycemic and antihyperglycemic effects of ten Sri Lankan medicinal plant extracts in healthy and streptozotocin-induced diabetic rats. *International Journal of Diabetes in Developing Countries* 2014. [doi.org/10.1007/s13410-014-0217-8](http://dx.doi.org/10.1007/s13410-014-0217-8).
13. Munasinghe MAAK, Abeysena C, Yaddhegige IS, Vidanapathirana T and Piyumal KPB: Blood sugar lowering effect of *Coccinia grandis* (L.) J. Voigt: Path for

- a new drug for diabetes mellitus. *Experimental Diabetes Research* 2011. <http://dx.doi.org/10.1155/2011/978762>.
14. Ndong M, Uehara M, Katsumata S and Suzuki K: Effects of oral administration of *Moringaoleifera* Lam on glucose tolerance in Goto-Kakizaki and Wistar Rats. *Journal of Clinical Biochemistry and Nutrition* 2007; 40(3): 229-233.
 15. Attanayake AP, Jayatilaka KAPW, Pathirana C and Mudduwa LKB: Antihyperglycemic activity of *Coccinia grandis* (L.) Voigt (Cucurbitaceae) in streptozotocin-induced diabetic rats. *Indian Journal of Traditional Knowledge* 2015; 14(3): 376-381.
 16. Pekamwar SS, Kalyankar TM and Kokate SS: Pharmacological Activities of *Coccinia grandis*: Review. *Journal of Applied Pharmaceutical Science* 2013; 3(05): 114-119.
 17. Deokate UA and Khadabadi SS: Pharmacology and phytochemistry of *Coccinia indica* Pharmacophore 2012; 3(3): 179-185.
 18. Balaraman AK, Singh J, Dash S and Maity TK: Antihyperglycemic and hypolipidemic effects of *Melothria maderaspatana* and *Coccinia indica* in streptozotocin-induced diabetes in rats. *Saudi Pharmaceutical Journal* 2010; 18(3): 173-178.
 19. Ashcroft FM and Rorsman P: Diabetes mellitus and the beta cell: the last ten years. *Cell* 2012; 148: 1160-1171.
 20. Ferrannini E: The stunned beta cell: a brief history. *Cell Metabolism* 2010; 11: 349-352.
 21. Hosseini A, Shafiee-Nick R and Ghorbani A: Pancreatic beta cell protection/regeneration with phytotherapy. *Brazilian Journal of Pharmaceutical Sciences* 2015; 51: 1-16.
 22. Vetere A, Choudhary A, Burns SM and Wagner BK: Targeting the pancreatic β -cell to treat diabetes. *Nature Reviews Drug Discovery* 2014; 14: 278 -289.
 23. Tangvarasittichai S: Oxidative stress, insulin resistance, dyslipidemia and type 2 diabetes mellitus. *World Journal of Diabetes* 2015; 6(3): 456-480.
 24. Farzaei MH, Rahimi R, Farzaei F and Abdollahi M: Traditional medicinal herbs for the management of diabetes and its complications: An evidence-based review. *International Journal of Pharmacology* 2015; 11: 874-887.
 25. Mohammed SI, Chopda MZ, Patil RH, Vishwakarma KS and Maheshwari VL: *In-vivo* antidiabetic and antioxidant activities of *Coccinia grandis* leaf extract against streptozotocin-induced diabetes in experimental rats *Asian Pacific Journal of Tropical Disease* 2016; 6(4): 298-304.
 26. Krishnakumari S, Bhuvaneshwari P and Rajeswari P: Ameliorative potential of *Coccinia grandis* extract on serum and liver marker enzymes and lipid profile in streptozotocin-induced diabetic rats. *Anc Sci Life* 2011; 31(1): 26-30.
 27. Manjula S and Ragavan B: Hypoglycemic and hypolipidemic effect of *Coccinia indica* Wight and Arn in alloxan-induced diabetic rats. *Ancient Science of Life* 2007; 27(2): 34-37.
 28. Tiwari BK, Pandey KB, Abidi AB, Rizvi SI: Markers of oxidative stress during diabetes mellitus. *Journal of Biomarkers* 2014. <http://dx.doi.org/10.1155/2013/378790>.
 29. Kurutas EB: The importance of antioxidants which play a role in cellular response against oxidative/nitrosative stress. *Nutrition Journal* 2015. doi: <http://10.1186/s12937-016-0186-5>.
 30. Attanayake AP, Jayatilaka KAPW, Mudduwa LKB and Pathirana C: *In vivo* antihyperlipidemic, antioxidative effects of *Coccinia grandis* (L.) Voigt (Cucurbitaceae) leaf extract: An approach to scrutinize the therapeutic benefits of traditional Sri Lankan medicines against diabetic complications. *International Journal of Pharmaceutical Sciences and Research* 2016; 7(10): 3949-3958.
 31. Perera HKI and Handuwalage CS: Analysis of glycation induced protein cross-linking inhibitory effects of some antidiabetic plants and spices. *BMC Complementary and Alternative Medicine* 2015. <http://doi.10.1186/s12906-015-0689-1.7>.
 32. Attanayake AP, Jayatilaka KAPW, Pathirana C and Mudduwa LKB: Phytochemical screening and *in-vitro* antioxidant potentials of extracts of ten medicinal plants used for the treatment of diabetes mellitus in Sri Lanka. *African Journal of Traditional Complementary and Alternative Medicines* 2015b; 12(4): 28-33.
 33. Kunle OF, Egharevba HO and Ahmadu PO: Standardization of herbal medicines - A review *International Journal of Biodiversity and Conservation*. 2012; 4(3): 101-112.
 34. Attanayake AP, Arawwawala LDAM and Jayatilaka KAPW: Chemical standardization of leaf extract of *Coccinia grandis* (L.) Voigt (Cucurbitaceae) of Sri Lankan origin, *Journal of Pharmacognosy and Phytochemistry* 2016; 5(5): 119-123.
 35. Ekor E: The growing use of herbal medicines: Issues relating to adverse reactions and challenges in monitoring safety. *Frontiers in Pharmacology* 2013. <http://doi:10.3389/fphar.2013.00177>.
 36. Jordan SA, Cunningham DG and Marles RJ: Assessment of herbal medicinal products: Challenges and opportunities to increase the knowledge base for safety assessment. *Toxicology and Applied Pharmacol* 2010; 243(2): 198-16.
 37. Attanayake AP, Jayatilaka KAPW, Pathirana C and Mudduwa LKB: Efficacy and toxicological evaluation of *Coccinia grandis* (Cucurbitaceae) extract in male Wistar rats. *Asian Pacific Journal of Tropical Disease* 2013(b); 3(6): 460-466.

How to cite this article:

Attanayake AP, Jayatilaka KAPW and Mudduwa LKB: Antidiabetic potential of ivy gourd (*Coccinia grandis*, Family: Cucurbitaceae) grown in Sri Lanka: a review. *Int J Pharmacognosy* 2017; 4(3): 79-83. doi link: [http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.4\(3\).79-83](http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.4(3).79-83).

This Journal licensed under a Creative Commons Attribution-Non-commercial-Share Alike 3.0 Unported License.

This article can be downloaded to **Android OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)