



Received on 08 March 2017; received in revised form, 21 May 2017; accepted, 30 May 2017; published 01 June 2017

A NEW NEO-CLERODANE DITERPENOID FROM *TEUCRIUM POLIUM* LINN.

M. W. M. Sadaka

Department of Chemistry, Applied Science Faculty, University of Kalamoon, Deir Atteia, Syria.

Keywords:

Teucrium polium L.,
neo-clerodane, Diterpenoid,
Cirsimartin, Cirsiliol, Apigenin

Correspondence to Author:

M. W. M. Sadaka

Department of Chemistry,
Applied Science Faculty, University
of Kalamoon, Deir Atteia, Syria.

E-mail: waleed.sadaka@gmail.com

ABSTRACT: *Teucrium polium* L. is a member of the Lamiaceae family and is represented in the Flora of Syria by two variety: *Teucrium polium* var. *angusti folium* and *Teucrium polium* var. *mollissimum*. *Teucrium polium* is a wild-growing flowering plant, found abundantly in Syria, and it is used traditional medicine for its diuretic, antipyretic, antispasmodic, tonic, anti-inflammatory, antihypertensive, anorexic, analgesic, antibacterial and antidiabetic effects. Several reports have demonstrated a wide range of beneficial biological and pharmacological activities of the phenylethanoid, phenylpropanoid, and flavonoid components, while the furano neo-clerodane diterpenoids present in germander have been implicated in the *in-vivo* hepatotoxicity of this botanical. Phytochemical studies of this plant have been carried out. Aerial parts of the plant were extracted with petroleum ether at room temperature, and then extract with methanol in Soxhlet, and finally extract with aqueous methanol at room temperature successively. Fractionation of the methanol extract by column chromatography and purification by crystallization yielded three known flavonoids Cirsimartin 2, Cirsiliol 3, and Apigenin 4. The aqueous methanolic extract yielded one new neo-clerodane type diterpenoid, Syrapolin I 1 (3,12- diacetoxy - 4 α ,18 α ; 15,16- diepoxy - 6 - oxo - neo - cleroda - 13 (16), 14-diene -7 α , 20 β - dihydroxy-19- hemiacetal). The structure of compounds was proposed based on spectroscopic methods IR, MS, and 1-D (¹H, ¹³C, and DEPT) and 2-D (COSY, HETCOR, HMBC) NMR experiment, and comparison with closely related compounds.

INTRODUCTION: *Teucrium polium* (Family Lamiaceae) is one of the popular fragrant plants in Syria and is distributed throughout the country. In the Flora of Syria, this genus is represented by 21 species such as *T. creticum* grow in Safita and Mussiaf, *T. oliverianum* grows in Deir-ez-Zor, *T. chamaedrys* grow in Lattakia and Baniyas, and *T. polium* grow in Al-Hassaka. *Teucrium polium* is a perennial shrub, 20-50 cm high, distributed widely in the dry and stony hills and deserts of almost all Mediterranean countries, South-western Asia, Europe, and North Africa ¹.

The bruised foliage releases a pleasant aromatic odor and the flowers are small, in clusters and range from pink to white. In the folk medicinal traditions, this plant is used for its antibacterial, anti-spasmodical, anti-rheumatic, anti-diabetic and antipyretic activities. In an admixture with other powdered herbs, it is claimed to be therapeutic for peptic ulcer.

The biological activity of *T. polium* is widely reported, and it has been shown to possess anti-inflammatory ², anti-nociceptive ^{3, 4} activities. The aqueous extract of *Teucrium polium* aerial parts, given intraperitoneally at dosages from 50 to 150 mg/kg for 10 days, reduced significantly the serum levels of cholesterol and triglycerides in hyperlipidemic rats ⁵. Recently, the high insulinotropic and anti-hyperglycemic activity of its crude extract using both animal and isolated rat pancreatic islets has been evaluated ⁶⁻⁸.

	<p>QUICK RESPONSE CODE</p>
	<p>DOI: 10.13040/IJPSR.0975-8232.IJP.4(6).185-92</p>
<p>The article can be accessed online on www.ijournal.com</p>	
<p>DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.4(6).185-92</p>	

Antibacterial activity against some Gram-positive and Gram-negative bacteria as well as antifungal activity against *Yarrowia lipolitica* and *Saccharomyces cerevisiae* has also been reported^{9, 10}. Recently, cytotoxic, anticancer, and antimutagenic effects of ethanol and aqueous extracts of *T. polium* have been shown on various cell lines¹¹⁻¹³. *T. polium* extract contains selenium¹⁴ and has shown antioxidant activity like α -tocopherol. In particular, their antioxidant activities were evaluated through the use of several commonly accepted in vitro assays, which included DPPH scavenging (DPPH), reducing power (RP), xanthine oxidase inhibitor effect (XOI) and antioxidant activity in a linoleic acid system (ALP)¹⁵⁻²¹.

The few adverse effects of *T. polium* have been reported which indicate the relatively safe nature of this medicinal herb²²⁻²⁷. The composition of the essential oil of *T. polium* has been the subject of several investigations²⁸⁻³⁶. The naturally occurring *neo*-clerodane diterpenoids have attracted interest because of their biological activity as insect antifeedant and as antifungal, anti-tumor, and antimicrobial agents. Although a large number of these compounds have been isolated from many plants in the last few years, the genus *Teucrium* is one of the richest sources of *neo*-clerodane diterpenes: more than 230 diterpenes have been described. Phytochemical investigations of *T. polium* have been shown that it contains a lot of diterpenes belong to *neo*-clerodane type³⁷⁻⁴⁵.

T. polium also rich in constituents such as flavonoids, iridoids and Phenylpropanoid glycosides with undoubted antioxidant properties^{18, 46-50}. Our previous work on the aerial parts of *T. polium* L. var. *mollissimum* Hand-Mazz. led to the isolation of one new *neo*-clerodane diterpenes Syrapolin II, and two known compounds 6'-O-caffeoyl-8-O-acetylharpagide, clerosterol-3-O-glucoside⁵¹. In continuation of our studies on *neo*-clerodane diterpenoids from the *Teucrium* species, we have investigated *T. podium*, a small shrub which grows widely in Syria. From the methanolic extract and aqueous methanolic extract of the aerial parts of this plant we have isolated the new *neo*-clerodane diterpenoids Syraoiln I 1, beside to three new known flavonoids, Cirsimartin 2, Cirsiliol 3, and Apigenin 4.

MATERIALS AND METHODS:

Plant Material: *Teucrium polium* L. Amouda, Al-Hassaka, Syria in April 2008, and identified by Pro. Dr. Anwer Al-Khateeb, Department of Botany, College of Science, Damascus University, Damascus, Syria. A voucher specimen is available in the herbarium of Department of Botany in Damascus University.

Chemicals: Silica gel 60 F₂₅₄ plates (Merck) were used for analytical and preparative TLC. Silica gel S (Merck; Mesh 230-400) was used for column chromatography (CC), and all the chemicals were purchased from Merck.

Instrumentation: The ¹H and ¹³C NMR spectra were recorded on Bruker Ultra Shield 400 MHz spectrometer (Faculty of Science, Al-Baath University, Homs, Syria) and reported in ppm. (δ) relative to TMS as internal standard and DMSO-*d*₆ as a solvent. The IR spectra were measured on JASCO (FT-IR)-410 spectrophotometer (Medico Medical Company, Homs, Syria). MS spectra were recorded on Shimadzu GCMS-QP 2010 (Syrian atomic energy commission, Damascus, Syria).

Experimental:

Extraction and Isolation: Dried and finely powdered *Teucrium polium* L. aerial parts (4 kg) were extracted with petroleum ether at room temperature for 2 days, then filtrate and the residual plant re-extracted with methanol in soxhlet. The methanolic extract (180 g) evaporated under reduced pressure and then extracted with diethyl ether. The diethyl ether extract (8g) was chromatographed on silica gel column and eluted with n-hexane, n-hexan-CHCl₃ (90:10), n-hexan-CHCl₃ (50:50), gradient CHCl₃-MeOH, and pure MeOH, and the fractions collected as 25 ml. Elution with CHCl₃-MeOH (90:10) afford 160 fractions. Fractions (92-100) gave 71 mg of an impure compound which were crystallized in dichloromethane to give 38 mg of compound 2, and Fractions (122-124) gave 64 mg of an impure compound which were crystallized in dichloromethane to gave 29 mg of compound 3. Elution with CHCl₃-MeOH (80:20) gave 48 mg residues which were crystallized in a mixture of acetone-methanol to gave 22 mg of pure compound 4. The plant residue from methanolic extract then re-extracted with the mixture of MeOH-H₂O

(70:30) at room temperature, and the filtrate was evaporated in vacuum, and during this evaporation, the white precipitate was accumulated, we filtrate it and washed with methanol to gave 80 mg of pure compound 1.

RESULTS AND DISCUSSION: Syrapolin I (1) (Fig. 1) gave a molecular ion peak in its EI-mass spectrum at m/z 478, in agreement with the

molecular formula $C_{24}H_{30}O_{10}$. The other prominent peaks were found to occur at m/z 460 $[M-H_2O]^+$, 418 $[M-AcOH]^+$, 135, 122, 105(100), 94, 91, 81, 77, 55, 43. The IR spectrum of (1) showed absorptions of furan ring (3141.5, 3124.1, 1504.2, 874.6 cm^{-1}), hydroxyl (3381.5 cm^{-1} , br), tow acetates (1754.9, 1731.8, 1280.5, 1239.0 cm^{-1}), and a strong band for a carbonyl group (1713.4 cm^{-1}).




FIG. 1: STRUCTURE OF COMPOUNDS 1, 2, 3, 4

The data from the ^{13}C -NMR spectrum **Table 1** display **24** signals. The multiplicities of carbon signals were determined by the DEPT experiment. Taking into account the results from our comprehensive 1D and 2D-NMR studies and previous knowledge derived from metabolites isolated from the genus *Teucrium*^{39, 51-53}. It is evident that (1) possesses a highly oxidized *neo*-clerodane nucleus. The 1H -NMR spectrum (Table 1) of compound (1) showed *add* at δ 6.56 ($J = 0.8$, 1.6 Hz, H-14), a broad singlet at δ 7.71 (H-15), and a triplet at δ 7.63 ($J = 1.7$ Hz, H-16) were readily assigned to a terminal furan ring.

Also, the 1H -NMR spectrum showed two AB system (δ 2.95, 4.14 and 2.96, 2.81) which we have assigned to H-19 and H-18 respectively. The broad doublet at δ 5.83 and the doublet at 4.87 were assigned to H-12 and H-20, respectively. The 1H -NMR spectrum also showed two doublets at δ 5.03 (5.6 Hz), 6.47 (4.2 Hz) which were not connected to any carbon atoms in HETCOR spectrum were assigned to two hydroxyl group, and disappeared from 1H -NMR spectrum when recorded in DMSO

with few drops of D_2O . Detailed examination of 1H -NMR spectrum indicated the presence of 1H - 1H spin system which could be traced from the secondary methyl protons δ 1.09 (*d*, $J = 6.8$ Hz, 3H-17) to δ 1.66 (*q*, $J = 6.4$ Hz, H-8) and from there to an oxymethine proton at δ 4.25 (*dd*, $J = 5.6$, 9.9 Hz, H-7). Mutual coupling with the signal at δ 5.03 (*d*, $J = 5.6$ Hz, OH) suggested by the common J value, was confirmed by a cross peak in the 1H -NMR, 2D COSY spectrum, and supported the presence of a hydroxyl group at C-7 with α configuration (equatorial). As expected, no cross peak was observed in the HETCOR spectrum for a carbon resonance and proton signal at δ 5.03, but in the HMBC spectrum, cross peaks were observed for C-6, C-8, and C-7. The HMBC spectrum also showed cross peaks between the signal at δ 6.47 (*d*, $J = 4.2$ Hz, C20-OH) and C-20, C-9, and C-10. The 1H -NMR spectrum also shows tow singlet at δ 1.90, and 2.03 indicate to the tow CH_3 in acetate groups.

The ^{13}C -NMR **Table 1** and DEPT spectra confirmed the presence of a furan ring (δ 126.70,

C-13; 109.38, C-14; 140.37, C-15 and 144.31, C-16), a carbonyl group (δ 208.23, C-6), two acetate (δ 169.90 COMe; 21.42, COMe, and 170.24 COMe; 21.70, COMe), one methyl signal (δ 13.68, C-17) and two oxygen-substituted methylene signals for the C-18 and C-19 at δ 52.39 and 54.00 respectively, three oxygen-substituted methine

signals for the C-3, C-7, C-12 and C-20 at δ 72.97, 78.85, 65.23 and 93.29 respectively. On the basis of these facts, the structure of (1) was established as 3, 12-diacetoxy-4 α , 18 α ; 15, 16-diepoxy-6-oxo-*neo*-cleroda-13(16),14-diene-7 α ,20 β -dihydroxy-19-hemiacetal.

TABLE 1: ^{13}C -NMR AND ^1H -NMR DATA OF 1 (400 MHz, DMSO- D_6)

S. no.	C	DEPT	HETCOR
1	19.05	CH ₂	2.17 <i>m</i> (5.2 Hz)
2	26.87	CH ₂	a 1.62 <i>brd</i> (6.4 Hz) b 2.14 <i>m</i> (5.2 Hz)
3	72.97	CH	4.29 <i>d</i> (6.1 Hz)
4	55.93	C	
5	48.20	C	
6	208.23	C	
7	78.85	CH	4.25 <i>dd</i> (5.6, 9.9 Hz)
8	46.90	CH	1.66 <i>q</i> (6.4 Hz)
9	41.12	C	
10	40.77	CH	2.20 <i>d</i> (4.4 Hz)
11	35.73	CH ₂	a 1.56 <i>dd</i> (16.2, 2.4 Hz) b 2.45 <i>dd</i> (12, 16 Hz)
12	65.23	CH	5.83 <i>brd</i> (8 Hz)
13	126.70	C	
14	109.38	CH	6.56 <i>dd</i> (0.8, 1.6 Hz)
15	140.37	CH	7.71 <i>brs</i>
16	144.31	CH	7.63 <i>t</i> (1.7 Hz)
17	13.68	CH ₃	1.09 <i>d</i> (6.8 Hz)
18	52.39	CH ₂	a 2.81 <i>d</i> (4.7 Hz) b 2.96 <i>d</i> (4.7 Hz)
19	54.00	CH ₂	a 2.95 <i>d</i> (12.3 Hz) b 4.14 <i>d</i> (12.4 Hz)
20	93.29	CH	4.87 <i>d</i> (4.3 Hz)
CH ₃ CO	21.42	CH ₃	1.88 <i>s</i>
CH ₃ CO	21.70	CH ₃	2.04 <i>s</i>
CH ₃ CO	169.90		
CH ₃ CO	170.24		
		OH	5.03 <i>d</i> (5.6 Hz)
		OH	6.47 <i>d</i> (4.2 Hz)

Compound (2) was obtained as a yellow granular powder and exhibiting a positive ferric chloride test and magnesium hydrochloric acid test. Compound (2) gave a molecular ion peak in its EI-mass spectrum at m/z ; M = 314 (100%), and M+1 (11%) in agreement with the molecular formula C₁₇H₁₄O₆. IR spectrum of (2) showed absorption band for hydroxyl groups 3300 cm⁻¹, α , β -unsaturated carbonyl carbon 1640 cm⁻¹, and aromatic double bond 1595 cm⁻¹ functionalities. The UV spectrum of (2) exhibited absorption maxima at 331 (Band I) and 275 nm (Band II) (in MeOH), bathochromic shifts of 67, 3 and 4 nm with NaOMe, NaOAc and AlCl₃ in Band I, respectively. These data revealed that (2) has a flavonoid type structure.

The ^1H -NMR spectrum of (2) **Table 2** showed a characteristic low-frequency signal at δ 12.90 (*s*, 5-OH) assignable to aromatic hydroxyl signal hydrogen bonded to carbonyl at C-5 of a flavonoid. Two *ortho*-coupled aromatic protons at δ 7.92 (1H, *d*, $J = 8.1$ Hz) and δ 6.92 (^1H , *d*, $J = 8.1$ Hz) were diagnostic for a C-2'/C-6' and C-3'/C-5' oxygenated ring B. The ^1H NMR spectrum showed two aromatic proton signals at (δ 6.86, *s*, H-8) and (δ 6.80, *s*, H-3). The ^1H -NMR also showed two singlet signal at δ 3.72 (*s*, 3H) and δ 3.90 (*s*, 3H) indicate the presence of two methoxy group. In ^{13}C -NMR spectra **Table 2** and DEPT-135, the signals at δ 164.53, 103.15, 182.10, 152.58 and 105.56 were typical of C-2, C-3, C-4, C-9 and C-10

of a flavonoid moiety. The position of two methoxy groups was authenticated through the HMBC spectrum between methoxy protons at δ 3.72 and C-6, and δ 3.90 and C-7. By comparison of ^1H - and

^{13}C -NMR chemical shifts with those reported in the literature, the flavonoid skeleton were in agreement to those of cirsimartin⁵⁴.

TABLE 2: ^{13}C -NMR AND ^1H -NMR DATA OF COMPOUND (2), (400 MHZ, DMSO- D_6)

No	2		3		4	
	C	H	C	H	C	H
2	164.53	---	164.77	---	164.77	---
3	103.15	6.80, 1H, <i>s</i>	103.18	6.71, 1H, <i>s</i>	103.32	6.76, 1H, <i>s</i>
4	182.10	---	183.64	---	182.25	---
5	153.11	---	153.13	---	164.23	---
6	132.33	---	132.35	---	99.44	6.18, 1H, <i>d</i> , <i>J</i> = 2.09 Hz
7	159.09	---	159.11	---	164.95	---
8	92.00	6.86, 1H, <i>s</i>	91.95	6.85, 1H, <i>s</i>	94.54	6.48, 1H, <i>d</i> , <i>J</i> = 2.09 Hz
9	152.58	---	152.57	---	157.86	---
10	105.56	---	105.54	---	104.14	---
1'	121.61	---	121.94	---	121.68	---
2'	129.01	7.92, 1H, <i>d</i> , <i>J</i> = 8.1 Hz	113.94	7.45, 1H, <i>s</i>	128.99	7.92, 1H, <i>d</i> , <i>J</i> = 8.9 Hz
3'	116.47	6.92, 1H, <i>d</i> , <i>J</i> = 8.1 Hz	146.30	---	116.50	6.92, 1H, <i>d</i> , <i>J</i> = 8.9 Hz
4'	161.79	---	150.38	---	161.99	---
5'	116.47	6.92, 1H, <i>d</i> , <i>J</i> = 8.1 Hz	116.49	6.90, 1H, <i>d</i> , <i>J</i> = 8.1 Hz	116.50	6.92, 1H, <i>d</i> , <i>J</i> = 8.9 Hz
6'	129.01	7.92, 1H, <i>d</i> , <i>J</i> = 8.1 Hz	119.59	7.44, 1H, <i>d</i> , <i>J</i> = 8.1 Hz	128.99	7.92, 1H, <i>d</i> , <i>J</i> = 8.9 Hz
OCH ₃	60.53	3.72, 3H, <i>s</i>	60.58	3.72, 3H, <i>s</i>		12.98, <i>s</i>
OCH ₃	56.91	3.90, 3H, <i>s</i>	56.93	3.91, 3H, <i>s</i>		
C4'-OH		10.40, <i>br s</i>		12.91, <i>s</i>		
C5'-OH		12.90, <i>s</i>				

Compound (3) **Fig. 1** isolated as a pale yellow amorphous powder. The UV spectrum of (3) in methanol showed three peaks at 274, 348 and 447 nm. The UV spectrum of (3) with NaOMe showed peaks at 266 and 408 nm, and with NaOAc at 254 and 338 nm, and with AlCl₃ at 271 nm. These data revealed that (3) has a flavonoid type structure. The IR spectrum of (3) showed absorption band for hydroxyl groups 3410 cm⁻¹, carbonyl group 1647 cm⁻¹ (C=O α,β -unsaturated), and an aromatic double bond at 1580 cm⁻¹. Compound (3) gave a molecular ion peak in its EI-mass spectrum at *m/z* 330 (100%), in agreement with the molecular formula C₁₇H₁₄O₇.

The ^1H -NMR spectrum of (3) **Table 2** showed a characteristic low-frequency signal at δ 12.91 (*s*, OH) assignable to aromatic hydroxyl. Two *ortho*-coupled aromatic protons at δ 7.44 (1H, *d*, *J* = 8.1 Hz) and δ 6.90 (1H, *d*, *J* = 8.1 Hz) were diagnostic for a C-5' and C-6' oxygenated ring B. The ^1H -NMR spectrum showed three aromatic proton

signals at (δ 6.85, *s*, H-8), (δ 6.71, *s*, H-3) and (δ 7.45, *s*, H-2'). The ^1H -NMR also showed two singlet signal at δ 3.72 (*s*, 3H) and δ 3.91 (*s*, 3H) indicate the presence of two methoxy group. In ^{13}C -NMR spectrum **Table 2** and DEPT-135, the signals at δ 164.77, 103.18, 183.64, 152.58 and 105.54 were typical of C-2, C-3, C-4, C-9 and C-10 of a flavonoid moiety. The position of two methoxy group was authenticated through the HMBC spectrum between methoxy protons at δ 3.72 and C-6, and δ 3.91 and C-7. By comparison of ^1H - and ^{13}C -NMR chemical shifts with those reported in the literature, the flavonoid skeleton were in agreement with those of Cirsiol⁵⁵.

Compound (4) **Fig. 1** isolated as a yellow granular powder. The UV spectrum of (4) in methanol showed three peaks at 266 and 334 nm. The UV spectrum of (4) with NaOMe showed peaks at 274, 322 and 390 nm, and with NaOAc at 276, 300 and 378 nm, and with AlCl₃ at 277, 302, 338 and 384 nm. These data revealed that (4) has a flavonoid

type structure. The IR spectrum of (4) showed absorption band for hydroxyl groups 3352 cm^{-1} , α , β -unsaturated carbonyl group 1647 cm^{-1} , and an aromatic double bond at 1602 cm^{-1} . Compound (4) gave a molecular ion peak in its EI-mass spectrum at m/z 270 (100%), in agreement with the molecular formula $\text{C}_{15}\text{H}_{10}\text{O}_5$.

The $^1\text{H-NMR}$ spectrum of (4) **Table 2** showed two *ortho*-coupled aromatic protons at δ 7.92 (1H, *d*, $J = 8.9$ Hz) and δ 6.92 (1H, *d*, $J = 8.9$ Hz) were diagnostic for a C-2'/C-6' and C-3'/C-5' oxygenated ring B. The $^1\text{H-NMR}$ spectrum of (4) **Table 2** also showed two *meta*-coupled aromatic protons at δ 6.48 (1H, *d*, $J = 2.09$ Hz) and δ 6.18 (1H, *d*, $J = 2.09$ Hz) were diagnostic for a C-8 and C-6 oxygenated ring A. The $^1\text{H-NMR}$ also showed one singlet signal at δ 6.76 (*s*, 1H) for C-3. In $^{13}\text{C-NMR}$ spectrum **Table 2** and DEPT-135, the signals at δ 164.77, 103.32, 182.25, 157.86 and 104.14 were typical of C-2, C-3, C-4, C-9 and C-10 of a flavonoid moiety. Analysis of the $^1\text{H-NMR}$, $^{13}\text{C-NMR}$, COSY, HETCOR, and HMBC experiments indicated that this compound bears a flavonoid skeleton, which was identified as Apigenin, by comparison of $^1\text{H-}$ and $^{13}\text{C-NMR}$ chemical shifts with those reported in the literature ⁵⁶.

CONCLUSION: Phytochemical study of the aerial parts of *Teucrium polium* resulted in the isolation of four compounds, a new *neo*-clerodane diterpenoid (1), and three known flavonoids. The known flavonoids were identified as Cirsimartin (2), Cirsiliol (3), and Apigenin (4). The structures were determined by physical, chemical and spectral techniques.

ACKNOWLEDGEMENT: The author is very grateful to Prof. Dr. Anwer Al-khateeb (Botany Department, Faculty of Science, Damascus University) for plant taxonomic identification. The author also grateful to the mass spectra unit at the Department of Chemistry, Atomic Energy Commission of Syria.

CONFLICT OF INTEREST: We declare that we have no conflict of interest.

REFERENCES:

- Mouterde P: Nouvelle Flore du Liban et de la Syrie. Dar el-Machreq, Beirut 1983; 3.
- Menichini F, Conforti F, Rigano D, Formisano C, Piozzi F and Senatore F: Phytochemical composition, anti-inflammatory and antitumor activities of four *Teucrium* essential oils from Greece. Food Chemistry 2009; 115: 679-686.
- Abdollahi M, Karimpour H and Monsef-Esfehani HR: Antinociceptive effects of *Teucrium polium* total extract and essential oil in mouse writhing test. Pharmacol Res 2003; 48: 31-35.
- Baluchnejadmojarad T, Roghani M and Roghani-Dehkordi F: Antinociceptive effects of *Teucrium polium* leaf extract in the diabetic rat formalin test. J Ethnopharmacol 2005; 97: 207-210.
- Rasekh HR, Khoshnood-Mansourkhani MJ and Kamalinejad M: Hypolipidemic effects of *Teucrium polium* in rats. Fitoterapia 2001; 72: 937-939.
- Afifi FU, Al-Khalidi B and Khalil E: Studies on the *in-vivo* hypoglycemic activities of two medicinal plants used in the treatment of diabetes in Jordanian traditional medicine following intranasal administration. J Ethnopharmacol 2005; 14: 314-318.
- Esmaceli MA and Yazdanparast R: Hypoglycaemic effect of *Teucrium polium*: studies with rat pancreatic islets. J Ethnopharmacol 2004; 95: 27-30.
- Vessal M, Zal F and Vasei M: Effects of *Teucrium polium* on oral glucose tolerance test, regeneration of pancreatic islets and activity of hepatic glucokinase in diabetic rats. Arch Iranian Med 2003; 6: 35-39.
- Autore G, Capasso F, De Fusco R, Fasulo MP, Lembo M, Mascolo N and Menghini A: Antipyretic and antibacterial actions of *Teucrium polium* L. Pharmacol Res Commun 1984; 16: 21-29.
- Aggelis G, Athanassopoulos N, Paliogianni A and Komaitis M: Effect of a *Teucrium polium* L. extract on the growth and fatty acid composition of *Saccharomyces cerevisiae* and *Yarrowia lipolytica*. Antonie Van Leeuwenhoek 1998; 73: 195-198.
- Khader M, Eckl PM and Bresgen N: Effects of aqueous extracts of medicinal plants on MNNG-treated rat hepatocytes in primary cultures, J Ethnopharmacol 2007; 112: 199-202.
- Khader M, Bresgen N and Eckl PM: Antimutagenic effects of ethanolic extracts from selected Palestinian medicinal plants. J Ethnopharmacol 2010; 127: 319-324.
- Nematollahi-Mahani SN, Rezazadeh-Kermani M, Mehrabani M and Nakhaee N: Cytotoxic effects of *Teucrium polium* on some established cell lines. Pharm Biol 2007; 45: 295-298.
- Juricic RV, Kalodera Z and Grgic J: Determination of selenium in *Teucrium* generation atomic absorption spectrometry. Z Naturforsch 2003; 58: 143-145.
- De Marino S, Festa C, Zollo F, Incollingo F, Raimo G, Evangelista G and Iorizzi M: Antioxidant activity of phenolic and phenylethanoid glycosides from *Teucrium polium*. Food Chemistry 2012; 133: 21-28.
- Tepe B, Degerli S, Arslan S, Malatyali E and Sarikurku C: Determination of chemical profile, antioxidant, DNA damage protection and antimicrobial activities of *Teucrium polium* and *Stachys iberica*. Fitoterapia 2011; 82: 237-246.
- Hasani P, Yasa N, Vosough-Ghanbari S, Mohammadirad A, Dehghan G and Abdollahi M: *In-vivo* antioxidant potential of *Teucrium polium* as compared to α -tocopherol. Acta Pharm 2007; 57: 123-129.
- Sharififar F, Dehghn-Nudeh G and Mirtajaldini M: Major flavonoids with antioxidant activity from *Teucrium polium* L. Food Chemistry 2009; 112: 885-888.

19. Esmaeili MA, Zohari F and Sadeghi H: Antioxidant and protective effects of major flavonoids from *Teucrium polium* on beta-cell destruction in a model of streptozotocin-induced diabetes. *Planta Med* 2009; 75: 1418-1420.
20. Ardestani A, Yazdanparast R and Jamshidi S: Therapeutic effects of *Teucrium polium* extract on oxidative stress in the pancreas of streptozotocin-induced diabetic rats. *J Med Food* 2008; 11: 525-532.
21. Suboh SM, Bilot YY and Aburjai TA: Protective effects of selected medicinal plants against protein degradation, lipid peroxidation and deformability loss of oxidatively stressed human erythrocytes. *Phytother Res* 2004; 18: 280-284.
22. Boumerfeg S, Baghiani A, Djarmouni M, Ameni D, Adjadji M, Belkhirri F, Charef N, Khennouf S and Arrar L: Inhibitory Activity on Xanthine Oxidase and Antioxidant Properties of *Teucrium polium* L. Extracts. *Chinese Medicine* 2012; 3: 30-41.
23. Starakis I, Siagris D, Leonidou L, Mazokopakis E, Tsamandasc A and Karatza C: Hepatitis caused by the herbal remedy *Teucrium polium* L. *Eur J Gastroenterol Hepatol* 2006; 18: 681-683.
24. Mattei A, Rucay P, Samuel D, Feray C, Reynes M and Bismuth H: Liver transplantation for severe acute liver failure after herbal medicine (*Teucrium polium*) administration. *J Hepatol* 1995; 22: 597.
25. Larrey D, Vial T, Pauwels A, Castot A and Michel H: Hepatitis after germander (*Teucrium chamaedrys*) administration, another instance of herbal medicine hepatotoxicity. *Ann Intern Med* 1992; 117: 129-132.
26. Mattei A, Bizollon T and Charles JD: Atteinte hepatique associee a la prise d'un produit de phytotherapie contenant de la germandee petit-chene. *Gastroenterol Clin Biol* 1992; 16: 798-800.
27. Wassel GM and Ahmed SS: On the essential oil of *Teucrium polium* L. *Pharmazie* 1974; 29: 351-352.
28. Hassan MM, Muhtadi FJ and Al-Badr AA: GLC-mass spectrometry of *Teucrium polium* oil. *J Pharm Sci* 1979; 68: 800-801.
29. Cakir A, Duru ME, Harmandar M, Ciriminna R, Passannanti S and Kazim K: Volatile constituents of *Teucrium polium* L. from Turkey. *J Essent Oil Res* 1998; 10: 113-115.
30. Eikani MH, Goodarznia L and Mirza M: Comparison between the essential oil of supercritical carbon dioxide extract of *Teucrium polium* L. *J Essent Oil Res* 1999; 11: 470-472.
31. Perez-Alonso MJ, Velasco-Negueruela A and Lopez-Saez JA: The essential oils of two Iberian *Teucrium* species. *J Essent Oil Res* 1993; 5: 397-402.
32. Aburjai T, Hudaib M and Cavrini V: Composition of the essential oil from Jordanian germander *Teucrium polium* L. *J Essent Oil Res* 2006; 18: 97-99.
33. Sadeghia H, Jamalpoorb S and Shirzadib MH: Variability in essential oil of *Teucrium polium* L. of different latitudinal populations. *Industrial Crops and Products* 2014; 54: 130-134.
34. Djabou N, Lorenzi V, Guinoiseau E, Andreani S, Giuliani MC, Desjobert JM, Bolla JM, Costa J, Berti L, Luciani A and Muselli A: Phytochemical composition of Corsican *Teucrium* essential oils and antibacterial activity against foodborne or toxi-infectious pathogens. *Food Control* 2013; 30: 354-363.
35. Purnavaba S, Ketabchia S and Rowshan V: Chemical composition and antibacterial activity of methanolic extract and essential oil of Iranian *Teucrium polium* against some of phyto-bacteria. *Natural Product Research* 2015; 29: 1376-1379.
36. Eguren L, Perales A, Fayos J, Savona G, Paternostro M, Piozzi F and Rodriguez B: New Clerodane Diterpenoid from *Teucrium polium* subsp. aureum X-ray Structure Determination. *J Org Chem* 1981; 46: 3364-3367.
37. Malakov P, Papanov G and Ziesche J: Teupolin III A Furanoid Diterpene from *Teucrium polium*. *Phytochemistry* 1982; 21: 2597-2598.
38. Malakov P and Papanov G: Furanoid Diterpene from *Teucrium polium*. *Phytochemistry* 1983; 22: 2791-2793.
39. Fernandez P, Rodriguez B, Savona G and Piozzi F: Neo-Clerodane diterpenoids from *Teucrium polium* subsp capitatum. *Phytochemistry* 1986; 25: 181-184.
40. De la Torre M, Piozzi F, Rizk AF, Rodriguez B and Savona G: 19-Acetylteupolin IV, A neo-Clerodane diterpenoid from *Teucrium polium* subsp pilosum. *Phytochemistry* 1986; 25: 2239-2240.
41. Malakov P, Boneva IM, Papanov G and Spassov SL: Teulamifin B, A neo-Clerodane diterpenoid from *Teucrium lamiifolium* and *T. polium*. *Phytochemistry* 1988; 27: 1141-1443.
42. Alcazar R, De La Torre MC, Rodriguez B, Bruno M, Piozzs F and Arnold GNA: Neo-Clerodane Diterpenoids from three Species of *Teucrium*. *Phytochemistry* 1992; 31: 3957-3960.
43. Bruno M, Maggiosa AM, Piozzi F, Puech S, Rosselli SG and Simmonds MSJ: Neoclerodane diterpenoids from *Teucrium polium* subsp. polium and their antifeedant activity. *Biochemical Systematics and Ecology* 2003; 31: 1051-1056.
44. Malakov P, Papanov GG and Mollov NM: Montanin A and B, new furanoid diterpenes of nor-clerodane type from *Teucrium montanum* L. *Tetrahedron Letters* 1978; 19: 2025-2026.
45. D'Abrosca B, Pacifico S, Scognamiglio M, D'Angelo G, Galasso S, Monaco P and Fiorentino A: A new acylated flavone glycoside with antioxidant and radical scavenging activities from *Teucrium polium* leaves. *Natural Product Research* 2013; 27: 356-363.
46. Rizk AM, Hammouda FM, Rimpler H and Kamel A: Iridoids and flavonoids of *Teucrium polium* Herb. *Planta Medica* 1986; 52: 87-88.
47. Verykokidou-Vitsaropoulou E and Vajias C: Methylated flavones from *T. polium*, *Planta Med* 1986; 52: 401-402.
48. Kawashty SA, Gamal El-din EM and Saleh NAM: The flavonoid chemo-systematics of two *Teucrium* species from Southern Sinai Egypt. *Biochemical Systematics and Ecology* 1999; 27: 657-660.
49. Oganessian GB, Galstyan AM, Mnatsakanyan VA, Shashkov AS and Agababyan PV: Phenylpropanoid Glycosides of *Teucrium polium*. *Khimiya Prirodnykh Soedinenii* 1977; 1: 630-634.
50. Moustapha C, Hasen T and Sadaka MWM: Chemical Constituents of *Teucrium polium* L. var. mollissimum Hand-Mazz, *Jordan J Chem* 2011; 6: 339-345.
51. Malakov P, Papanov G and Ziesche J: Teupolin III, A furanoid diterpene from *Teucrium polium*, *Phytochemistry* 1982; 21: 2597-2598.
52. De la Torre MC, Piozzi F, Rizk AF, Rodriguez B and Savona G: 19-Acetylteupolin IV, A neo-clerodane diterpenoid from *Teucrium polium* subsp. Pilosum. *Phytochemistry* 1986; 25: 2239-2240.
53. Yim SH, Kim HJ and Lee IS: A Polyacetylene and Flavonoids from *Cirsium rhinoceros*. *Arch Pharm Res* 2003; 26: 128-131.

54. Marder M, Viola H, Wasowaki C, Wolfman C, Waterman PG, Medina JH and Paladini AC: Cirsiliol and caffeic acid ethyl ester, isolated from *Salvia guaranitica*, are competitive ligands for the central benzodiazepine receptors. *Phytomedicine* 1996; 3: 29-31.
55. Kim NM, Kim J, Chung HY and Choi JS: Isolation of luteolin 7-o-rutinoside and esculetin with potential antioxidant activity from the aerial parts of *Artemisia montana*. *Arch Pharm Res* 2000; 23: 237-239.

How to cite this article:

Sadaka MWM: A new neo-clerodane diterpenoid from *Teucrium polium* L.. *Int J Pharmacognosy* 2017; 4(6): 185-92. doi link: [http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.4\(6\).185-92](http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.4(6).185-92).

This Journal licensed under a Creative Commons Attribution-Non-commercial-Share Alike 3.0 Unported License.

This article can be downloaded to **Android OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)