



Received on 03 January 2018; received in revised form, 03 February 2018; accepted, 13 February 2018; published 01 May 2018

IDENTIFICATION OF BIOACTIVE COMPOUNDS OF *ALCHEMILLA CAUCASICA* USING GAS CHROMATOGRAPHY-MASS SPECTROMETRY

Esen Sezen Karaoglan ^{*1} and Bilal Yilmaz ²

Department of Pharmaceutical Botany ¹, Department of Analytical Chemistry ², Faculty of Pharmacy, Ataturk University, 25240, Erzurum, Turkey.

Keywords:

Alchemilla caucasica,
Rosaceae, Bioactive compounds, Gas
chromatography-mass spectrometry

Correspondence to Author:

Dr. Esen Sezen Karaoglan

Assistant Professor,
Department of Pharmaceutical
Botany, Faculty of Pharmacy, Ataturk
University, 25240, Erzurum, Turkey.

E-mail: esen.karaoglan@atauni.edu.tr

ABSTRACT: *Alchemilla* species (Rosaceae) are locally known as "aslanpençesi" in Turkey. *Alchemilla* plants (lady's mantle) are used in traditional medicine for different indications. In this study, the methanol extract of *Alchemilla caucasica* was analyzed using gas chromatography-Mass Spectrometry, while the mass spectra of the compounds found in the extract were matched with the National Institute of Standards and Technology Library. The study revealed the presence of fourteen phyto-components. The most abundant compounds in the methanol extract were 13-docosenamide (Z)-, 5- octadecenoic acid 2,3-dihydroxy-propyl ester and hexadecanoic acid 2-hydroxy-1-(hydroxyl-methyl) ethyl ester which had concentrations of 59.77%, 15.34%, and 7.59% respectively. Results indicated that polarity of the solvents used in extraction influenced the relative abundance and type of compounds extracted. Various compounds identified are known to have varying bioactivities such as being anticancer and antioxidant. Therefore, the presence of these compounds in *Alchemilla caucasica* could be useful in the preparation of functional foods and nutraceuticals in preventing and treating debilitating and chronic human diseases.

INTRODUCTION: Plants play a major role in primary health care as therapeutic remedies in developing countries ¹. They serve as a source of medicine and an important component in health care system ². This stems from the fact that aside from providing vitamins and minerals, plants contain phytochemicals which are secondary metabolites. These phytochemicals have various bioactivities such as being antioxidants, anti-nutritional or cytotoxic.

Antioxidants are substances used by the body to protect itself from damage caused by free radicals, whose accumulation has been linked to several diseases such as heart diseases ³, liver diseases and cancers ⁴.

The anti-nutritional activity of a bioactive compound is based on its ability to inhibit the absorption and utilization of nutrients. Currently, there is upsurge interest in the knowledge of bioactive compounds present in plants and hence their beneficial potentials as nutraceuticals and functional foods; hence, having positive health-promoting effects. Herbal medicine, as a major part of traditional medicine, has been used in medical practice since antiquity and is a common element of Ayurvedic, homeopathic, and naturopathic medicine.

	QUICK RESPONSE CODE DOI: 10.13040/IJPSR.0975-8232.IJP.5(5).287-93
	The article can be accessed online on www.ijpjournal.com
DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5(5).287-93	

World health organization notes that 74% of the plant-derived medicines are used in modern medicine, in a way that their modern application directly correlates with their traditional use as herbal medicines by native cultures^{5,6}.

Alchemilla species belonging to Rosaceae family are locally known as "aslanpençesi" in Turkey. These species include fatty acids, esters, aldehydes, terpenes, hydrocarbons, mainly phenolic compounds, including flavonoids and tannins⁷⁻¹⁰. *Alchemilla* plants (lady's mantle) are used in traditional medicine for different indications which are minimizing the symptoms of sore throat, promoting wound healing, stopping bleeding, gynecological diseases, alleviating nausea and vomiting¹¹. Different studies showed that the phenolic compounds (tannins, flavonoids, etc.) present in the plant are responsible for the pharmacological activity of Lady's mantle¹²⁻¹⁴. *Alchemilla caucasica* is a member of this genus. This plant has short, erect flowering stems, wide and reniform leaves¹⁵ **Fig. 1**.



FIG. 1: ALCHEMILLA CAUCASICA PLANT

Knowledge of the chemical constituents of *Alchemilla caucasica* is desirable not only for the discovery of therapeutic agents but also because such information may be of great value in disclosing new sources of economic phyto-compounds for the synthesis of complex chemical substances and for discovering the actual significance of folkloric remedies. Hence, a thorough validation of the herbal drugs has emerged as a new branch of science emphasizing and prioritizing the standardization of the natural drugs and products because several of the phytochemical have the complementary and over-

lapping mechanism of action. Mass spectrometry coupled with chromatographic separations such as gas chromatography (GC-MS) is normally used for direct analysis of components existing in traditional medicines and medicinal plants. In previous studies, it was reported that the seeds contained mainly palmitic, stearic, oleic and linoleic fatty acids¹⁶. In this study, the fourteen chemical content of the aerial part of the plant was tried to be determined by GS-MS method.

MATERIALS AND METHODS:

Chemicals and Reagents: Glycolaldehyde dimethyl acetal, furfural, phthalic acid dibutyl ester, n-hexadecanoic acid, linoleic acid, linolenic acid, octadecanoic acid, hexadecanoic acid 2-hydroxy- 1- (hydroxymethyl) ethyl ester, dicyclohexyl phthalate, octadecanoic acid, 2,3-dihydroxypropyl ester, 13- docosenamide (Z)-, octadecanamide, 24, 25 dihydroxycholecalciferol and β -sitosterol were purchased from Sigma-Aldrich (St. Louis, MO, USA).

GC-MS System: Chromatographic analysis were carried out on an Agilent 7820A gas chromatography system equipped with 5977 series mass selective detector, 7673 series autosampler and chemstation (Agilent Technologies, Palo Alto, CA). HP-5 MS column with 0.25 μ m film thickness (30 m \times 0.25 mm I.D., USA) was used for separation. The temperatures of the inlet, transfer line, and detector were 250, 250 and 300 $^{\circ}$ C, respectively.

GC-MS Conditions: Different temperature programs were investigated for GC-MS method. The end of this investigation, the temperature program of the GC/MS was as follows: initial temperature was 50 $^{\circ}$ C, held for 1 min, increased to 100 $^{\circ}$ C at a rate of 20 $^{\circ}$ C/min held for 1 min, increased to 180 $^{\circ}$ C at a rate of 10 $^{\circ}$ C/min held for 1 min, increased to 220 $^{\circ}$ C at a rate of 5 $^{\circ}$ C/min held for 5 min, and finally to 300 $^{\circ}$ C at a rate of 10 $^{\circ}$ C/min and held for 5.5 min.

Extraction Procedure: *Alchemilla caucasica* were collected from Konaklı Mountain (Erzurum province, Turkey) in June 2016. A voucher specimen was deposited in the Herbarium of Ataturk University, Faculty of Pharmacy (AUEF1023). The air-dried and powdered aerial

parts of the plant were extracted three times with methanol in the mantle heater at 40 °C (3×1 L). After filtration, 1 μ l of this solution was employed for GC-MS analysis.

Identification of Components: Identification was based on the molecular structure, molecular mass, and calculated fragments. Interpretation on mass spectrum GC-MS was conducted using the database on National Institute Standard and Technology having more than 62,000 patterns. The name, molecular weight, and structure of the components of the test materials were ascertained. The relative percentage amount of each component was calculated by comparing its average peak area to the total areas. The spectrum of the unknown component was compared with the spectrum of the component stored in the National Institute of Standards and Technology Library Version (2005), Software, Turbomass 5.2.

RESULTS AND DISCUSSIONS:

Method Development and Optimization: The method development for the assay of phyto-components was based on their chemical properties. In this study, the capillary column coated with 5% phenyl, 95% dimethylpolysiloxane is a good choice for separation of these analytes since they elute as symmetrical peaks at a wide range of concentrations. Different temperature programs were investigated for the GC oven. The end of this investigation, the best temperature

program was selected for good separation. The temperature programs of the GC oven were as follows: initial temperature 50 °C, held for 1 min, increased to 100 °C at a rate of 20 °C/min held for 1 min, increased to 180 °C at a rate of 10 °C/min held for 1 min, increased to 220 °C at a rate of 5 °C/min held for 5 min, and finally to 300 °C at a rate of 10 °C/min and held for 5.5 min. The splitless injection mode was chosen. The injector volume was 1 μ l in splitless mode, and the carrier gas was helium at a flow rate of 1 ml/min.

GC-MS Analysis: The more precise information in qualitative analysis can be obtained by GC-MS. For quantitative determination, gas chromatography with flame ionization detector (GC-FID) and GC-MS are preferred. GC-MS is one of the best techniques to identify the constituents of volatile matter, long chain, branched chain hydrocarbons, alcohols, acids, esters *etc.*

The GC-MS analysis of *Alchemilla caucasica* extracts revealed the presence of fourteen compounds (phytochemical constituents) that could contribute to the medicinal quality of the plant. The identification of the phytochemical compounds was confirmed based on the peak area, retention time and molecular formula. The major phytochemical constituent's present in methanolic extract of *Alchemilla caucasica* is presented as compound chromatogram is in **Fig. 2**.


















FIG. 2: GC-MS CHROMATOGRAM OF METHANOLIC EXTRACT OF *ALCHEMILLA CAUCASICA*

GC-MS analysis revealed the presence of fourteen compounds in methanol extracts of *Alchemilla caucasica*. Peak number, retention time, compound name, molecular weight, and molecular formula are stated in **Table 1**. It shows compounds identified in methanol extract of *Alchemilla caucasica* which represented the hydrophobic and hydrophilic bioactive fraction. Three compounds were identified as the major bioactive compounds. They are 13-docosenamide (Z)- and it had a relative abundance of 59.77% and eluted as represented at peak 11, octadecenoic acid 2, 3-dihydroxy-propyl ester ester had a relative abundance of 15.34% and eluted as represented at peak 10 and hexadecanoic acid 2-hydroxy-1-(hydroxymethyl)ethyl ester had a

relative abundance of 7.59% and eluted as represented at peak 8. Major fatty acids have biological activities such as being anti-inflammatory, anti-androgenic, cancer preventing, dermatogenic, hypocholesterolemic, anaemiagenic, insectifuge as well as being a 5-alpha-reductase inhibitor¹⁷. n-hexadecanoic acid has been reported to show selective toxicity to human leukemic cells as well as antitumor activity and is suggested to be a lead compound in anti-cancer drugs¹⁸. Other fatty acids and fatty acid esters identified were linoleic acid (0.69%), linolenic acid (1.96%), octadecanoic acid (0.56%) and n-hexadecanoic acid (1.81%).

TABLE 1: CHEMICAL COMPOSITION OF THE BIOACTIVE COMPOUNDS OF *ALCHEMILLA CAUCASICA*

Peak	Retention time (minute)	Compound	Molecular formula	Chemical formulation	% of the total
1	3.252	Glycolaldehyde dimethyl acetal	C ₄ H ₁₀ O ₃		0.319
2	3.459	Furfural	C ₅ H ₄ O ₂		0.319
3	17.187	Phthalic acid dibutyl ester	C ₁₆ H ₂₂ O ₄		0.980
4	18.675	n-Hexadecanoic acid	C ₁₆ H ₃₂ O ₂		1.811
5	21.581	Linoleic acid	C ₁₈ H ₃₂ O ₂		0.688
6	21.712	Linolenic acid	C ₁₈ H ₃₀ O ₂		1.963
7	22.070	Octadecanoic acid	C ₁₈ H ₃₆ O ₂		0.555

8	29.728	Hexadecanoic acid 2-hydroxy-1-(hydroxymethyl) ethyl ester	$C_{19}H_{38}O_4$		7.594
9	29.838	Dicyclohexyl phthalate	$C_{20}H_{26}O_4$		2.031
10	32.396	Octadecanoic acid 2,3-dihydroxypropyl ester	$C_{21}H_{42}O_4$		15.335
11	33.097	13-Docosamide, (Z)-	$C_{22}H_{43}NO$		59.771
12	13.246	Octadecanamide	$C_{18}H_{37}NO$		1.728
13	34.867	24, 25 dihydroxycholecalciferol	$C_{27}H_{44}O_3$		0.711
14	38.534	β - sitosterol	$C_{29}H_{50}O$		1.561

Phthalic acid and phthalate derivatives are major industrial materials used to manufacture plastic products like toys and bottles, also being widely used as plasticizers, adhesives, films, polymers, etc. However, some studies revealed that these compounds could affect the male reproductive systems producing testicular atrophy¹⁹ and alter the normal development of fetuses in pregnant rats²⁰. In view of these adverse effects on living beings, the US Environmental Protection Agency (EPA) classified the phthalic acid and some industrial

phthalates as priority pollutants²¹ and so, for example, the maximum admissible content in water for a common phthalic acid derivative such as the phthalic acid dibutyl ester was established in 6 g/dm.

Sterols are important constituents of all eukaryotes and play a vital role in plant cell membranes. Plant sterols possess valuable physiological activities; they are biogenetic precursors of many hormones and oviposition stimulants of some insects²².

Among all phytochemicals, β -sitosterol is the main phytosterol found in many plants. It has been reported to show anti-inflammatory, antineoplastic, antipyretic, and immunomodulating activity²³.

The chemical composition of the total methyl esters of fatty acids from the extracts of *Alchemilla caucasica* showed to have a very similar profile. Fatty acid methyl esters are an alternative diesel fuel (namely, biodiesel) derived from vegetable oils or animal fats²⁴. The main components of vegetable oils and animal fats are triglycerides or also known as esters of fatty acids attached to a glycerol. The most widely used industrial method for the commercial production of fatty acid methyl esters from vegetable oils/fats is a base catalyzed trans-esterification process using sodium hydroxide or potassium hydroxide as the homogeneous catalyst and methanol as the lower alcohol²⁵.

CONCLUSION: The source of many plants can often be identified from the peak pattern of the chromatograms obtained directly from GC-MS analysis. Similarly, unique qualitative and quantitative patterns from a GC analysis will often help identify the source of many alcoholic beverages. The technique of fingerprint could identify false herbal products. The construction of chromatographic fingerprints aims at evaluating the quality of herbal medicines. The fundamental reason of quality control of herbal medicines is based on the concept of phytoequivalence of herbs, and then to use this conception to identify the real herbal medicine and the false one, and further to do quality control. The importance of the study is due to the biological activity of some of these compounds. The present study, which reveals the presence of components in *Alchemilla caucasica* suggest that the contribution of these compounds on the pharmacological activity should be evaluated.

ACKNOWLEDGEMENT: Nil

CONFLICT OF INTEREST: Nil

REFERENCES:

- Jonathan G, Ohimain E and Kigigha L: Antagonistic effects of extracts of some Nigerian higher fungi against selected microorganisms. American-European Journal of Agriculture Environment and Science 2007; 2: 364-68.

- Kathirvel A and Sujatha V: Phytochemical studies, antioxidant activities and identification of active compounds using GC-MS of *Dryopteris cochleata* leaves. Arabian Journal of Chemistry 2016; 9: 1435-42.
- Aviram M: Review of human studies on oxidative damage and antioxidative damage and antioxidant protection related to cardiovascular disease. Free Radical Research 2000; 33: 585-97.
- Owen RW, Giacosa WE, Hull R, Haubner BS and Bartsh H: The antioxidant/anticancer potential of phenolic compounds isolated from olive oil. European Journal of Cancer 2000; 36: 1235-47.
- Kumar V and Parmar NS: Herbs: A potential source for the development of new phytomedicines. The Pharma Review 2003; 1: 59-63.
- Mukherjee PK: Quality control of herbal drugs. An approach to the evaluation of botanicals. Business Horizons Pharmaceutical Publishers 2002.
- Babayana MS, Chelombit'ko VA: Study of polysaccharides of herb *Alchemilla tredecimloba* from Rosaceae family. Izvestiya Vysshikh Uchebnykh Zavedenii, Severo-Kavkazskii Region, Estestvennye Nauki 2009; 4: 68-69.
- Falchero L, Coppa M, Fossi A, Lombardi G, Ramella D and Tava A: Essential oil composition of lady's mantle (*Alchemilla xanthochlora* Rothm.) growing wild in Alpine pastures. Natural Product Research 2009; 23: 1367-72.
- Kaya B, Menemen Y and Saltan FZ: Flavonoids in the endemic species of *Alchemilla* L., (section *Alchemilla* L. subsection *Calycanthum* Rothm. ser. *elatae* Rothm.) from North-East Black sea region in Turkey. Pakistan Journal of Botany 2012; 44: 595-97.
- Kaya B, Menemen Y and Saltan FZ: Flavonoid compounds identified in *Alchemilla* L. species collected in the north-eastern Black Sea region of Turkey. African Journal of Traditional, Complementary and Alternative Medicines 2012; 9: 418-25.
- Makau JN, Watanabe K and Kobayashi N: Anti-influenza activity of *Alchemilla mollis* extract: Possible virucidal activity against influenza virus particles. Drug Discoveries and Therapeutics 2013; 7: 189-95.
- Jonadet M, Meunier MT, Villie F, Bastide JP and Lamaison JL: Flavonoids extracted from *Ribes nigrum* L. and *Alchemilla vulgaris* L.: 1. *In-vitro* inhibitory activities on elastase, trypsin and chymotrypsin. 2. Angioprotective activities compared *in-vivo*. Journal of Pharmacology 1986; 17: 21-27.
- Filipek J: Effect of *Alchemilla xanthochlora* Water Extract on Lipid Peroxidation and Superoxide Anion Scavenging Activity. Pharmazie 1992; 47: 717-18.
- Schimmer O and Lindenbaum M: Tannins with anti-mutagenic properties in the herb of *Alchemilla* species and *Potentilla anserina*. Planta medica 1995; 61: 141-45.
- PH Davis: Flora of Turkey and the East Aegean Islands. Edinburgh University Press 1972; 4: 88.
- Matthaus B and Özcan MM: Fatty acid, tocopherol and squalene contents of Rosaceae seed oils. Botanical Studies 2014; 55(48): 1-6.
- Duke JA: Phytochemical and ethnobotanical databases; Available: <https://phytochem.usda.gov/phytochem/search/list>, 1998.
- Harada H, Yamashita U, Kurihara H, Fukushi E, Kawabata J and Kamei Y: Antitumor activity of palmitic and found as the selective cytotoxic substance in marine red algae. Anticancer Research 2002; 22: 2587-90.
- Oishi S and Hiraga K: Testicular atrophy induced by phthalic acid esters: Effect on testosterone and zinc

- concentrations, Toxicology and Applied Pharmacology 1980; 53: 35-41.
20. Makoto E, Emiko M, Akira H and Kunio K: Developmental toxicity evaluation of phthalic acid, one of the metabolites of phthalic acid esters, in rats. Toxicology Letters 1997; 93: 109-15.
 21. The United States Environmental Protection Agency (7407), Pollution Prevention and Toxics, EPA 749-F-95-016a, 1994.
 22. Harborne JB: Twenty-five years of chemical ecology. Natural Product Reports 2001; 18: 361-79.
 23. Fraile L, Crisci E, Córdoba L, Navarro MA, Osada J and Montoya M: Immunomodulatory properties of beta-sitosterol in pig immune responses. International Immunopharmacology 2012; 13: 316-21.
 24. Vasudevan PT and Briggs M: Biodiesel production current state of the art and challenges. The Journal of Industrial Microbiology and Biotechnology 2008; 35: 421-30.
 25. Semwal S, Arora AK, Badoni RP and Tuli DK: Biodiesel production using heterogeneous catalysts. Bioresource Technology 2011; 102: 2151-61.

How to cite this article:

Karaoglan ES and Yilmaz B: Identification of bioactive compounds of *Alchemilla caucasica* using gas chromatography-mass spectrometry. Int J Pharmacognosy 2018; 5(5): 287-93. doi link: [http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5\(5\).287-93](http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5(5).287-93).

This Journal licensed under a Creative Commons Attribution-Non-commercial-Share Alike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)