



Received on 12 January 2022; received in revised form, 23 January 2022; accepted, 28 January 2022; published 31 January 2022

## ANTI-PLASMODIAL ACTIVITY OF *BLIGHIA SAPIDA* LEAF EXTRACTS IN *PLASMODIUM BERGHEI* INFECTED MICE

Olumekun Victor Olugbenga\*, Odukomaiya, Samuel Odubajo and Sunday, Matthew Damilola

Department of Plant Science and Biotechnology, Adekunle Ajasin University, Akungba-Akoko, Nigeria.

### Keywords:

*Blighia sapida*, Ackee Apple, *Plasmodium berghei*, Anti-plasmodial activity, Medicinal plant

### Correspondence to Author:

**Olumekun, Victor Olugbenga**

Professor,  
Department of Plant Science and  
Biotechnology, Adekunle Ajasin  
University, Akungba-Akoko, Nigeria.

**E-mail:** victor.olumekun@aau.edu.ng

**ABSTRACT:** *Blighia sapida* a tropical evergreen plant with a dense crown, is commonly called ackee apple. Traditionally, the stem barks and leaves of the plant are used in the treatment of high fever, strokes, malaria, headache and some eye ailments. This study examined the antiplasmodial activity of *Blighia sapida* leaf extracts in comparison with the standard antimalarial drug Artemether-Lumefantrine combination, with a view to determining the changes in the parasitemia level in mice infected with *Plasmodium berghei*. The results showed a significant decrease in parasitemia level over six days period for mice treated with Artemether-Lumefantrine, ethanolic, and aqueous leaf extracts of *B. sapida*, whereas the controls showed a continuous increase. The parasitemia level was significantly reduced ( $P < 0.05$ ) in mice treated with 6.0mg/kg body weight of Artemether-Lumefantrine with 88% suppression; those treated with the ethanolic leaf extract at 50mg/kg and 30mg/kg body weight had parasitemia suppressed by 59% and 43.67% respectively. Parasitaemia count in mice treated with 50mg/kg and 30mg/kg body weight of aqueous leaf extract of *B. sapida* was suppressed by 43.83% and 40% respectively. This study reveals that ethanolic and aqueous leaf extracts of *Blighia sapida* have promising anti-malaria activity at the concentrations used, although these were not as effective as the Artemether-Lumefantrine combination. However, the results lend credence to the ethnobotanical use of the plant for the treatment of malaria.

**INTRODUCTION:** Malaria is the most devastating human disease on earth; nearly half of the world's population is at risk of malaria. According to the World Health Organization<sup>1</sup>, in 2015, there were roughly 212 million malaria cases and an estimated 429,000 malaria deaths. Increased prevention and control measures have led to a 29% reduction in malaria mortality rates globally since 2010.

Sub-Saharan Africa still carries a disproportionately high share of the global malaria burden. In 2015, the region was home to 90% of malaria cases and 92% of malaria deaths<sup>1</sup>. In 2010, about 3.3 billion people were exposed to malaria. The highest risk was for people living in Africa, where approximately 91% of the cases were recorded, while 86% of deaths globally occur among children under five years of age<sup>2,3</sup>.

In adults, its common symptoms are headaches, weakness, fever, aches and pains, high body temperature, and bitterness of the mouth with loss of appetite, while in children, in addition to the above-mentioned symptoms, it may also manifest with nausea and vomiting<sup>4</sup>. The high rate of deaths in Africa is traceable to several factors such as poor

	<p><b>QUICK RESPONSE CODE</b></p>
	<p><b>DOI:</b> 10.13040/IJPSR.0975-8232.IJP.9(1).17-24</p>
<p>The article can be accessed online on: <a href="http://www.ijournal.com">www.ijournal.com</a></p>	
<p>DOI link: <a href="http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.9(1).17-24">http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.9(1).17-24</a></p>	

economic standards, the high cost of synthetic medicines, the development of resistance by the causative protozoan parasite, and the side effects of existing orthodox drugs, which have made the choice of herbal remedies against these infections inevitable and more economical<sup>5</sup>. The severe long-term effects of malaria parasites have compelled continued pharmacological investigations into plants used for treating malaria disease in search of potent and cost-effective solutions to mitigate the effect of the disease.

*Plasmodium berghei* is a protozoan parasite that infects mammals other than humans. They are not of direct concern to man practically; man's interest in these parasites is that they are practical model organisms in the laboratory for the experimental study of human malaria<sup>6</sup>. Rodent malaria parasites are used in many Research Institutes for studies to develop new drugs or vaccines against malaria<sup>7</sup>. Like all malaria parasites of mammals, *P. berghei* is transmitted by the anopheles mosquito, and it infects the liver after being released into the bloodstream by a bite from an infected female mosquito<sup>8</sup>.

From ancient times, plants and their extracts have been used in the treatment of diseases in Africa, Asia and North America. Traditional medicine has been the focus for wider coverage of primary health care delivery in Africa and the rest of the world<sup>9</sup>. It comprises therapeutic practices for hundreds of years before the development of modern scientific medicine. It is still in use today without much-documented evidence of adverse effects<sup>10</sup>.

Several antimalarial drugs have been developed from plant-based materials, for example, the alkaloid (quinine) derived from the bark of *Cinchona spp.* (Rubiaceae) and lately, artemisinin derived from *Artemisia annua* (Asteraceae). Historically, the majority of antimalarial drugs are of plant origin<sup>11</sup>. Several plants with anti-plasmodial properties have been proved as sources of novel anti-plasmodial compounds. Because the continuous resistance of malaria parasites to synthetic drugs like chloroquine is a major cause for concern<sup>12</sup> and due to the limited availability and affordability of orthodox medicine in many tropical countries, most indigenous populations depend on traditional medical remedies<sup>13</sup>, mainly

from plants that have been discovered to be a rich source of novel drugs<sup>14</sup>. Botanicals have played a significant role in human societies throughout history and prehistory<sup>14</sup>. In the tropical regions of Asia, South America, and Africa, where malaria is endemic, there is still widespread use of medicinal plants to treat the disease. It has been reported that *B. sapida*, among other plants, is used in southwest Nigeria for malaria treatment<sup>15</sup>.

*Khaya senegalensis* and *Khaya grandifoliola* stem and bark have been extensively used as antimalarial remedies but with adverse effects<sup>16</sup>. *Alstoniaboonei* (Cheese wood) has been a highly valuable plant, especially in situations where affordable antimalarial drugs are found ineffective due to drug-resistant malaria parasites<sup>17</sup>. Several other natural products having antimalarial activities have been documented in Nigeria<sup>18</sup>. Before any plant can be recommended, there is a need for scrutiny because there are indications of deleterious effects of many phytochemicals on human health<sup>19</sup>.

Apart from contemporary antimalarial drugs that have been used for years and the conventional traditional herbs, artemisinin-based combination therapy (artemether and lumefantrine) is very active<sup>20, 21</sup> in the treatment of malaria caused by *Plasmodium falciparum* that is resistant to chloroquine, with common side effects which may include muscle and joint pains, fever, loss of appetite and headaches<sup>22</sup>.

The need for an alternative drug has initiated intensive efforts to develop new antimalarial drugs from indigenous plants. Various studies have been documented with over 1200 plant species from 160 families used to treat malaria or fever<sup>23</sup>. Research and investigations have been carried out in many African nations such as Ethiopia<sup>24</sup>, Ghana<sup>25</sup>, Cameroon<sup>26</sup> and Nigeria<sup>27, 18, 28, 29</sup>.

*Blighia sapida* is indigenous to the forests of most West African countries, where the fruits are rarely eaten but are used for other purposes<sup>30</sup>. The fruit is edible when ripe but toxic when unripe due to hypoglycin A and hypoglycin B. Ackee arils have been reported to have a comparable proximate composition of moisture, ash, carbohydrate, crude protein and crude fiber to many known legumes

and oil seeds<sup>31, 32</sup>. In Ghana, several uses are recorded for *B. sapida*; the bark is one of the ingredients in a concoction administered for epilepsy. The leaf juice is used for washing or as drops for sore eyes, conjunctivitis, and trachoma<sup>33</sup>. The bark and leaf decoctions are administered to treat oedema, intercostal pain, dysentery, diarrhoea, chest pain<sup>34</sup>, malaria<sup>15</sup> in South-Western Nigeria. The pulp of twiggy leaves is applied on the forehead to treat migraine/headache. The ash obtained from dried husks and seeds is used for soap making in West Africa and the fruits contain saponin, which lathers in water and is used for washing<sup>35</sup>. *B. sapida* has also been used by the Centre for Scientific Research into Plant Medicine (CSRPM), Ghana, for the treatment of diarrhoea for over twenty years<sup>36</sup>. The leaves and seeds of the plant have also been reported by the people of Itak Ikot Akap Community, in Ikono Local Government Area of Akwa Ibom State, Nigeria to possess piscicidal properties<sup>37</sup>.

#### **Phytochemical Screening of *Blighia sapida*:**

Alkaloids, saponins, tannins, phlobatannins, flavonoids, terpenes, cardiac glycosides, and combined anthraquinones were detected in the leaf of *B. sapida*, whereas only saponins, flavonoids, combined anthraquinones, and cardiac glycosides were detected in the stem bark<sup>31</sup>. It has also been reported that *B. sapida* tested positive for the presence of some groups of phytochemicals such as reducing sugars, phytosterols and Polyphenol<sup>38</sup>.

**Research Objectives:** Despite the recent successes recorded with the development of a malaria vaccine<sup>39</sup>, therapeutic interventions must be encouraged, and plants have presented a vast array of beneficial chemicals in the treatment of diseases. To this end, mankind must stay ahead of the malaria disease by presenting a wide array of interventions. Therefore, this study was conducted to assess the antimalarial property of *Blighia sapida* leaf extracts compared to Artemether-Lumefantrine, a standard antimalarial drug in continuation of the search for cheap and affordable treatments for malaria.

#### **MATERIALS AND METHODS:**

**Collection and Preparation of the Plant Material:** Leaves of *Blighia sapida* were obtained from a single tree identified in the Department of Plant Science and Biotechnology, Adekunle Ajasin

University, Akungba-Akoko, Ondo State, Nigeria, and the herbarium specimen was deposited at the Herbarium Unit of the Department. The harvested leaves were washed thoroughly with tap water and air-dried at room temperature on the laboratory bench. The dried leaves were milled using an electric blender and stored in specimen bags.

**Extraction of Plant Materials:** A decoction was made by placing 100g of the milled sample in 2 liters of boiling water. The supernatant was collected and concentrated, and the filtrate was lyophilized in the laboratory. The dry extract was collected and stored in a moisture-free container for use. Another portion of the milled leaves weighing 100g was also measured into 2 liters of absolute ethanol, stirred continuously, and left for 72 h. The filtrate was separated using Whatman filter paper (18cm diameter and 0.1 µm pore diameter). The filtrate was then dried in an oven at 40°C and stored in a vial at 4°C in a refrigerator.

#### **Experimental Animals and Malaria Parasite:**

Eight-week-old Swiss albino mice used for this study were obtained from the Animal Unit of the Institute for Advanced Medical Research and Training (IAMRAT), College of Medicine, University of Ibadan, Nigeria. The animals were kept in well-aerated wired cages, fed with standard mouse feed (Top Feeds Ltd, Nigeria), and allowed to drink water freely for one week before they were infected with *Plasmodium berghei* NK 65, donated by the Institute for Advanced Medical Research and Training (IAMRAT), College of Medicine, University of Ibadan, Nigeria. The animals were infected by the serial passage of blood collected from a donor mouse to the naïve recipients. This study was undertaken in line with the internationally accepted laboratory animal use and care guideline<sup>40</sup>.

**In-vivo Antimalarial Assay:** The animals were infected intraperitoneally with an aliquot of 0.2ml of standard inoculum ( $1 \times 10^7$  *Plasmodium berghei* strain NK 65 parasitized erythrocytes). The infected blood was obtained from a donor mouse with a parasitemia of 35 to 40% by heart puncture into a test tube having anticoagulant (0.5% trisodium citrate). The blood was then diluted with normal saline (0.9% NaCl). Thirty Mice weighing between 28g to 34g were distributed evenly into six

groups. Group 1 (control) was not given any antimalarial treatment. Group 2 was treated with 6.0mg/kg body weight of Artemether-Lumefantrine, Groups 3 and 4 were treated with 50 mg/kg body weight of ethanolic and aqueous leaf extract of *B. sapida*, respectively. Groups 5 and 6 were treated with 30 mg/kg body weight of ethanolic and aqueous leaf extract of *B. sapida*, respectively. All the treatments were administered once daily by gavages using an intubator for five consecutive days. Observation for changes in parasitaemia level began twenty-four hours after parasitising the animals and ended twenty-four after the last treatment. Blood was taken daily from the tail vein of the mice for six days.

**Drugs and Reagents:** Artemether 80mg - Lumefantrine 480mg (GETZ Pharma PVT Ltd. Pakistan), Giemsa (BioLab Diagnostics Ltd., India), Ethanol (Nosak Distilleries Limited, Nigeria), Trisodium citrate (Annexe Chem Private Limited, India) and normal saline (Biomedical Limited, Nigeria) were used. All reagents were analytically graded and procured from legal pharmaceutical suppliers.

**Parasitological Study:** Blood films were prepared from blood collected from each mouse over six days. The slides were screened for malaria parasites using 10% Giemsa's stain at pH 7.2. The number of parasites counted per 200 white blood cells was recorded and used to calculate parasite density based on 8000 assumed leucocytes/ $\mu$ l of blood as previously described<sup>41</sup>.

**Statistical Analysis:** The one-way Analysis of Variance (ANOVA) was used to analyze the differences at  $P < 0.05$  among the treatments (SPSS 23.0, SPSS Inc., Chicago, Illinois, USA). The results were expressed as mean  $\pm$  Standard Error (SE).

**RESULTS AND DISCUSSION:** Treatment of *Plasmodium berghei* infected mice elicited a change in parasitaemia. A reduction was regarded as an indicator of the antimalarial properties of any compound. Results obtained in this study **Fig. 1** and **2** indicated a reduction in parasitaemia induced by *B. sapida* extracts. The mean $\pm$ SE for parasitaemia count in mice treated with 6.0mg/kg body weight of Artemether-Lumefantrine

significantly reduced from 528 $\pm$ 44.54 to 60 $\pm$ 8.94 over six days' observation and treatment period. Mice treated with the ethanolic leaf extract of *B. sapida* at 50mg/kg and 30mg/kg body weight had parasitaemia suppressed from 608 $\pm$ 23.32 to 248 $\pm$ 14.97 and 536 $\pm$ 27.13 to 320 $\pm$ 28.2 respectively, while the mean $\pm$ SE value of parasitaemia count in the group treated with 50mg/kg and 30mg/kg body weight of Aqueous leaf extract of *B. sapida* was suppressed from 584 $\pm$ 20.40 to 328 $\pm$ 14.97 and 560 $\pm$ 52.15 to 336 $\pm$ 24.00 respectively. This indicates that Artemether-Lumefantrine significantly expressed an effect on the red blood cell count/parasitaemia count<sup>42</sup>.


Limited availability and affordability of orthodox medicine in many tropical countries have left much of the population depending on traditional medical remedies<sup>13</sup>. Of the many diseases that afflict tropical countries where the greater majority of the world's poor reside, malaria remains a major global health problem with high mortality and morbidity than any other infection<sup>2</sup>.

This is further exacerbated by its high degree of transmissibility. A concerted international effort has been put into obtaining a cure for this deadly disease culminating in the recent breakthrough in vaccine therapy<sup>39</sup>. This, however, does not prevent the need for further chemotherapeutic interventions<sup>43</sup>. There has been a remarkable help in administering several compounds, but progress has been challenging due to the resilience of the causative parasite in developing resistance to available medications<sup>44</sup>. Chloroquine had been the drug of choice for several years until widespread resistance demanded the search for more potent formulations<sup>45</sup>.


Therefore, the race is on to identify natural remedies for the treatment of malaria. Currently, the standard drug is the Artemether-Lumefantrine combination which was formulated from artemisinin extracted from *Artemisia annua*<sup>46</sup> a member of the Asteraceae family. *Blighia sapida*, has been identified as one of the local plants commonly used in southwest Nigeria for malaria treatment<sup>15</sup>. It is imperative to seek comparison with the standard treatment in use if the effects are to be measured appropriately.

This is why this study focused on the comparative effects of leaf extracts of *B. sapida*, with the standard Artemether-Lumefantrine combination. Although the results obtained in Fig. 1 show that leaf extracts of *Blighia sapida*, as well as the

Artemether-Lumefantrine combination, effectively reduced the *P. berghei* parasite level, the leaf extracts of *Blighia sapida* were not as efficacious as the Artemether-Lumefantrine combination.



**FIG. 1: COMPARISON BETWEEN THE EFFICACY OF THE 50MG/KG PLANT EXTRACT TREATMENTS AND THE STANDARD DRUG ON P. BERGHEI.** The graph compares the mean value and standard error difference for all treatment groups. The effects of ethanolic and aqueous leaves extract at 50mg/kg body weight compared to 6mg/kg body weight of Artemether-Lumefantrine (Standard) and the control group. Control = control group, A-L = group treated with Artemether-Lumefantrine, ETH-50 = group treated with 50mg/kg body weight of ethanolic extract, AQ-50 = group treated with 50mg/kg body weight of aqueous extract.



**FIG. 2: COMPARISON BETWEEN THE EFFICACY OF THE 30MG/KG PLANT EXTRACT TREATMENTS AND THE STANDARD DRUG ON P. BERGHEI.** The graph compares the mean value and standard error difference for all treatment groups. The effects of ethanolic and aqueous leaves extract at 30mg/kg body weight compared to 6mg/kg body weight of Artemether-Lumefantrine and the control group. CNTRL = control group, A-L = group treated with Artemether-Lumefantrine, ETH-30 = group treated with 30mg/kg body weight of ethanolic extract, AQ-30 = group treated with 30mg/kg body weight of aqueous extract.

**TABLE 1: EFFECT OF THE TREATMENTS ON PARASITAEMIA IN FIVE-DAY SUPPRESSIVE TEST**

Treatment	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6
CNTRL	0	+7.4627	+10.4480	+19.4030	+28.3580	+41.7910
A-L	0	-13.6364	-31.8182	-54.5455	-68.9394	-88.6364
ETH-50	0	-6.5790	-17.1053	-30.2632	-46.0526	-59.2105
AQ-50	0	-6.8493	-13.6986	-21.9178	-31.5068	-43.8356
ETH-30	0	-5.6338	-15.4930	-22.5352	-32.3944	-43.6620
AQ-30	0	-10.0000	-17.1429	-24.2857	-34.2857	-40.0000

The table shows percentage (%) parasitaemia level change for all groups. CNTRL = control group, A-L = standard drug group treated with Artemether-Lumefantrine, ETH-50 = group treated with 50mg/kg body weight of ethanolic extract, AQ-50 = group treated with 50mg/kg body weight of aqueous extract, ETH-30 = group treated with 30mg/kg body weight of ethanolic extract, AQ-30 = group treated with 30mg/kg body weight of aqueous extract.

Percentage of parasitemia increase and decrease =  $\frac{\text{initial} - \text{final parasitemia level}}{\text{Initial parasitemia level}} \times 100$

The results in **Table 1** indicate that the effect of the extracts on *P. berghei* was concentration-dependent, although the threshold was not determined. The ethanolic extract exhibited greater effectiveness than the aqueous extract; this can be attributed to the solvent potency to extract more phytochemicals from medicinal plants than its aqueous counterpart<sup>47, 48</sup>.

The percentage parasitemia count as shown in **Table 1** shows that the group of infected mice treated with 6.0mg/kg body weight of Artemether-Lumefantrine combination reflects a significant reduction of 88.63% in the parasitaemia level of the mice, which was significantly better than reduction obtained with the leaf extracts. Treatment with 50mg/kg body weight of ethanolic leaf extract of *B. sapida*, elicited a reduction of up to 59.21% in the parasitemia level which was better than the 43.84% reduction obtained from the aqueous leaf extract of *B. sapida*. At the lower dosage of 30mg/kg body weight, the ethanolic and aqueous extracts had comparable suppression levels of 43.66% and 40%, respectively, whereas the controls had a 41.79% increase in parasite density.

This agrees with previous study<sup>49</sup>, which reported that parasitemia level increases progressively after inoculation until the point of death in the absence of suitable treatment. Although the results obtained were encouraging, this study indicates that leaf extracts of *Blighia sapida*, although efficacious in reducing parasitemia, did not show greater potency than the Artemether-Lumefantrine combination. However, there is a need for further studies to explore the use of other extracting solvents and to determine the response to increased levels of the extracts in concert with toxicological studies.

**CONCLUSION:** Based on the results, this study confirms that the ethanolic and aqueous leaf extracts of *Blighia sapida* have promising anti-malaria activity at the concentrations used. The results lend credence to the ethnobotanical use of the plant for the treatment of malaria. Further research using higher concentrations of the ethanolic and aqueous leaf extracts of the plant, and/or using other solvents can be employed to test the efficiency in treating malaria disease.

**ACKNOWLEDGEMENT:** We acknowledge the help of the Central Laboratory of the Federal University of Technology, Akure, Ondo State, Nigeria, in lyophilizing the aqueous extracts of the plant materials and the Animal Care Unit of the Institute for Advanced Medical Research and Training (IAMRAT), College of Medicine, University of Ibadan, Nigeria for providing the parasitized laboratory animals used for this study.

**Funding:** This research received no external funding.

**Ethics Approval:** Ethical clearance to experiment on mice was obtained from the Centre for Research and Development, Adekunle Ajasin University, Akungba-Akoko, Nigeria. The mice were handled in accordance with “The care and Use of Animals for Scientific purposes” guidelines of the Animal Use and Care Committee (AUCC) and the legal provisions captured in the constitution of the Federal Republic of Nigeria, Animal Diseases (Control) Act. Cap A17 LFN, 2004.

**CONFLICT OF INTEREST:** Nil

## REFERENCES:

1. WHO: World Malaria Report. Geneva. World Health Organization 2015.
2. WHO: World Malaria Report. Geneva. World Health Organization 2011.
3. Akanbi OM, Omonkhu AA and Cyril-Olutayo CM and Fasimoye RY: The antiplasmodial activity of *Anogeissus leiocarpus* and its effect on oxidative stress and lipid profile in mice infected with *Plasmodium berghei*. Parasitology Research 2012; 110(1): 219-226.
4. Jimoh A, Sofola O, Petu A and Okorosobo T: Quantifying the economic burden of malaria in Nigeria using the willingness to pay approach. Cost Effectiveness and Resources Allocation 2007; 5: 6-13.
5. Okokon JE, Antia BS, Igboasoiji, AC, Essien EE and Mbagwu, HO: Evaluation of anti-plasmodial activity of ethanolic seed extract of *Picralimanitida*. Journal Ethnopharmacol 2007; 111(3): 464-467.
6. Aja PM, Odeh CO, Uraku AJ and Ofor CE: Evaluation of antioxidant activities of ethanol leaf extract of *Cymbopogon citratus* and *Hyptis spicigera* in mice exposed to *Plasmodium berghei*. International Journal of Current Microbiology and Applied Science 2015; 4(2): 382-393.
7. Amino R, Menard R and Frischknecht F: *In-vivo* imaging malaria parasites-recent advances and future directions. Current Opinion in Microbiology 2005; 8: 407-414.
8. Cox FE: History of the discovery of the malaria parasites and their vectors. Parasites Vectors 2010; 3(5): 1-9. <<https://doi.org/10.1186/1756-3305-3-5>>
9. Elujoba AA, Odeleye OM and Ogunyemi CM: Traditional Medical Development for medical and Dental Primary Health Care Delivery System in Africa. African Journal of Traditional, Complementary and Alternative Med 2005; 2: 46-61.

10. Okigbo RN and Mmeka EC: An appraisal of phytomedicine in Africa. *Ladkrabang Science and Technology Journal* 2006; 6: 83–94.
11. Chiyaka C, Garira W and Dube S: Effects of treatment and drug resistance on the transmission dynamics of malaria in endemic areas. *Theoretical Population Biology* 2009; 75: 14–29.
12. Trape JF: Combating malaria in Africa. *Trends in Parasitology* 2002; 18: 224-230.
13. Bankole AE, Adekunle AA, Sowemimo AA, Umebese CE, Abiodun O and Gbotosho GO: Phytochemical screening and *in-vivo* antimalarial activity of extracts from three medicinal plants used in malaria treatment in Nigeria. *Parasitology Research* 2016; 115(1): 299–305.
14. Lewis WH and Elvin-Lewis MP: *Plants Affecting Human. Edition 2<sup>nd</sup> Medical Botany* 2003; 812.
15. Oladele AT and Adewunmi CO: Medicinal plants used in the management of malaria among the Traditional medicine practitioners (TMP's) in South-Western Nigeria. *African Journal of Infectious Diseases* 2008; 2(1): 51-59.
16. Adebayo JO, Yakubu MT, Egwim EC, Owoyele BV and Enaibe BU: Effect of ethanolic extract of *Khaya senegalensis* on some biochemical parameters of rat kidney. *Journal of Ethnopharmacology* 2003; 88: 69–72.
17. Majekodunmi SO, Adegoke OA and Odeku OA: Formulation of the extract of the stem bark of *Alstoniabooneias* tablet dosage form. *Tropical Journal of Pharmaceutical Research* 2008; 7: 987–994.
18. Bode AM and Dong Z: Toxic phytochemicals and their potential risks for human cancer. *Cancer prevention research (Philadelphia, Pa.)* 2015; 8(1): 1–8. <https://doi.org/10.1158/1940-6207>.
19. Idowu OA, Soniran OT, Ajana O and Aworinde DO: Ethnobotanical survey of antimalarial plants used in Ogun State, Southwest Nigeria. *African Journal of Pharmacy and Pharmacology* 2009; 4(2): 055–060.
20. Makanga M and Krudsood S: The clinical efficacy of artemether-lumefantrine (Coartem®). *Malaria Journal* 2009; 8: 1-12
21. Mueller M: Efficacy and safety of the six-dose regimen of artemether–lumefantrine for treatment of uncomplicated *Plasmodium falciparum* malaria in adolescents and adults: A pooled analysis of individual patient data from randomized clinical trials. *Acta Tropical* 2006; 100(1-2): 41–53.
22. Morita H, Wu J and Zipes DP: The QT syndromes: long and short. *Lancet* 2008; 372(9640): 750–763.
23. Willcox M, Burford G and Bodeker G: An overview of ethnobotanical studies on plants used for the treatment of malaria. *Tradit Med Plants and Malaria* 2004; 187-197.
24. Bekalo TH, Woodmatas SD and Woldemariam ZA: An ethnobotanical study of medicinal plants used by local people in the lowlands of Konta Special Woreda, southern nations, nationalities and peoples regional state. *Ethiopia J of Ethnobiology and Ethnomedicine* 2009; 5(1): 1-15.
25. Asase A, Oteng-Yeboah AA, Odamten TG and Simmonds MSJ: Ethnobotanical study of some Ghanaian antimalarial plants. *Journal of Ethnopharmacology* 2005; 99: 273–279.
26. Titanji VPK, Zofou D and Ngemenya MN: The antimalarial potential of medicinal plants used for the treatment of malaria in Cameroonian folk medicine. *African Journal of Traditional Complementary and Alternative Medicine* 2008; 5: 302–321.
27. Ajibesin KK, Ekpo BA, Bala DN, Essien EE and Adesanya SA: Ethnobotanical survey of Akwa Ibom State of Nigeria. *Journal of Ethnopharmacology* 2008; 115(3): 387–408.
28. Kayode J, Ige OE, Adetogo TA and Igbakin AP: Conservation and biodiversity erosion in Ondo State, Nigeria: Survey of Plant barks used in native pharmaceutical extraction in Akoko region. *Ethnobotanical Leaflets* 2009; 13: 665–667.
29. Dike PI, Obembe OO and Adebisi EF: Ethnobotanical survey for potential antimalarial plants in South-Western Nigeria. *J of Ethnopharmacology* 2012; 144(3): 618–626.
30. Ubulom, Peace ME, Udobi CE, Akpabio EI and Eshiet U: Antimicrobial Activities of Leaf and Stem Bark Extracts of *Blighia sapida*. *Journal of Plant Studies* 2013; 2(2): 47-52.
31. Oyeleke GO, Oyetade OA, Afolabi F and Adegoke BM: Nutrients, antinutrients and physicochemical compositions of *Blighia sapida* pulp and pulp oil (ackee apple). *Journal of Applied Chemistry* 2013; 4(1): 05-08.
32. Blake OA, Jackson JC, Jackson MA and Gordon CA: Assessment of dietary exposure to the natural toxin hypoglycin in ackee (*Blighia sapida*) by Jamaican consumers. *Food Research Inter* 2004; 37(8): 833-838.
33. Iyang E: *Ethnobotany: Conventional and traditional uses of plants.* The Verdict Press Uyo 2003; 1: 191.
34. Ige OE: Preliminary investigations on the ethnomedicinal plants of Akoko Division, South West Nigeria. *Global Journal of Health Science* 2011; 3(2): 84–89.
35. Orwa C, Mutua A, Kindt R, Jamnadass R and Anthony S: *Agroforestry Database: a tree reference and selection guide version 4.0.* Diaksestanggal 2009; 10(9): 2016. <<http://www.worldagroforestry.org/sites/treedbs/treedatabases.asp>>
36. Antwi S, Martey ONK, Donkor K, Nii-Ayitey and Okine LK: Diarrhoeal activity of *Blighia sapida* (Sapindaceae) in rats and mice. *J of Pharma and Toxi* 2009; 4(3): 117-125.
37. Peace MEU, Godwin N, Imandeh Ette O, Ettebong and Chinweizu EU: Potential Larvicidal Properties of *Blighia sapida* Leaf Extracts against Larvae of *An. Gambiae*, *Cu. Quinqüefasciatus* and *Ae. Aegypti*. *British Journal of Pharmaceutical Research* 2012; 2(4): 259-268.
38. Ajayi AF and Ajayi LO: Ethanolic Extract of *Blighia sapida* Improved Male Wistar Rats Reproductive Variables. *Res J of Medical Sciences* 2014; 8(3): 109-115.
39. World Health Organization. WHO recommends groundbreaking malaria vaccine for children at risk. *World Health Organization* 2021. <<https://www.who.int/news/item/06-10-2021-who-recommends-groundbreaking-malaria-vaccine-for-children-at-risk>>(accessed 16 November 2021).
40. National Research Council (US) Committee for the Update of the Guide for the Care and Use of Laboratory Animals. *Guide for the Care and Use of Laboratory Animals.* 8th edition. Washington (DC): National Academies Press (US); 2011. Available from: <<https://www.ncbi.nlm.nih.gov/books/NBK54050/> DOI: 10.17226/12910>
41. Christiana I, Ifeoma O, Mercy A and Martins E: Antiplasmodial activity of the mixed stem bark extracts of *Anogeissus leiocarpus* and *Prosopis africana* and *in-vitro* evaluation of its tablet dosage form. *Journal of Herbs Spices and Medicinal Plants* 2011; 17(4): 419-435.
42. Walker SP, Washs TD, Gardner JM, Lozoff B, Wasserman GA and Pollitt E: Child development: risk factors for adverse outcomes in developing countries. *Lancet* 2007; 369(9555): 145–157.
43. Greenwood B, Cairns M, Chaponda M, Chico RM, Dicko A, Ouedraogo JB, Phiri KS, Ter Kuile FO and Chandramohan D: Combining malaria vaccination with chemoprevention: a promising new approach to malaria control. *Malaria Journal* 2021; 20(1): 1-7.

44. White NJ: Antimalarial drug resistance. The Journal of Clinical Investigation 2004; 113(8): 1084–1092.
45. Peter BB: Drug Resistance in Malaria. Malaria Epidemiology Branch, Centers for Disease Control and Prevention Chamblee, GA, USA, 2001. <<https://www.who.int/csr/resources/publications/drugresist/malaria.pdf>>
46. Premji ZG: Coartem®: the journey to the clinic. Malaria Journal 2009; 8(1): S3. <<https://doi.org/10.1186/1475-2875-8-S1-S3>>.
47. Cowan MM: Plant products as antimicrobial agents. Clinical Microbiology Reviews 1999; 12(4): 564–582.
48. Emad MA, Amna SK and Nazlina I: Antibacterial activity of oleo-gum resins of Commiphoramolmol and Boswellia papyrifera against methicillin resistant Staphylococcus aureus (MRSA). Scientific Research and Essay 2009; 4(4): 351-356.
49. Breman JG: Ears of the hippopotamus: manifestations, determinants and estimates of the malaria burden. in: Breman JG, Egan A and Keusch GT (eds.): The intolerable burden of malaria: a new look at the numbers: of the American Journal of tropical medicine and hygiene. Supplement to the American Society of Tropical Medicine and Hygiene 2001; 64(1): 1-4.

**How to cite this article:**

Olugbenga OV, Odukamaiya, Odubajo S and Sunday, Damilola M: Anti-plasmodial activity of *Blighia sapida* leaf extracts in *Plasmodium berghei* infected mice. Int J Pharmacognosy 2022; 9(1): 17-24. doi link: [http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.9\(1\).17-24](http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.9(1).17-24).

This Journal licensed under a Creative Commons Attribution-Non-commercial-Share Alike 3.0 Unported License.

This article can be downloaded to **Android OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)