



Received on 24 June, 2017; received in revised form, 07 July, 2017; accepted, 17 September, 2017; published 01 December, 2017

## EVALUATION OF ANTIOXIDANT, ANTI-INFLAMMATION AND CHEMICAL COMPOSITION OF METHANOLIC EXTRACT OF *AMOMUM MASTICATORIUM* THWAITES (ZINGIBERACEAE)

K. Sinitha\* and J. E. Thoppil

Cell and Molecular Biology Division, Department of Botany, University of Calicut - 673635, Kerala, India.

### Keywords:

*Amomum masticatorium*,  
Anti-inflammation, Antioxidant

### Correspondence to Author:

**K. Sinitha**

Cell and Molecular Biology Division,  
Department of Botany, University of  
Calicut - 673635, Kerala, India.

**E-mail:** sinithak@gmail.com

**ABSTRACT:** The anti-inflammatory and antioxidant activities of the methanol extract of *Amomum masticatorium* rhizome were observed in various experimental models. Phytochemical constituents like phenols and flavonoids were also estimated. The antioxidant activity was evaluated by DPPH and super oxide radical scavenging assays. The extract showed significant antioxidant activity. The *in vitro* anti-inflammatory activity of the extract was investigated by various methods *viz.* albumin denaturation, membrane stabilization and proteinase inhibitory assays. Results showed that, the extract exhibited significant anti-inflammatory activity by inhibiting the heat induced albumin denaturation with  $IC_{50}$  value  $34.15 \pm 0.404 \mu\text{g/ml}$ , effectively stabilized RBC membrane with  $IC_{50}$  value of  $51.47 \pm 1.56 \mu\text{g/ml}$  and significantly inhibited proteinase activity with  $IC_{50}$  value of  $52.8 \pm 0.61 \mu\text{g/ml}$ . *In vivo* activity was evaluated in carrageenan and formalin induced paw edema test in mice and compared with that of standard drug diclofenac. In both the models the extract showed significant reduction in edema. The extract at 200mg/kg b. wt showed significant reduction in carrageenan induced paw edema at 3<sup>rd</sup> hour and chronic edema on 6<sup>th</sup> day. From the results, it is concluded that, methanol extract of *Amomum masticatorium* rhizome exhibited significant anti-inflammatory and antioxidant activity, which may be due to the presence of active phytochemicals in it.

**INTRODUCTION:** Inflammation is a protective response of cell which involves several mechanisms and complex array of enzyme activation, mediator release, cell migration, tissue breakdown and repair. Inflammation results in the release of endogenous mediators like histamine, serotonin, bradykinin and prostaglandin <sup>1</sup>.

It plays a major role in most chronic illnesses like neurodegenerative, cardiovascular, pulmonary, metabolic, autoimmune and neoplastic diseases <sup>2</sup>. Sometimes inflammation may lead to events that are quite serious and become chronic, like occurrence of rheumatoid arthritis <sup>3</sup>. Many of the anti-inflammatory drugs available have many side effects which impose a growing concern for more natural alternatives. *Amomum masticatorium* of Zingiberaceae is found as under growth in the evergreen forest at high altitudes. It has slender stoloniferous rhizomes. The rhizome of this plant is aromatic and carminative. It is used to chew with betel in Sri Lanka <sup>4</sup>.



Several species of *Amomum* has traditional medicinal uses. Some are prescribed for the treatment of indigestion, vomiting, abdominal pains and rectal diseases. They are also ingredients of Ayurvedic and Unani medicines. A closely related species *Amomum subulatum* have reported to possess valuable properties like allopathic, analgesic, anti-inflammatory, antimicrobial, antioxidant, antiulcer, cardio-adaptogen and hypolipidaemic activities<sup>5</sup>.

## MATERIALS AND METHODS:

**Plant material:** The plant material was collected from Wagamon, Idukki district, Kerala, India. The material was herbarised (CALI 123766) and deposited at CALI, University of Calicut. Rhizome of the plant was chopped into small pieces, shade dried and powdered. The powdered material was run in soxhlet apparatus for 6 hrs with methanol for extraction. The extract was evaporated in vacuum to completely remove the methanolic content to get a final thick extract.

### Determination of Phytochemical Constituents:

**Total Phenolic Content:** The total phenolic content in the extract was determined by Folin-Ciocalteu assay<sup>6</sup>. Gallic acid was used as standard. The content of phenolic compounds of the sample was expressed as gallic acid equivalents in mg per gram dry weight (mg GAE/g DW). All the samples were analyzed in triplicates.

**Determination of Total Flavonoid Content:** The AlCl<sub>3</sub> method<sup>6</sup> was used for the determination of total flavonoid content of the methanolic plant extract. Quercetin was used as the standard. Total flavonoid content was expressed as quercetin equivalents in mg per gram dry weight (mg QE/g DW). All the determinations were performed in triplicates.

**Antioxidant Potential:** The antioxidant property of the extract was determined by superoxide radical scavenging by the NBT reduction method<sup>7</sup>. The optical density was measured at 560 nm and percentage inhibition was evaluated by comparing the absorbance value of the control and samples. DPPH radical scavenging potential was measured by the method proposed by Coruh *et al.*, 2007<sup>8</sup>. DPPH has an absorption peak at 515 nm which disappears on reduction by an antioxidant

compound. Extract was added to 1.5ml of freshly prepared DPPH solution and absorbance measured at 515 nm 20 min after the reaction was started. The percentage inhibition was calculated by comparing with that of the control.

### *In vitro* Anti-inflammation:

**Inhibition of Albumin Denaturation:** Inhibition of albumin denaturation was studied according to methods previously described<sup>9, 10</sup>. The reaction mixture containing test extracts of different concentrations (50µg, 100µg and 200µg/ml) and 5% aqueous solution of bovine albumin fraction were incubated at 37 °C for 20 min and then heated to 57 °C for 20 min. After cooling the samples, the turbidity was measured at 660 nm. The experiment was performed in triplicate. The percentage inhibition of protein denaturation was then calculated. Diclofenac was used as the standard.

**Proteinase Inhibition:** The potential for inhibiting proteinase was evaluated according to the method of Oyedepo *et al.*, (1995)<sup>11</sup>. The reaction mixture (2ml) with 0.06mg trypsin, 1ml 20mM Tris HCl buffer (pH 7.4) and 1ml test sample of different concentrations (50 – 200µg/ml) was incubated at 37 °C for 5 min. Then 1ml of 0.8% (w/v) casein was added and was incubated for an additional 20 min. 2ml of 70% perchloric acid was added to stop the reaction. Cloudy suspension was centrifuged and the absorbance of the supernatant was read at 210 nm against buffer as blank. The experiment was performed in triplicate. The percentage inhibition of proteinase inhibitory activity was calculated.

**Membrane Stabilization:** Preparation of Red Blood cells (RBCs) suspension was done according to the standard protocol<sup>12</sup>. The blood sample was collected and transferred to the centrifuge tubes. The tubes were centrifuged at 3000 rpm for 10 min and were washed three times with equal volume of normal saline. The volume of blood was measured and reconstituted as 10% v/v suspension with normal saline.

**Heat Induced Haemolysis:** The reaction mixture consisted of 1ml test sample of different concentrations (50 - 250µg/ml) and 1ml of 10% RBCs suspension, instead of test sample only saline was added to the control test tube<sup>13</sup>. Diclofenac was used as standard drug. The centrifuge tubes

were incubated in water bath at 56 °C for 30 min. The tubes were then cooled under running tap water. The reaction mixture was centrifuged at 2500 rpm for 5 min and the absorbance of the supernatants was read at 560 nm. The experiments were performed in triplicates. The percentage of inhibition of haemolysis was calculated.

#### **In vivo Anti - inflammation:**

**Animals:** Young male swiss albino mice were purchased from small animal breeding station, Kerala Agriculture University, Mannuthy, Kerala, India. They were maintained in well ventilated cages under controlled temperature and provided with 12 h light/dark cycle. The experimental protocol was approved by the Institutional Animal Ethical Committee.

#### **Carrageenan Induced Acute Inflammation:**


Male swiss albino mice were used for the experimental study. The animals were divided into four groups with five animals in each group. Group I was treated with carrageenan alone and was taken as the control. Group II was treated with standard drug diclofenac (10mg/ kg/ b. wt). Group III and IV were treated with different doses of the extract (50mg / kg and 200 mg/kg b. wt). The extract was administered orally for five consecutive days. One hour after the fifth dose of the extract, acute inflammation was induced in the right hind paw of the mice by sub plantar injection of 0.02ml freshly prepared 0.1% carrageenan in normal saline<sup>14</sup>. Paw edema volume was measured using vernier callipers. Reading was taken every 1 hr up to 6 hr, and finally reading at 24 hr was also taken.

**Formalin Induced Chronic Inflammation:** Male swiss albino mice were divided into four groups with five animals in each group. Group I was treated as the control. Group II was treated with standard drug diclofenac (10mg/ kg/ b. wt). Group III and IV were treated with different concentrations of the extract (50mg / kg and 200mg/kg b. wt). The extract was administered orally for five consecutive days. One hour after the fifth dose of the extract, chronic inflammation was induced in the right hind paw of the mice by sub plantar injection of 0.02ml of 2% formalin<sup>15</sup>. Paw edema thickness was measured using vernier callipers. Reading was taken for six consecutive days.

## **RESULTS AND DISCUSSION:**


### **Phytochemical Constituents:**

**Total Phenol Content:** The total phenolic content of *A. masticatorium* was determined by Folin-Ciocalteu assay<sup>16</sup>. A standard calibration curve for gallic acid was determined for a range of concentrations 20-120 mg/ml (**Fig. 1**). Total phenolic content of the extract was found to be  $89.54 \pm 4.67$ mg GAE/g dry weight of the extract.



**FIG. 1: STANDARD CALIBRATION CURVE OF GALLIC ACID**

**Total Flavonoid Content:** Total flavonoid content was determined by AlCl<sub>3</sub> method. Standard calibration curve of quercetin for a range of concentrations 20-100mg/ml was determined (**Fig. 2**). Total flavonoid content of the extract was found to be  $58.28 \pm 3.32$ mg QE/g dry weight of the extract.



**FIG. 2: STANDARD CALIBRATION CURVE OF QUERCETIN**

The total phenolic and flavonoid content of the extract was found to be in considerable high amounts. Flavonoids are used in the treatment of many common diseases. Their ability to inhibit specific enzymes, simulate some hormones and neurotransmitters, and scavenge free radicals is already proven<sup>17</sup>. Phenolic compounds are also reported to have several biological properties, including antioxidant activity<sup>18</sup>.




FIG. 3: GRAPH SHOWING ANTIOXIDANT ACTIVITY OF THE EXTRACT

**Antioxidant:** The methanolic rhizome extract has significant radical scavenging activity which exhibits its high antioxidant prospective. The IC<sub>50</sub> values are given in Table 1.

TABLE 2: PERCENTAGE INHIBITION OF THE EXTRACT IN VARIOUS ANTI-INFLAMMATORY ASSAYS

Treatment	Proteinase inhibition	Albumin denaturation	Membrane stabilisation
<i>A. masticatorium</i> 50µg/ml	31.56 ± 2.31 <sup>a</sup>	39.51 ± 1.68 <sup>a</sup>	33.86 ± 6.56 <sup>a</sup>
<i>A. masticatorium</i> 100µg/ml	48.43 ± 5.84 <sup>b</sup>	58.63 ± 2.22 <sup>b</sup>	55.30 ± 7.15 <sup>b</sup>
<i>A. masticatorium</i> 200µg/ml	92.33 ± 1.36 <sup>c</sup>	89.06 ± 0.84 <sup>c</sup>	91.72 ± 2.99 <sup>c</sup>
Diclofenac 100µg/ml	90.21 ± 1.22 <sup>c</sup>	91.83 ± 0.52 <sup>c</sup>	94.40 ± 1.55 <sup>c</sup>

Each value (mean ± S.E.) represents mean of three replicates. Means in a column followed by the same superscript letters are not significantly different (p < 0.05, one way ANOVA, DMR test).

TABLE 3: IC<sub>50</sub> OF THE EXTRACT IN *IN VITRO* ANTI-INFLAMMATORY ASSAYS

Assay	IC <sub>50</sub> value
Proteinase inhibition	52.80 ± 0.61
Albumin denaturation	34.15 ± 0.40
Membrane stabilisation	51.47 ± 1.56

The *in vitro* anti-inflammatory assays provide considerable evidence for the anti-inflammatory ability of the extract. Protein denaturation and inflammation are concomitantly related. Tissue damage during inflammatory reactions occurs due to the release of leukocytes proteinase<sup>21</sup>. Significant level of protection was provided by the extract which may have inhibited the activity of proteinase. The erythrocyte membrane is analogous to the lysosomal membrane<sup>22, 23</sup>. The extra cellular activity of lysosomal enzymes released during inflammation may lead to severe disorders. The extract might have stabilised lysosomal membrane thereby inhibiting the activity of these enzymes.

***In vivo* Anti-inflammation:** Carrageenan and formalin induced edema models are important assays for establishing the *in vivo* anti-inflammatory activity of the extract.

It shows a dose dependent increase in the activity in both DPPH and superoxide radical scavenging assays (Fig. 3). Antioxidants have many health benefits and are considered to be important nutraceuticals<sup>19</sup>. The antioxidant potential of the plant extract is mainly from the flavonoids which contain hydroxyl functional groups<sup>20</sup>.

TABLE 1: IC<sub>50</sub> OF THE EXTRACT IN ANTIOXIDANT ASSAYS

Assay	IC <sub>50</sub> value
Super oxide radical scavenging	32.59 ± 1.22
DPPH radical scavenging	46.0 ± 0.21

***In vitro* Anti-inflammation:** In all the three *in vitro* anti-inflammatory assays, the extract showed significant activity. The results are given in Table 2 and 3.

**Carrageenan Induced Inflammation:** The extract significantly reduced inflammation in a dose dependent manner (Fig. 4). The extract inhibited carrageenan induced edema formation by 51% at a concentration of 50mg/kg b. wt and by 68% at 200 mg/kg b. wt (Table 4). The extract showed significant inhibition starting at 3<sup>rd</sup> hour after the administration of the drug.




FIG. 4: GRAPH SHOWING THE EFFECT OF EXTRACT ON CARRAGEENEN INDUCED INFLAMMATION

**TABLE 4: EFFECT OF THE EXTRACT ON CARRAGEENAN INDUCED PAW EDEMA IN MICE**


Treatment groups	Initial paw edema thickness (mm)	Paw edema thickness at 3 <sup>rd</sup> hour (mm)	Difference in thickness (mm)	% inhibition
Control	2.15 ± 0.01 <sup>b</sup>	3.71 ± 0.03 <sup>c</sup>	1.56 ± 0.04 <sup>b</sup>	--
Diclofenac	2.20 ± 0.07 <sup>b</sup>	2.62 ± 0.16 <sup>b</sup>	0.04 ± 0.13 <sup>a</sup>	73 ± 8.39 <sup>b</sup>
<i>A. masticatorium</i> 50 mg/kg b. wt	1.89 ± 0.07 <sup>a</sup>	2.64 ± 0.07 <sup>b</sup>	0.75 ± 0.12 <sup>a</sup>	51 ± 7.42 <sup>b</sup>
<i>A. masticatorium</i> 200 mg/kg b. wt	1.84 ± 0.07 <sup>a</sup>	2.33 ± 0.07 <sup>a</sup>	0.49 ± 0.13 <sup>a</sup>	68 ± 7.92 <sup>b</sup>

Each value (mean ± S.E.) represents mean of five replicates. Means in a column followed by the same superscript letters are not significantly different (p < 0.05, one way ANOVA, DMR test).

**Formalin Induced Inflammation:** The extract also effectively inhibited formalin induced edema (Fig. 5). The extract at higher concentration showed 53.97% inhibition which was comparable with the effect of standard drug (Table 5).

Edema formation in the paw is the result of a synergism between various inflammatory mediators that increases vascular permeability and blood flow. The probable activity of the extract may be by the inhibition of inflammatory mediators such as histamine, serotonin and prostaglandin. Anti-inflammatory properties of plants are attributed to the presence of phenolics and flavonoids in them which is also responsible for their antioxidant

properties. ROS is scavenged by the antioxidants thus decreasing the disorders.

**FIG. 5: GRAPH SHOWING THE EFFECT OF EXTRACT ON FORMALIN INDUCED INFLAMMATION****TABLE 5: EFFECT OF THE EXTRACT ON FORMALIN INDUCED PAW EDEMA IN MICE**

Treatment groups	Initial paw edema thickness (mm)	Paw edema thickness at 6 <sup>th</sup> day (mm)	Difference in thickness (mm)	% inhibition
Control	2.08 ± 0.05 <sup>b</sup>	3.61 ± 0.08 <sup>c</sup>	1.53 ± 0.11 <sup>b</sup>	---
Diclofenac	1.77 ± 0.06 <sup>a</sup>	2.41 ± 0.08 <sup>a</sup>	0.64 ± 0.11 <sup>a</sup>	58.04 ± 6.95 <sup>a</sup>
<i>A. masticatorium</i> 50 mg/kg b. wt	1.94 ± 0.04 <sup>a,b</sup>	2.81 ± 0.1 <sup>b</sup>	0.87 ± 0.11 <sup>a</sup>	43.01 ± 7.12 <sup>a</sup>
<i>A. masticatorium</i> 200 mg/kg b. wt	1.84 ± .07 <sup>a</sup>	2.54 ± 0.09 <sup>a</sup>	0.70 ± 0.1 <sup>a</sup>	53.97 ± 6.39 <sup>a</sup>

Each value (mean ± S.E.) represents mean of three replicates. Means in a column followed by the same superscript letters are not significantly different (p < 0.05, one way ANOVA, DMR test).

**CONCLUSION:** The bioactivities of this species have been assessed for the first time. The extract has considerably high phenolic and flavonoid content. The therapeutic applications of flavonoids on inflammation have previously been reported<sup>24</sup>. Flavonoids and phenolics are also responsible for antioxidant activity. The anti-inflammatory property is highly related to the antioxidant capacity. This suggests that *A. masticatorium* may be a potential source of chemical compounds which can be utilised in pharmaceutical industry.

**ACKNOWLEDGEMENT:** The first author gratefully acknowledges University Grants Commission for the award of FIP for doing this research work.

**CONFLICT OF INTEREST:** The authors declare that there are no conflicts of interest.

#### REFERENCES:

- Vane JO and Botting R: Inflammation and the mechanism of action of anti-inflammatory drugs. The FASEB journal 1987; 1: 89-96.
- Buadonpri W, Wichitnithad W, Rojsitthisak P and Towiwat P: Synthetic curcumin inhibits carrageenan-induced paw edema in rats. Journal of Health Research 2009; 23: 11-16.
- Denko CW: A role for neuropeptides in inflammation. Biochemistry of inflammation. Springer Netherlands 1992; 177-181.
- Jayaweera DMA: Medicinal plants (indigenous and exotic) used in Ceylon: part IV 1982.
- Gautam N, Bhattarai RR, Khanal BKS and Oli P: Technology, Chemistry and Bioactive Properties of Large Cardamom (*Amomum subulatum* Roxb.): An Overview.

- International Journal of Applied Sciences and Biotechnology 2016; 4: 139-149.
6. Oueslati S, Ksouri R, Falleh H, Pichette A, Abdelly C and Legault J: Phenolic content, antioxidant, anti-inflammatory and anticancer activities of the edible halophyte *Suaeda fruticosa* Forssk. Food Chemistry 2012; 132: 943-947.
  7. McCord JM and Fridovich I: The utility of superoxide dismutase in studying free radical reactions I. radicals generated by the interaction of sulfite, dimethyl sulfoxide, and oxygen. Journal of Biological Chemistry 1969; 244: 6056-6063.
  8. Coruh N, Celep AS and Özgökçe F: Antioxidant properties of *Prangos ferulacea* (L.) Lindl., *Chaerophyllum macropodum* Boiss. and *Heracleum persicum* Desf. from Apiaceae family used as food in Eastern Anatolia and their inhibitory effects on glutathione-S-transferase. Food Chemistry 2007; 100: 1237-1242.
  9. Mizushima Y and Kobayashi M: Interaction of anti-inflammatory drugs with serum proteins especially with some biologically active proteins. Journal of Pharmacy and Pharmacology 1968; 20: 169-173.
  10. Sakat S, Juvekar AR and Gambhire MN: *In vitro* antioxidant and inflammatory activity of methanol extract of *Oxalis corniculata* Linn. International Journal of Pharmacy and Pharmaceutical Sciences 2010; 2: 146-155.
  11. Oyedepo OO and Femurewa AJ: Anti-protease and membrane stabilizing activities of extracts of *Fagra zanthoxiloides*, *Oxalis subscorpioides* and *Tetrapleura tetraptera*. International Journal of Pharmacognosy 1995; 33: 65-69.
  12. Sadique J, Al-Rqobahs WA, Bughaith MF and EI-Gindi AR: The bioactivity of certain medicinal plants on the stabilization of RBS membrane system. Fitoterapia 1989; 60: 525-532.
  13. Shinde UA, Kulkarni KR, Phadke AS, Nair AM, Dikshit Mungantiwar VJ and Saraf MN: Mast cell stabilizing and lipoxygenase inhibitory activity of *Cedrus deodara* (Roxb) Loud. wood oil. Indian Journal of Experimental Biology 1999; 37: 258-261.
  14. Adeyemi OO, Okpo SO and Ogunti OO: Analgesic and anti-inflammatory effects of the aqueous extract of leaves of *Persea americana* Mill (Lauraceae). Fitoterapia 2002; 73: 375-380.
  15. Roy A, Gupta JK and Lahiri SC: Further studies on anti-inflammatory activity of two potent indan-1-acetic acids. Indian Journal of Physiology and Pharmacology 1982; 26: 207-214.
  16. Ainsworth EA and Gillespie KM: Estimation of total phenolic content and other oxidation substrates in plant tissues using Folin-Ciocalteu reagent. Nature protocols 2007; 2: 875-877.
  17. Havsteen BH: The biochemistry and medical significance of the flavonoids. Pharmacology and Therapeutics 2002; 96: 67-202.
  18. Brown JE and Rice ECA: Luteolin-rich Artichoke extract protects low density lipoprotein from oxidation *in vitro*. Free Radical Research 1998; 29: 247-255.
  19. Lee J, Koo N and Min DB: Reactive oxygen species, aging and antioxidative nutraceuticals. Comprehensive reviews in food science and food safety 2004; 3: 21-33.
  20. Das NP and Pereira TA: Effects of flavonoids on thermal autooxidation of Palm oil: structure- activity relationship. Journal of the American Oil Chemist' Society 1990; 67: 255-258.
  21. Das SN and Chatterjee S: Long term toxicity study of ART-400. Indian Journal of Indigenous Medicines 1995; 16: 117-123.
  22. Gandhidasan R, Thamarachelvan A and Baburaj S: Anti-inflammatory action of *Lannea coromandelica* by HRBC membrane stabilization Fitoterapia 1991; 62: 81- 83.
  23. Shenoy S, Shwetha K, Prabhu K, Maradi R, Bairy KL and Shanbhag T: Evaluation of anti-inflammatory activity of *Tephrosia purpurea* in rats. Asian Pacific Journal of Tropical Medicine 2010; 3: 193-195.
  24. Middleton E, Kandaswami C, Theoharides TC: The effects of plant flavonoids on mammalian cells: implications for inflammation, heart disease and cancer. Pharmacological reviews 2000; 52: 673-751.

**How to cite this article:**

Sinitha K and Thoppil JE: Evaluation of antioxidant, anti-inflammation and chemical composition of methanolic extract of *Amomum masticatorium* Thwaites (Zingiberaceae). Int J Pharmacognosy 2017; 4(12): 413-18;.doi link: [http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.4\(12\).413-18](http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.4(12).413-18).

This Journal licensed under a Creative Commons Attribution-Non-commercial-Share Alike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)