



Received on 24 June, 2017; received in revised form, 11 July, 2017; accepted, 17 September, 2017; published 01 December, 2017

## EVALAUTION OF ANTHELMINTIC ACTIVITY OF *RICINUS COMMUNIS* LINN. LEAVES BY USING DIFFERENT TYPES OF SOLVENT

Nirmale Dnyaneshwar Mahadev\* and Balakshe Prajkta Vitthal

Yashoda Technical Campus, Satara Faculty of Pharmacy, Satara - 415015, Maharashtra, India.

### Keywords:

*Pheretima posthuma*,  
*Ricinus communis*, Anthelmintic  
activity, Helminthiasis

### Correspondence to Author:

Mr. Nirmale D. Mahadev

Assistant Professor,  
Yashoda Technical Campus, Satara  
Faculty of Pharmacy, Satara - 415015,  
Maharashtra, India.

E-mail: balakshep@gmail.com

**ABSTRACT:** Aqueous and ethanol ethyl acetate chloroform methanol extracts from the leaves of *Ricinus communis* were investigated for their anthelmintic activity against *Pheretima posthuma* and three concentrations (50, 75 and 100mg/ml) of each extract were studied for activity, in which paralysis and death time of the worm. Both the extract exhibited remarkable anthelmintic activity at all three concentrations. Albendazole suspension is used as standard reference (20mg/ml) and distilled water as control. Therefore the aqueous and ethanol methanol extract of leaves of *Ricinus communis* has been demonstrated for the first time.

**INTRODUCTION:** Helminthiasis or worm infection is one of the most prevalent diseases in the World<sup>1</sup>. The disease is highly prevalent particularly in third world countries due to poor management practices<sup>2</sup>. The word Helminthes has been derived from the Greek which means 'worm'. Helminthes infections are among the most common infections in man, affecting a large proportion of the world's population. The parasites can be acquired by contact with a) infected water b) infected meal c) infected animal<sup>3</sup>. In developing countries they pose a large threat to public health, and contribute to the prevalence of malnutrition, anemia, eosinophilia, and pneumonia<sup>4</sup>. Helminthiasis is a macro parasitic disease of humans and animals in which a part of the body is infested with parasitic worms such as pinworm, roundworm, or tapeworm<sup>5</sup>.

The World health Organization (WHO) estimated that 80% of the population of developing countries relies on traditional medicines, mostly plant drugs for their primary health care needs. The use of medicinal plant is growing worldwide because of the increasing toxicity and allergic manifestations of the synthetic drugs. Hence there is an increasing demand towards natural Anthelmintic<sup>6</sup>.

*Ricinus communis* also known as Castor oil plant belongs to family Euphorbiaceae. This plant is native of India, and it has spread thence over all the warmer countries of the world<sup>7</sup>. It is an annual or perennial bush or occasionally a soft-wooded small tree grown up to 6m or more, found nearly throughout India, Brazil, South Africa and Russia<sup>8</sup>. In the Indian system of medicine, the leaf, root and seed oil of this plant have been used for the treatment of inflammation, liver disorders, hypoglycemic and laxative<sup>9</sup>. Castor oil is widely used as a cathartic, and also for lubrication and illumination. Roots are administered in the form of a decoction for lumbago and in the form of a paste for toothache<sup>7</sup>. Stem of *Ricinus communis* have anticancer, antidiabetic and antiprotozoal activity<sup>10</sup>.

	<b>DOI:</b> 10.13040/IJPSR.0975-8232.IJP.4(12).419-21
	<b>Article can be accessed online on:</b> www.ijournal.com
<b>DOI link:</b> <a href="http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.4(12).419-21">http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.4(12).419-21</a>	

Literature review indicates that the Anthelmintic activity on the leaves of *Ricinus communis* has not been evaluated so far and thus this paper reports the Anthelmintic activity on the extract of leaves of *Ricinus communis* L. against earthworms.

**Authentication of Plant Material:** The fresh leaves of *Ricinus communis* were collected from the Malshiras District Solapur, Maharashtra and authenticated head of department of Yashvantrao chavan institute of sciences, Satara.

**Preparation of Extract:** The leaves of *Ricinus communis* was subjected to shade drying. The dried bark was further crushed into powder and then stored in air tight containers for further analysis. Crude extract of leaves was prepared by ultra sonication extraction method. About 125mg of powdered plant material was uniformly packed into a thimble and extracted with solvents (2 liters each). Solvents used were ethanol, water, ethyl acetate chloroform methanol then the beaker are keep in sonicator for sonication extraction for time 60 minutes respectively. After the sonication filter the extract then collect in different Petri plates and keep the petriplates for evaporation at room temperature. Then collect all the extract keep in refrigerator.

**Standard used for the Activity:** Albendazole suspension in the concentration of 20mg/ml mfg. by Cipla Ltd. used as reference standard.

**Animals:** Earthworms (*Pheretima posthuma*) of about 4-5cm long were used for anthelmintic activity, collected from Department of Agriculture, Satara.

**Method:** All the glass wares previously washed were sterilized in hot air oven. Petri-dishes,

pipettes, test tube were wrapped separately in the paper and kept in the hot air oven for sterilization at 180 °C for 1h. Take 10gm of powdered drug of night jasmine in 4 different beakers and then add 50ml of ethanol in each baker. Then the beaker is kept in sonicator for sonication extraction for time 60 minutes. The Anthelmintic assay was carried out as per the method of Deore *et al.*, with minor modifications<sup>12</sup>. Earthworms were used for Anthelmintic activity grouped into control, standard, and aqueous, ethanol, methanol ethyl acetate, chloroform, extract, four animals in each group having length 4-5cm. Albendazole was used as standard, where distilled water as control. In the first set of experiment, seven groups of four earthworms were released in to 25ml of solutions of aqueous and ethanol extracts of bark of *Ricinus communis* Linn. (50, 75 and 100mg/ml each) in distilled water. They were observed for their spontaneous motility and evoked responses. Time required for paralysis and death of animals were noted for each sample. Time for paralysis was noted when no movement of any sort could be observed except when the worms were shaken vigorously. The death of the worm was ascertained by transferring it into a beaker containing hot water (50 °C), which stimulated and induced movements if the worm was alive. The death and / or total paralysis time was recorded at room temperature.

**RESULT AND DISCUSSION:** The result of Anthelmintic activity is shown in **Table 1**. In the present study ethanol, aqueous, methanol, ethyl acetate chloroform the extracts showed remarkable Anthelmintic activity against intestinal parasitism. The all extract shows paralysis as well as death of worms in less time than ethanol, methanol, and aqueous extract especially at higher concentration of 100 mg/ml.

**TABLE 1: RESULTS OF ANTHELMINTIC ACTIVITY OF *RICINUS COMMUNIS* L. EXTRACTS**

Extract	Concentration (mg/ml)	Time taken for paralysis (min)	Time taken for death (min)
Ethanol	50	20.50±1.25	70.01±0.50
	75	17.20±0.54	32.50±1.25
	100	9.50±0.69	30.50±0.44
Aqueous	50	18.56±0.58	81.50±1.55
	75	17.22±0.25	43.45±0.76
	100	9.05±0.62	31.07±0.56
Ethyl acetate	50	40.30±1.22	98.24±0.77
	75	35.41±0.12	60.01±0.41
	100	22.05±0.56	40.99±0.52
Chloroform	50	35.03±0.67	97.11±0.76
	75	28.44±1.20	70.05±0.44

	150	20.22±0.70	45.90±0.72
Methanol	50	26.03±0.60	85.60±0.46
	75	19.47±1.12	60.05±0.76
	100	11.75±0.70	35.50±0.44
Control	-	-	-
Albendazol	20	8.30±0.60	40.21±1.04

**CONCLUSION:** It can be concluded that the leaves of *Ricinus communis* possess Anthelmintic activity against the worms and hence further investigation of active principles might help in finding of new compounds, which will be effective against parasitic infections.

**ACKNOWLEDGEMENT:** The author is thankful to all the management of Yashoda Technical Campus, Faculty of Pharmacy and Satara for providing experiment facilities.

**CONFLICTS OF INTEREST:** Nil.

#### REFERENCES:

- Gaikwad SA, Kale AA, Jadhav BG, Deshpande NR and Salvekar JP: Scholars Research Library, J. Nat. Prod. Plant Resour 2009; 1(2): 62-66.
- Vidyadhar S, Saidulu M, Gopal TK, Chamundeeswari D, Umamaheswara R and David B: International Journal of Applied Biology and Pharmaceutical Technology 2010; 1(3): 1119-1125.
- Aleemuddin MA, Karthikeyan M and Priya PK: Scholars Research Library J. Nat. Prod. Plant Resour 2012; 2(2): 267-271.
- Surana AR; Aher AN, Pal SC and Deore UV: International Journal of Pharmacy and Life Sciences 2011; 2(6): 813-814.
- Van RE, Hartqers FC and Yazdanbakshi M: Immunology, 2007; 212(6): 475-90.
- Tuse TA, Bidkar AA, Bhale SA and Patanka RD: International Journal of Pharmacological Research 2011; 1(1): 10-13.
- Wealth of India. Rawmaterials. Published by National Institute of science Communication Resource, New Delhi, 1972; 9: 26-47.
- Bentley R and Trimen H: A Textbook of Medicinal Plants, 2nd Edition, Asiatic Publishing House, New Delhi 2007; 237.
- Mary VK and Yasmin SS: Plant Science Feed 2011; 1(9): 167-173.
- Singh RK, Gupta MK, Singh AK and Kumar S: International Journal of Pharmaceutical Sciences and Research 2010; 1(6): 89-94.
- Rana M, Kumar H and Parashar B: Journal of Chemical and Pharmaceutical Research 2013; 5(6): 4042.

#### How to cite this article:

Mahadev ND and Vitthal BP: Evalaution of anthelmintic activity of *Ricinus communis* linn. leaves by using different types of solvent. Int J Pharmacognosy 2017; 4(12): 419-21;.doi link: [http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.4\(12\).419-21](http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.4(12).419-21).

This Journal licensed under a Creative Commons Attribution-Non-commercial-Share Alike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)