



Received on 16 October, 2017; received in revised form, 13 November, 2017; accepted, 18 November, 2017; published 01 February, 2018

## THE ROLE OF PLANT EXTRACTS IN THE TREATMENT OF LEUKEMIA TYPES

Seyyed Hossein Hassanpour\*<sup>1</sup> and Seyyedeh Zeinab Karami <sup>2</sup>

Young Researchers and Elite Club <sup>1</sup>, Yasooj Branch, Islamic Azad University, Yasooj, Iran.  
Department of Biology <sup>2</sup>, Faculty of Basic Sciences, Yasouj University, Yasouj, Iran.

### Keywords:

Cancer, Leukemia,  
Herbs, Apoptosis, Free radical

### Correspondence to Author: Seyyed Hossein Hassanpour

Young Researchers and Elite  
Club, Yasooj Branch, Islamic  
Azad University, Yasooj, Iran.


**Email:** Dr.hossein1366@yahoo.com

**ABSTRACT:** Cancer is one of the most common reasons mortality in worldwide. Leukemia as one type of cancer is a serious threat for human. Although there are many synthetic drug to treat leukemia but the use of them is limited due to side effect, therefore there is a dire need to find a promising solution. Herbs and their compound can be promising candidates for control leukemia. In this study, we review new studies entitled role of extracts of herbs to inhibit leukemia since 2014 to now. Our review study showed that herbs have anticancer activities through cell proliferation inhibition. This feature is due to induction of apoptosis and increase of free radical formation in cancer cell. Finally, we suggest that need to special attention to herbs in order to treat leukemia.

**INTRODUCTION:** Cancer is considered as main reason of mortality in developed and developing countries due to changing in lifestyle such as lack of exercise and physical activity, smoking and consuming of fast food <sup>1</sup>. Leukemia as one type of cancer is occurred due to uncontrolled growth of abnormal white blood cells and inability of blood-forming cells to differentiate into functional white blood cells so that these cancer cell transport to other organ such as lymph nodes, spleen and central nervous system by bloodstream <sup>2, 3</sup>. It can be divided two types: acute or chronic with fast or slow growing, respectively <sup>4</sup>. Although, the Wnt signaling is considered as a common pathway to induce other hematological malignancies and solids tumors but it is one of the pathways involved to leukemia development. In addition, Notch and SHH pathways have pivotal role to develop leukemia <sup>4</sup>.

In 2012, it has been diagnosed about 13,780 new patients with leukemia, interestingly, the mortality related to leukemia was 10,200 in United States <sup>5</sup>. The National Cancer Institute recently reported that it has been diagnosed 52380 leukemia patients so that 3% of all new cancer cases belonged to leukemia <sup>6</sup>. According to finding obtained from several European CML registries, the annual incidence of chronic myeloid leukemia is 0.7-1.3/100,000 <sup>7</sup>. Oxidative stress followed by imbalance in reactive oxygen species (ROS) formation and inactivity of their scavenging by antioxidant defense systems has prominent role in cancer pathophysiology <sup>8</sup>.

Increase of free radical leads to damage on macromolecule and ultimately increase of malondialdehyde, dityrosine formation, aggregation of protein and DNA fragmentation <sup>9, 10</sup>. Interestingly, it has been observed that ROS induce uncontrolled proliferation due to genetic instability during acute myeloid leukemia <sup>11 - 13</sup>. In conjunction with role of oxidative stress in relapse of acute myeloid leukemia, Zhou *et al.*, 2010 conducted a study on 102 patients with acute myeloid leukemia. The results revealed that levels of advanced

	<b>QUICK RESPONSE CODE</b>
	<b>DOI:</b> 10.13040/IJPSR.0975-8232.IJP.5(2).74-81
<b>Article can be accessed online on:</b> www.ijpjournal.com	
<b>DOI link:</b> <a href="http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5(2).74-81">http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5(2).74-81</a>	

oxidation protein products, malondialdehyde and 8-hydroxydeoxyguanosine and adenosine deaminase and xanthine oxidase activities as well as human thioredoxin (TRX) and indoleamine 2, 3-dioxygenase mRNA levels were significantly high in relapse status. In contrast, there was not showed reasonable results for parameters such as activities of glutathione peroxidase, monoamine oxidase and superoxide dismutase as well as level of total antioxidant capacity (T-AOC) <sup>14</sup>.

Evaluation of oxidative stress condition in 80 children with acute lymphoblastic leukemia was confirmed higher concentrations of plasmatic thiobarbituric acid reactive substances (TBARS) and serum protein carbonylation in these patients, while they had lower levels of parameters such as activities of whole blood catalase (CAT) and superoxide dismutase (SOD) and concentration of serum Vitamin E. Given that this study, there is continues oxidative stress in acute lymphoblastic leukemia <sup>13</sup>. In fact, potential ability of cancer cells to produce free radicals particularly  $O_2^{\cdot-}$  is a promising idea to treat cancer because their susceptibility to injury induced by ROS-generating agents is higher than normal cell. Thus, this strategy is considered as therapeutic selectivity <sup>15</sup>. Today, Use of traditional medicine to design new drugs is a promising idea to treat diseases <sup>16</sup>. Although, there are many anticancer drugs but their use is limited due to toxicity, side effect and non-selective targets <sup>17</sup>. Meanwhile, herbs have pivotal role to promote health society caused by high demand by people in worldwide <sup>17</sup>. In association with leukemia, it have been reported that herbs and their bioactive compound were served for treatment of types of leukemia such as CML, AML, lymphoid leukemia and Hodgkin's and non-Hodgkin's lymphomas <sup>18-20</sup>.

Also, natural products and plant extracts have been investigated for their ability to protect against radiation-induced toxicity <sup>21</sup>. Based on believe of scientists, many compound identified from plants are considered as anticancer drug such as quinine, salicylates ergotamine and digitalis. Furthermore, plants are good source from bioactive substance with anti-tumor activity that need to evaluate their ability in this field <sup>4</sup>. Here, we reviewed the role of extracts prepared from plants in treatment of type of leukemia.

**Review Method:** The aim of this study was to review new studies in association with effect of extracts prepared from plants to treat leukemia. We collected related studies by searching keywords such as “extract, herb, and leukemia”, “extract and herb and apoptosis and leukemia treatment”, etc from databases web of science, PubMed and Scopus since 2014 to now. Then, papers were read and their findings written.

**RESULTS:** Given that oil extract obtained from *Argania spinosa* has anti-proliferative effect against T-cell acute lymphoblastic leukemia human after its treatment into cell lines including JURKAT, MOLT3 and DND41, therefore, it can be a new useful treatment for acute lymphoblastic leukemia. In fact, this study was showed that this extract leads to growth inhibition of mentioned cell lines and obvious reduction of mRNA level and activity of ERK1/2 and Notch1 intracellular domain, as proliferation-related proteins <sup>22</sup>.

Probably, *Albizia zygia* be a potent anticancer herb because incubation of its aqueous and hydro-ethanolic extracts inhibits cell growth in Jurkat cells. Interestingly, its hydroethanolic extract results in prominent cell morphological changes and increase of DNA fragmentation due to induction of apoptosis in Jurkat cells <sup>23</sup>. It has been suggested that *Moringa oleifera* Lam is candidate plant to treat leukemia due to inhibition of cell proliferation subsequently its incubation into K562 cell line. They were showed that crude ethanolic extract prepared from *Moringa oleifera* Lam decreases WT1 protein level and has anti-proliferation property <sup>24</sup>. In a study, it has been found that besides inhibition of cell growth in jurkat cell line, it had good stability.

Therefore, it could be consider as promising therapy against leukemia cancer cells <sup>25</sup>. Souid *et al.*, 2016 showed that incubation of K562 cells by dehydrated aqueous extract obtained from *Allium roseum* results in inhibition of cell viability (by BCR-ABL kinase dephosphorylation and interference in ERK1/2, Akt, and STAT5 pathways), FOXO3 transcription factor activation (through Akt kinase inactivation). It also could improve expression pattern of FOXO3-regulated proapoptotic effectors, Bim and Bax, and cell cycle inhibitor p27 and abrogate vascular endothelial growth factor

secretion. Indeed, given that these finding should be special attention to this herb due to its anticancer property<sup>26</sup>.

A research group was incubated HL-60 (human promyelocytic leukemia) cell line with aqueous extracts of *Morinda lucida* and *Taraxacum officinale*. The results showed that both extract are cytotoxic against HL-60 due to proliferation inhibition through reduction of cell viability and apoptosis induction through increase of DNA fragmentation and changing of cell morphology<sup>27</sup>. Examination on antileukemic effects related to dimethyl sulfoxide extract prepared from *Withania somnifera* was revealed its cytotoxic and genotoxic activity against human T-lymphoblastoid cell line. Indeed, it has been demonstrated that this extract increases intracellular  $Ca^{2+}$  accumulation and reactive oxygen species formation as well as induces apoptosis so that leads to immunogenic cell death (ICD)<sup>28</sup>.

Study of effect of different extracts obtained from *Urtica dioica* on acute myelogenous leukemia was showed that chloroform has strongest effect in reduction of cell viability according to MTT assay on KG-1 cell line. In addition, both chloroform and ethyl acetate induced apoptotic pathway based on Flow cytometric analysis. Generally, this study revealed that *Urtica dioica* is a therapeutic potential source to overcome leukemia<sup>29</sup>. Tawil *et al.*, 2015 reported that incubation of *Daucus carota* oil extract into several acute myeloid leukemia (AML) cell lines results in apoptosis induction. Interestingly, cytotoxicity related to oil extract was diminished after MAPK pathway inhibition, therefore anticancer effect of *Daucus carota* is involved to MAPK pathway<sup>30</sup>.

Study conducted by Fan *et al.*, 2015 confirmed anticancer activity ethanol extract of *Meconopsis horridula* Hook so that it could induces cytotoxicity in L1210 cell line. This effect was through obvious alteration in cell morphology, increase of DNA fragmentation, apoptosis induction and arresting of G2/M related to cell cycle. Given that this extract had a potential effect to produce reactive oxygen species, thus induction of oxidative stress has central role in association with antitumor property of *Meconopsis horridula* Hook<sup>31</sup>. Given that deficiency of apoptosis is main

reason resistance to chemotherapy during B cell chronic lymphocytic leukemia (B CLL), Alhosin *et al.*, 2015 conducted a study about apoptotic effect of anthocyanin-rich dietary (*Vaccinium myrtillus*) bilberry extract on B CLL cells obtained from thirty patients. The results indicated that this extract leads to caspase 3 activation and down-regulation of Bcl-2 and UHRF1 (a rapid dephosphorylation of Akt and Bad).

In addition, it had prominent effect in increase of reactive oxygen species level. Moreover, when PEG-catalase was incubated to B CLL cells, the apoptosis induced by extract and its related signaling diminished that confirm the effect of extract on apoptosis induction in B CLL cells<sup>32</sup>. *Lepidium sativum* is one of the herbs that considered as a antitumor plant because its hydro-alcoholic extract incubation to K562 cells leads to obvious cytotoxic effect based on evaluation of cell viability by MTT test<sup>33</sup>. For determination of mechanisms related to anti-cancer activity of *Punica granatum* (pomegranate) against K562 cell line (chronic myeloid leukemia (CML) cells) was done a study by Asmaa *et al.*, 2015.

According to findings related to this study, this extract leads to either cell cycle arresting or apoptosis induction through inhibition of G2/M phase, increase of p21 and p53 levels and up-regulation of caspases and cytochrome c, respectively<sup>34</sup>. Given that chemotherapy and radiation lead to damage to normal cell during treatment of acute myeloid leukemia, thus they consider as non-selective therapies to overcome leukemia. In addition, there is dire requirement to find new anticancer drug for example *Myrothamnus flabellifolius* could be a promising candidate for this purpose. Indeed, it has been reported that methanol extract of *Myrothamnus flabellifolius* has cytotoxic effect in HL-60 cell line and reduces cell viability by induction of caspase-7 cleavage<sup>35</sup>.

*Basella Alba* considers as a natural edible source with antitumor activity so that its methanol extract results in reduction of cell viability and changing of cell morphology. In addition, MTT assay confirmed its cytotoxic effect against U937 cell line and it also could induce apoptosis in this cell line<sup>3</sup>. It has well been demonstrated that treatment of HL-60 cell line by ethanol extract prepared from orange

(*Citrus aurantium*) leads to cell viability inhibition. Based on WST assay, this extract had cytotoxic activity and based on agarose gel electrophoresis, it was confirmed DNA fragmentation followed by treatment. This study indicated that orange has potential ability to reduce cancer cell <sup>36</sup>.

In 2014, Assadollahi *et al.*, suggested that aqueous extract related to *Cinnamon zeylanicum* has prominent effect in reduction of human myelocytic leukemia by induction of apoptosis. In other word, they found that incubation of THP-1 cell line with mentioned extract increases Apoptotic cells after determination of Hoechst33342 staining. Moreover, flow cytometry and MTT assay confirmed cell cycle arresting and inhibition of cell proliferation. Finally, this study introduced an anticancer potential source that need to perform more investigations in further studies <sup>37</sup>.

Investigation of anticancer activity related to *Zanthoxylum heitzii* was confirmed that it has anti-proliferative effect in HL-60 cells treated by ethanol extract of *Zanthoxylum heitzii*. This study also revealed that mechanisms related to antitumor activity of this extract are obvious increase of reactive oxygen species (ROS), disturbance of mitochondrial membrane potential (MMP), DNA fragmentation and cell cycle arresting in G1/G0 phase <sup>38</sup>. Jantova *et al.*, 2014 reported that ethanol extract of *Salvia officinalis* has cytotoxic and anti-proliferative properties after its treatment in leukemia L1210 cell line. Interestingly, it induced apoptosis through pathway related to mitochondrial /caspase <sup>39</sup>.

Samet *et al.*, 2014 found that ethanol extract of olive (*Olea europaea*) has potential effect in inhibition of cell growth of K562 cell line. The evaluation of related mechanism was revealed that this extract leads to cells proliferation inhibition, cell cycle arresting (at G0/G1, and then at G2/M phase) and induction of apoptosis. Besides, microarray analysis confirmed high expression of genes related to K562 cells differentiation to monocyte/

macrophage lineage (IFI16, EGR1, NFYA, FOXP1, CXCL2, CXCL3, and CXCL8) <sup>40</sup>. In an experiment, it has been indicated that reduction of living cells number, induction of apoptosis occur in Jurkat Cell line incubated by ethanol extract of *Convolvulus arvensis*.

Indeed, this study confirms that *Convolvulus arvensis* has potential anti-cancer activity against leukemia cells <sup>41</sup>. Evaluation of role of n-hexane extract of *Cichorium intybus* in treatment of leukemia confirmed that its treatment is affective in reduction of cell viability and induction of apoptosis in Jurkat cells as a cell line related to human leukemia <sup>42</sup>. Moreover, it has been showed that reduction of mitochondrial outer membrane permeability and nuclear translocation of apoptosis-inducing factor, induction of LC3-I cleavage result from treatment with leaf extract obtained from *Azadirachta indica*.

In addition, according to biochemical tests this extract diminished BCl-2 and p53 proteins. Given that anti-tumor activity of this plant, it can be a promising treatment for patients with CLL (B cell chronic lymphocytic leukemia) <sup>43</sup>. In order to find a useful treatment against chronic myeloid leukemia was evaluated the effect of hydro-alcoholic extract prepared from *Mentha pulegium* on K562 cell line. At the end of study, it was found promising cytotoxic effect by this extract so that it significantly reduced cell viability <sup>44</sup>. Comparison of effect of two herbs (*Thymus vulgaris* Linn. and *Origanum syriacum* Linn.) against THP-1 leukemia cell line was revealed that both them have similar ability to reduce cell viability. In association with their effects on peripheral blood mononuclear cell (PBMC) was demonstrated that *Origanum syriacum* Linn. has more potent affect against PBMC but *Thymus vulgaris* L. leads to selective moderation. Based on findings of this study, given that *Thymus vulgaris* Linn. has selective therapeutic activity therefore its anticancer property is more than *Origanum syriacum* Linn. <sup>45</sup>.

**TABLE 1: SUMMARIZE OF TREATMENT OF LEUKEMIA BY EXTRACTS OF HERBS**

Plant	Extract	Animal model/cell line	Result(s)	References
<i>Argania spinosa</i>	Oil	Three T-ALL cell lines (JURKAT, MOLT3 and DND41)	To have anti-proliferative effects due to inactivation and expression reduction of ERK1/2 and Notch1	22
<i>Albizia zygia</i>	Aqueous and	JURKAT cell line	To have cytotoxic effects in both	23



	hydroethanolic		extract, apoptosis induction by hydroethanolic extract	
<i>Moringa oleifera</i>	Ethanolic	K562 cell line	Reduction of WT1 protein	24
<i>Solanum aethiopicum</i>	µEm Labrasol-crude extrac	JURKAT cell line	To have anti-proliferative effect and good stability	25
<i>Allium roseum</i>	Dehydrated aqueous	K562 cell line	Inhibition of cell viability and VEGF secretion, arresting of cell cycle in molecular level	26
<i>Morinda lucida</i>	Aqueous	HL-60 cell line	Reduction of proliferation, induction of apoptosis	27
<i>Taraxacum officinale</i>				
<i>Withania somnifera</i>	Dimethyl sulfoxide	Human T-lymphoblastoid cell line	Increase of intracellular Ca <sup>2+</sup> accumulation and ROS level	28
<i>Urtica dioica</i>	Aqueous , hydro-alcoholic, chloroform, ethyl acetate	KG-1 cell line	Chloroform extract; most effective to reduce cell viability, induction of apoptosis by chloroform and ethyl acetate	29
<i>Daucus carota</i>	Oil	AML cell lines (HL60, U937, ML1, ML2, Mono-Mac-1, Mono-Mac-6, KG-1, MV-4-11, TF1-vRaf, TF1-vSrc and TF1-HaRas )	To have apoptotic and cytotoxic effects probably by MAPK pathway	30
<i>Meconopsis horridula</i> Hook	Ethanol	L1210 cell line	Induction of apoptosis and inhibition of cell cycle by increase of oxidative stress	31
<i>Vaccinium myrtillus</i>	Anthocyanin-rich dietary extract	B CLL cell obtained from patients	Induction of apoptosis and oxidative stress	32
<i>Lepidium Sativum</i>	Hydro-alcoholic	K562 cell line	To have cytotoxic effect	33
<i>Punica granatum</i>	Ethanol	K562 cell line	Inhibition of proliferation and induction of apoptosis	34
<i>Myrothamnus flabellifolius</i>	Methanol	HL-60 cell line	To have apoptotic effect by induction of caspase-7 cleavage	35
<i>Basella alba</i>	Methanol	U937 cell line	Cell viability reduction, cell morphology changing, to have cytotoxic and apoptotic effects	3
<i>Citrus aurantium</i>	Ethanol	HL-60 cell line	Cell viability inhibition, to have cytotoxic activity DNA fragmentation increase	36
<i>Cinnamon zeylanicum</i>	Aqueous	THP-1	induction of apoptosis, anti-proliferative effect, inhibition of cell cycle	37
<i>Zanthoxylum heitzii</i>	Methanol	HL-60 cell line	ROS levels increase, MMP disturbance, DNA fragmentation and G1/G0 phase arresting	38
<i>Salvia officinalis</i>	Ethanol	L1210 cell line	Cytotoxic activity and reduction of cell proliferation by induction of apoptosis	39
<i>Olea europaea</i>		K562 Cells	Cell cycle arresting, apoptosis induction and K562 cell line differentiation to monocyte/macrophage lineage	40
<i>Convolvulus arvensis</i>	Ethanol	JURKAT cell line	Cell viability reduction, apoptosis induction	41
<i>Cichorium intybus</i>	n-hexane	JURKAT cell line	Reduction of living cell number and apoptotic activity	42
<i>Azadirachta indica</i>	Extract related to its leaf	PBMC obtained from patients with CLL	Reduction of mitochondrial outer membrane permeability, factors related to apoptosis as well as Bcl-2 and p53 proteins	43
<i>Mentha pulegium</i>	Hydro-alcoholic	K562 cell line	Reduction of cell viability	44

<i>Thymus vulgaris</i> L. <i>Origanum syriacum</i> L.	Ethanol	leukemia THP-1 cell line	To have cytotoxic effect in both extract, <i>Thymus vulgaris</i> L is safer than <i>Origanum syriacum</i> L due to selective therapeutic activity	45
--	---------	--------------------------	---	----

T-ALL: T-cell acute lymphoblastic leukemia human; VEGF: vascular endothelial growth factor; ROS: reactive oxygen species; AML: acute myeloid leukemia; B CLL: B cell chronic lymphocytic leukemia; MMP: mitochondrial membrane potential; PBMC: peripheral blood mononuclear cell




FIG. 1: SCHEMATIC PLAN OF EFFECTS RELATED TO HERBS EXTRACT IN REDUCTION OF CELL VIABILITY

**CONCLUSION:** Here, we reviewed the role of plants in treatment of leukemia based on their evaluations on different cell lines, animal models and cancer cells obtained from patients. In these studies, extracts prepared from plants had anticancer activity and leads to reduction of cell viability (**Fig. 1**). Induction of apoptosis was main mechanism to reduce cell viability through increase of caspases activities, increase of intracellular Ca<sup>2+</sup>, disturbance of mitochondrial membrane potential (MMP), reduction of mitochondrial outer membrane permeability, cytochrome c releasing and increase of BCl-2 protein level. In most studies, induction of apoptosis was confirmed by cell morphology changing and DNA fragmentation. Meanwhile, It should be noted this point that main factor to induce apoptosis was imbalance oxidant-antioxidant by increase of reactive oxygen species (ROS) levels and free radical formation. Cancer cells have potential effect for growth therefore; strategies related to cell growth inhibition can be helpful to treat leukemia.

Here, we found treatment with extracts inhibit cell proliferation by mechanism involved to cell cycle such as arresting of G1/G0 and G2/M phases. In addition, the some extract involved in MAPK pathway and moderation of p53 protein level.

Interestingly, according to finding obtained from microarray analysis, during anticancer activity related to olive occurred high expression of genes related to K562 cells differentiation to monocyte/macrophage lineage (IFI16, EGR1, NFYA, FOXP1, CXCL2, CXCL3, and CXCL8) that can be consider as an alternative strategy rather than other extracts. However, here it was confirmed that herbs have anticancer property but a treatment when is promising that performs as selective. Although these extracts had potential effects to reduce cancer cells but it was not examined selective treatment ability except for few cases. Finally, we suggest that should evaluate further studies to understand their ability as a selective treatment.

**ACKNOWLEDGEMENT:** Me acknowledgement and gratefulness at the beginning and at last is to god who gave us the gift of the mind. The author thanks Young Researchers and Elite Club, Yasooj Branch, Islamic Azad University due to cooperation in this study.

**CONFLICT OF INTEREST:** The author declares that there is no conflict of interest regarding this study.

## REFERENCES:

1. Jemal A, Bray F, Center MM, Ferlay J, Ward E and Forman D: Global cancer statistics. CA: a cancer journal for clinicians. 2011; 61(2): 69-90.
2. Petrie K, Zelent A and Waxman S: Differentiation therapy of acute myeloid leukemia: past, present and future. Current Opinion in Hematology. 2009; 16(2): 84-91.
3. Pal K: Cell growth inhibition and apoptosis by extract of *Basella alba* plant on U937 cells. 2015.
4. Kanga PT and Kamdje AHN: Signaling pathways in leukemia: Any role for medicinal plants in leukemia therapy. Journal of Diseases and Medicinal Plants. 2015; 1(5): 76-9.
5. Siegel R, Naishadham D and Jemal A: Cancer statistics, 2012. CA: A Cancer Journal for Clinicians. 2012; 62(1): 10-29.
6. Spore B. National Cancer Institute. 2014.
7. Höglund M, Sandin F and Simonsson B: Epidemiology of chronic myeloid leukemia. Chronic myeloid leukemia: Springer 2016; 99-110.
8. Valko M, Rhodes C, Moncol J, Izakovic M and Mazur M: Free radicals, metals and antioxidants in oxidative stress-induced cancer. Chemico-Biological Interactions. 2006; 160(1): 1-40.
9. Shanmugam N, Figarola JL, Li Y, Swiderski PM, Rahbar S and Natarajan R: Proinflammatory effects of advanced lipoxidation end products in monocytes. Diabetes. 2008; 57(4): 879-88.
10. Selmeçi L, Seres L, Soós P, Székely M and Acsady G: Kinetic assay for the determination of the oxidative stress biomarker, advanced oxidation protein products (AOPP) in the human blood plasma. Acta Physiologica Hungarica. 2008; 95(2): 209-18.
11. Zhou F, Zhang W, Wei Y, Zhou D, Su Z, Meng X, *et al.*, The changes of oxidative stress and human 8-hydroxyguanine glycosylase1 gene expression in depressive patients with acute leukemia. Leukemia research. 2007; 31(3): 387-93.
12. Austin C: Does oxidative damage contribute to the generation of leukemia? Leukemia Research. 2009; 33(10): 1297.
13. Battisti V, Maders LD, Bagatini MD, Santos KF, Spanevello RM, Maldonado PA, *et al.*, Measurement of oxidative stress and antioxidant status in acute lymphoblastic leukemia patients. Clinical Biochemistry. 2008; 41(7): 511-8.
14. Zhou F-L, Zhang W-G, Wei Y-C, Meng S, Bai G-G, Wang B-Y, *et al.*, Involvement of oxidative stress in the relapse of acute myeloid leukemia. Journal of Biological Chemistry. 2010; 285(20): 15010-5.
15. Hileman EO, Liu J, Albitar M, Keating MJ and Huang P: Intrinsic oxidative stress in cancer cells: a biochemical basis for therapeutic selectivity. Cancer Chemotherapy and Pharmacology. 2004; 53(3): 209-19.
16. Mule SN, Patil SB, Naikwade NS and Magdum CS: Evaluation of antinociceptive and anti-inflammatory activity of stems of *Gynandropsis pentaphylla* Linn. International Journal of Green Pharmacy (IJGP). 2008; 2(2).
17. Houghton P, Fang R, Techatanawat I, Steventon G, Hylands PJ and Lee C: The sulphorhodamine (SRB) assay and other approaches to testing plant extracts and derived compounds for activities related to reputed anticancer activity. Methods. 2007; 42(4): 377-87.
18. Ramkumar K, Manjula C, Elango B, Krishnamurthi K, Saravana Devi S and Rajaguru P: *In-vitro* cytotoxicity of *Gymnema montanum* in human leukaemia HL-60 cells; induction of apoptosis by mitochondrial membrane potential collapse. Cell proliferation. 2013; 46(3): 263-71.
19. Omoregie SN, Omoruyi FO, Wright VF, Jones L and Zimba PV: Antiproliferative activities of lesser galangal (*Alpinia officinarum* Hance Jam), turmeric (*Curcuma longa* L.), and ginger (*Zingiber officinale* Rosc.) against acute monocytic leukemia. Journal of Medicinal Food. 2013; 16(7): 647-55.
20. Asmaa M, Al-Jamal H, Ang CY, Asan JM, Seeni A and Johan MF: Apoptosis induction in MV4-11 and K562 human leukemic cells by *Pereskia sacharosa* (Cactaceae) leaf crude extract. Asian Pac J Cancer Prev. 2014; 15: 475-81.
21. Kma L: Plant extracts and plant-derived compounds: promising players in a countermeasure strategy against radiological exposure. Asian Pacific journal of cancer prevention: APJCP. 2013; 15(6): 2405-25.
22. Aribi B, Zerizer S, Kabouche Z, Screpanti I and Palermo R: Effect of *Argania spinosa* oil extract on proliferation and Notch1 and ERK1/2 signaling of T-cell acute lymphoblastic leukemia cell lines. Food and Agricultural Immunology. 2016; 27(3): 350-7.
23. Appiah-opong R, Asante IK, Safo DO, Tuffour I, Ofori-attah E, Uto T, *et al.*, Cytotoxic effects of *Albizia zygia* (dc) JF Macbr, a ghanaiian medicinal plant, against human t-lymphoblast-like leukemia, prostate and breast cancer cell lines. International Journal of Pharmacy and Pharmaceutical Sciences. 2016; 8(5): 392-6.
24. Semsri S, Anuchapreeda S and Janwitayanuchit W: Inhibitory effects of crude ethanolic leave extract from *Moringa oleifera* Lam. on Wilms' tumor 1 protein expression in K562 leukemic cell line. Bulletin of Chiang Mai Associated Medical Sciences. 2016; 49(1): 53.
25. Christian KK, Anatolievna SE, Akhanovna PM-BJ and Yves-Alain PB: Extract-based microemulsion formulation with *Solanum aethiopicum* peduncle: characterization by DLS, relative humidity and F0 anti-proliferative effect on leukemia cancer cells. 2016.
26. Souid S, Najjaa H, Riahi-Chebbi I, Haoues M, Neffati M, Arnault I, *et al.*, *Allium roseum* Linn. Extract exerts potent suppressive activities on chronic myeloid leukemia K562 cell viability through the inhibition of BCR-ABL, PI3K/Akt, and ERK1/2 Pathways and the Abrogation of VEGF Secretion. Nutrition and Cancer. 2016; 1-14.
27. Appiah-Opong R, Tuffour I, Annor GK, Doris A, Blankson-Darku PC, Kissi-Twum AA, *et al.*, anti-proliferative, antioxidant activities and apoptosis induction by *Morinda lucida* and *Taraxacum officinale* in human HL-60 leukemia cells. Journal of Global Biosciences. 2016; 5(7): 4281-91.
28. Turrini E, Calcabrini C, Sestili P, Catanzaro E, de Gianni E, Diaz AR, *et al.*, *Withania somnifera* induces cytotoxic and cytostatic effects on human t leukemia cells. Toxins. 2016; 8(5): 147.
29. Keshavarz S, Ardekani MRS, Safavi M, Chahardouli B and Nadali F: *In-vitro* cytotoxic effect of *Urtica dioica* extracts on acute myelogenous leukemia cell line (KG-1). Archives of Medical Laboratory Sciences. 2016; 2(1).
30. Tawil M, Bekdash A, Mroueh M, Daher CF and Abi-Habib RJ: Wild carrot oil extract is selectively cytotoxic to human acute myeloid leukemia cells. Asian Pac J Cancer Prev. 2015; 16: 761-7.

31. Fan J, Wang Y, Wang X, Wang P, Tang W, Yuan W, *et al.*, The antitumor activity of *Meconopsis horridula* Hook, a traditional Tibetan medical plant, in murine leukemia L1210 cells. *Cellular Physiology and Biochemistry*. 2015; 37(3):1055-65.
32. Alhosin M, León-González AJ, Dandache I, Lelay A, Rashid SK, Kevers C, *et al.*, Bilberry extract (Antho 50) selectively induces redox-sensitive caspase 3-related apoptosis in chronic lymphocytic leukemia cells by targeting the BCL-2/Bad pathway. *Scientific reports*. 2015; 5.
33. Aslani E, Naghsh N and Ranjbar M: Cytotoxic effects of hydro-alcoholic extracts of cress (*Lepidium sativum*)-made from different stages of the plant-on k562 leukemia cell line. *Hormozgan Med J*. 2015; 18: 411-9.
34. Asmaa MJS, Ali A-JH, Farid JM and Azman S: Growth inhibitory effects of crude pomegranate peel extract on chronic myeloid leukemia, K562 cells. *International Journal of Applied and Basic Medical Research*. 2015; 5(2): 100.
35. Badiab A, Dhillon J, Nabbe F and Peethambaran B: Selective cytotoxicity and mechanism of *Myrothamnus flabellifolius*, an edible medicinal plant, on acute myeloid leukemia cells. *Cancer Research*. 2015; 75(15-S): 3802.
36. Diab KA, Shafik RE and Yasuda S: *In-vitro* antioxidant and antiproliferative activities of novel orange peel extract and its fractions on leukemia HL-60 Cells. *Asian Pacific Journal of Cancer Prevention*. 2015; 16(16): 7053-60.
37. Assadollahi V, Gholami M and Zendedel A: *C. zeylanicum* aqueous extract induced apoptosis in the human myelocytic leukemia cell line (THP-1). *Bratislavské lekárske listy*. 2014; 116(2): 132-5.
38. Pieme CA, Santosh GK, Tekwu EM, Askun T, Aydeniz H, Ngogang JY, *et al.*, Fruits and barks extracts of *Zanthoxylum heitzii* a spice from cameroon induce mitochondrial dependent apoptosis and G0/G1 phase arrest in human leukemia HL-60 cells. *Biological research*. 2014; 47(1): 1.
39. Jantová S, Hudec R, Sekretár S, Kučerák J and Melušová M: *Salvia officinalis* Linn. extract and its new food antioxidant formulations induce apoptosis through mitochondrial/caspase pathway in leukemia L1210 cells. *Interdisciplinary toxicology*. 2014; 7(3): 146-53.
40. Samet I, Han J, Jlaiel L, Sayadi S, Isoda H: Olive (*Olea europaea*) leaf extract induces apoptosis and monocyte/macrophage differentiation in human chronic myelogenous leukemia k562 cells: Insight into the underlying mechanism. *Oxidative medicine and cellular longevity*. 2014; 2014.
41. Saleem M, Qadir MI, Ahmad B, Saleem U, Naseer F, Schini-Kerth V, *et al.*, Cytotoxic effect of ethanol extract of *Convolvulus arvensis* Linn. (Convolvulaceae) on lymphoblastic leukemia Jurkat cells. *Tropical Journal of Pharmaceutical Research*. 2014; 13(5): 705-9.
42. Saleem M, Abbas K, Naseer F, Ahmad M, Javed F, Hussain K, *et al.*, Anticancer activity of n-hexane extract of *Cichorium intybus* on lymphoblastic leukemia cells (jurkat cells). *African Journal of Plant Science*. 2014; 8(6): 315-9.
43. Chitta KS, Khan ANH, Ersing N, Swaika A, Masood A, Paulus A, *et al.*, Neem leaf extract induces cell death by apoptosis and autophagy in B-chronic lymphocytic leukemia cells. *Leukemia and lymphoma*. 2014; 55(3): 652-61.
44. Aslani E, Naghsh N and Ranjbar M: Cytotoxic effect of hydroalcoholic extract of *Mentha Pulegium* before flowering on k562 leukemia cell line. 2014.
45. Ayesh BM, Abed AA and Doa'a MF: *In-vitro* inhibition of human leukemia THP-1 cells by *Origanum syriacum* L. and *Thymus vulgaris* Linn. extracts. *BMC research notes*. 2014; 7(1): 612.

**How to cite this article:**

Hassanpour SH and Karami SZ: The role of plant extracts in the treatment of leukemia types. *Int J Pharmacognosy* 2018; 5(2):74-81. doi link: [http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5\(2\).74-81](http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5(2).74-81).

This Journal licensed under a Creative Commons Attribution-Non-commercial-Share Alike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)