



Received on 24 November, 2017; received in revised form, 05 January, 2018; accepted, 13 February, 2018; published 01 April, 2018

## PHARMACOGNOSTIC STUDY OF *ERANTHEMUM NIGRUM* ROOT

D. S. N. B. K Prasanth\* and A. Lakshmana Rao

Department of Pharmacognosy, V.V. Institute of Pharmaceutical Sciences, Gudlavalleru - 521356, Andhra Pradesh, India.

### Keywords:

Pharmacognostic, *Eranthemum nigrum*, Lignified xylem vessels, Phytochemical and Physicochemical analysis

### Correspondence to Author:

**D. S. N. B. K Prasanth**

Department of Pharmacognosy, V. V. Institute of Pharmaceutical Sciences, Gudlavalleru - 521356, Andhra Pradesh, India.


**Email:** dsnbkprasanth@gmail.com

**ABSTRACT: Objective:** To analyze the pharmacognostic characteristics and physiochemical parameters of the root of *Eranthemum nigrum* (*E. nigrum*). **Methods:** Microscopic characters and powder analysis had been carried out with the help of a microscope. The physiochemical properties such as loss on drying, total ash value, acid insoluble ash value, water soluble ash value, extractive values and fluorescence of *E. nigrum* had been performed. **Results:** The color, shape, size, odor, and surface characteristics were reported from the root and powdered root material of *E. nigrum*. Light microscope images of cross section and powdered root revealed the presence of lignified xylem fibers, xylem vessels, cork cells and parenchyma cells. Phytochemical testing confirmed the presence of steroids, alkaloids, tannins, saponins, carbohydrates, glycosides, amino acids and proteins. Physicochemical parameters such as moisture content, ash value, extractive value and fluorescent behavior of root powder have also been established. **Conclusion:** The morphological, microscopical and physicochemical parameter results provided in this paper may be utilized as a basis for the preparation of a monograph on *E. nigrum* root.

**INTRODUCTION:** Medicinal plants are usually playing a significant part in traditional medicines intended for therapy of various health issues. However a crucial hurdle, which has impeded the promotion in the usage of alternative medications in the developed countries, is lack of evidence of documentation and absence of stringent quality control measures. Additionally, there is a dependence on the data of all study meted out on traditional medicines by way of documentation. Keeping this issue, it is now quite necessary to generate assurance about the standardization of the plant as well as its parts to be used like a medication.

During the process of standardization, we are able to take advantage of various techniques and methodology to achieve our goal in a phase wise approach e.g. pharmacognostic and phytochemical studies. These techniques and methods are helpful in recognition and standardization of the plant material. Appropriate characterization and quality assurance of starting material is a crucial step to ensure reproducible quality of herbal medicine to assist people in order to justify its safety and effectiveness. Because of this reason, we have executed pharmacognostic studies of *Eranthemum nigrum* belongs to family Acanthaceae<sup>1</sup>. This sort of research is not going to help in authentication but additionally ensures reproducibility of herbal goods in promoting<sup>2</sup>.

In the present study, we have been focusing our exploration on one of the commonly available plant in India i.e., *Eranthemum nigrum*, belongs to family Acanthaceae. The family Acanthaceae consists of almost 4000 species of exotic plants.



Various species of genus *Eranthemum* being utilized traditionally for extensive kinds of ethno medicinal purposes. The genus *Eranthemum*, with around 138 species, some of the important species include *E. austrosinensis*, *E. burmanicum*, *E. capense*, *E. ciliatum*, *E. erythrochilum*, *E. griffithii*, *E. macrophyllum*, *E. macrostachyus*, *E. obovatum*, *E. pulchellum*, *E. purpurascens*, *E. roseum*, *E. strictum*, *E. tapingense*, *E. tubiflorum* and *E. watti*. The *Eranthemum nigrum* (Acanthaceae) is native to Pacific Islands. The shrub attains height a height of 1.5 - 1.8 m. The upper surface of leaves is blackish purple and the lower surface purplish with dark veins. The flowers are in terminal erect spikes, white and spotted rose at the base<sup>3</sup>. Plants are adapted to partial shade. The leaves are elliptical, glossy or dull with smooth margins and acute tips<sup>4,5</sup>. All parts of this plant are widely used as a folklore medicine for the treatment of various ailments by the Indian traditional healer. Ethnomedicinally, the genus *Eranthemum* has been documented various pharmacological activities including antipyretic<sup>6</sup>, antidiabetic<sup>7</sup>, antiulcer<sup>8</sup>, antimicrobial<sup>9</sup>, larvicidal, ovicidal and pupicidal against *Anopheles stephensi*<sup>10</sup>, gastroprotective<sup>11</sup> and anti-inflammatory<sup>12</sup>.

A literature study and screening of scientific data says a lot of native medicines have already been investigated as regards their botany and chemistry is concerned, however a systematic standardization including pharmacognostical and physicochemical study is still lacking. The present investigation of *Eranthemum nigrum*. (Acanthaceae) is therefore taken up to establish certain botanical and chemical standards which would help in crude drug identification as well as in checking adulteration, if any. Further, the study will greatly help in quality assurance of finished products of herbal drugs<sup>13,14</sup>.

## MATERIALS AND METHODS:

**Plant Collection and Authentication:** The plant was obtained from V.V. Institute of Pharmaceutical Sciences, Gudlavalleru, Krishna District of Andhra Pradesh, India during the month of September 2017 and authenticated by Dr. K. Madhava Chetty, Taxonomist at Sri Venkateswara University, Tirupati, India. The plant was deposited at the herbarium for future reference. One portion of the root is preserved in formalin: acetic acid: alcohol mixture for histological studies and the remaining

portion was shade dried, powdered and sieved through 20 mesh and kept in an air tight container for future use.

**Chemicals:** All analytical grade chemicals were utilized in this study were procured from E. Merck, Germany absolute alcohol, phloroglucinol, acetic acid, chloral hydrate, H<sub>2</sub>SO<sub>4</sub>, NaOH, HNO<sub>3</sub>, FeCl<sub>3</sub>, distilled water, Conc. HCl and chloroform.

## Pharmacognostic Evaluation:

**Morphological Evaluation:** Organoleptic evaluation of *E. nigrum* root has been carried out in accordance the colour, size, odour, shape and taste as per WHO quality control methods of herbal medicine<sup>15</sup>.

## Microscopic Evaluation:

**Preparation of Sections:** Microscopic studies had been done by preparing thin hand section of the root with the help of sharp cutting edge of the blade, then cleared with chloral hydrate solution, stained with phloroglucinol-hydrochloric acid (1:1) and mounted in glycerin.

**Powdered Microscopy:** The powder microscopy was carried out in accordance with the procedure described in Khandelwal<sup>16</sup>.

**Quantitative Analysis:** The quantitative examinations including stomatal number, stomatal index, vein islet number and vein termination number were studied using standard method<sup>2</sup>.

**Preparation of Extracts and Preliminary Phytochemical Analysis:** The powdered material had been extracted with various solvents according to its polarity *i.e.*, chloroform, methanol and water. 5 g root powder was extracted with 20 ml of the respective solvent by maceration at room temperature for 24 hours. Then, filtered through Whatmann filter paper and collect the filtrate, concentrated with rota-evaporator. Then, the extracts had been subjected to preliminary phytochemical screening according to standard methods<sup>16,17</sup>.

**Physicochemical Analysis:** Physicochemical parameters such as ash value, moisture content and extractive values were determined according to the procedures mentioned in WHO quality control methods for herbal materials<sup>18</sup>.

**Fluorescence Analysis:** Various reagents were utilized to check the fluorescence activity. In this, 0.1 g of root powder was blended with 1.5 ml of respective reagent **Table 4**. The mixture was placed on slide for a minute and observed under visible light, short ultra-violet light (254 nm) and long ultraviolet light (365 nm) (19).

## RESULTS:

**Morphological Characteristics:** The morphological characteristics of *E. nigrum* root were described in **Fig. 1** and **Table 1**.



**FIG. 1: MORPHOLOGICAL FEATURES OF ROOT OF ERANTHEMUM NIGRUM**


**TABLE 1: MORPHOLOGICAL CHARACTERISTICS OF ROOT OF ERANTHEMUM NIGRUM**

Characters	Observation
Colour	Buff
Odour	Characteristic
Taste	Characteristic
Texture	Smooth
Thickness	0.2-4 cm

## Anatomical Description:


**Root:** The transverse section of the root of *E. nigrum* showed the presence of cortex (Phellem) shows 3- 4 polygonal thick walled parenchymatous cells filled with brown content. Cork cambium (Phellogen) was made up of 3 - 5 layered narrow, tangentially elongated parenchymatous cells. Secondary cortex (Phelloderm) is 4 - 6 layered rows of tangentially elongated thin walled cells. The endodermis showed the presence of phloem and xylem. The phloem is present in between the medullary rays. The medullary rays are parenchymatous and are uniseriate to triseriate,

majorly biseriate. Radially arranged vascular bundles were present in which, Phloem is well developed and shows the presence of phloem fibers, which are non-lignified. It also showed the presence of phloem parenchyma. The xylem region was similar to phloem region and was also surrounded by uniseriate to triseriate medullary rays. Xylem tissue consists of spiral xylem vessels, xylem fibers and xylem parenchyma **Fig. 2**.



**FIG. 2: TRANSVERSE SECTION OF ROOT OF ERANTHEMUM NIGRUM**


Ph: Phellem; Pg: Phellogen; Pd: Phelloderm Cx: Cortex; MR: Medullary rays; XV: Xylem Vessels; XP: Xylem parenchyma; PI: Pith.




**FIG. 3: T.S OF ERANTHEMUM NIGRUM ROOT SHOWED MEDULLARY RAYS AND XYLEM VESSELS**

MR: Medullary Rays; XV: Xylem Vessels







**FIG. 4: DETAILED T.S OF ROOT SHOWED CENTRAL REGION OF ERANTHEMUM NIGRUM**  
MR: Medullary rays; XV: Xylem vessels; PI: Pith



**FIG. 5: T.S SECTION OF ERANTHEMUM NIGRUM ROOT SHOWED CORK REGION**  
C: Cork; Pg: Phellogen; Pd: Phelloderm

**Powder Microscopy:** The crude powder of root was buff in color with characteristic odor and taste. Microscopic study of the powder showed revealed

different characters such as diacytic stomata, covering trichomes, xylem vessels and parenchyma cells **Fig. 8.**



**FIG. 8: POWDER MICROSCOPY OF ROOT OF ERANTHEMUM NIGRUM**  
(a) Parenchyma cells (b) lignified xylem vessels (c) Cork cells (d) Medullary rays with xylem vessels

**Preliminary Phytochemical Analysis:** The results of qualitative phytochemical analysis of crude powder of *E. nigrum* root are shown in **Table 2**.

**TABLE 2: PRELIMINARY PHYTOCHEMICAL ANALYSIS OF ERANTHEMUM NIGRUM ROOT**

Phytoconstituents	Method	Aqueous Extract	Methanolic Extract	Chloroform Extract	Pet. ether Extract
Flavonoids	Shinoda Test	-	-	-	-
	Zn. Hydrochloride test	+	+	-	-
	Lead acetate Test	+	+	-	-
Volatile oil	Stain test	-	-	-	-
	Wagner Test	+	+	+	-
Alkaloids	Hager's Test	+	+	+	-
	FeCl <sub>3</sub> Test	-	-	-	-
	Potassium dichromate test	-	-	-	-
Saponins	Foaming Test	+	+	-	-
Steroids and Triterpenoids	Salkowski test	+	+	-	+
Carbohydrates	Molish test	+	+	-	-
Acid compounds	Litmus test	-	-	-	-
Glycoside	Keller-Killani Test	+	+	-	-
Amino acids	Ninhydrin test	+	+	-	-
Proteins	Biuret	+	+	-	-

“+” -Present; “-” -Absent

**Physicochemical Parameters:** The results attained from various determinations of physicochemical analysis are produced in **Table 3**.

**TABLE 3: PHYSICOCHEMICAL PARAMETERS OF ROOT POWDER OF ERANTHEMUM NIGRUM**

Parameters	Values % w/w
Moisture content (Loss on drying)	8.25 ± 0.63
Total ash	6.56 ± 0.23
Acid insoluble ash	4.12 ± 1.22
Petroleum ether soluble extractive value	0.63 ± 0.05
Chloroform soluble extractive value	2.12 ± 0.06
Ethyl acetate soluble extractive value	6.24 ± 0.05
Alcohol soluble extractive value	8.66 ± 0.25
Water soluble extractive value	10.12 ± 0.22

**Fluorescence Analysis:** Fluorescence analysis of root powder was performed out after treating with different solvents. Fluorescence was observed at 254 and 365 nm comparing its change of color in the visible light. The observations are presented in **Table 4** shows the variation in color.

**TABLE 4: FLUORESCENCE ANALYSIS OF ERANTHEMUM NIGRUM ROOT POWDER**

Solvent used	Visible light	UV light	
		At short (254 nm)	At long (365 nm)
Distilled water	Buff	Black	Black
Methanol	Brown	black	Greenish black
1N HCl	Black	Black	Black
50% HNO <sub>3</sub>	Black	Black	Blue
FeCl <sub>3</sub>	Brownish yellow	Dark blue	Black
CHCl <sub>3</sub>	Pale green	Black	Black
Picric acid	Brownish yellow	Dark blue	Black
Ethyl acetate	Black	Black	Greenish black

**DISCUSSION:** Indian systems of medicine utilize majority of the crude drugs which are of plant origin. It is important that standards need to be set down to control and check the identity of the plant and confirm its quality before use. Hence a detailed pharmacognostic assessment is extremely an important prerequisite. In accordance with World health organization (WHO) the organoleptic and histological description of a medicinal plant could be the first step towards establishing its identity and

purity and should be performed before to any tests tend to be undertaken<sup>20</sup>. *E. nigrum*, extensively utilized in conventional medicines has tremendous therapeutical potential due to its various biological activities. The prominent diagnostic characteristics of root were parenchyma cells, lignified xylem vessels, cork cells, medullary rays with xylem vessels.

These characters can be utilized for standardization of drugs as well as useful for preparation of plant monograph and also reduces the possibilities of adulteration, when the drug is available in the powdered form studies of physicochemical parameters can serve as an important source to judge the purity and quality of crude drugs.

Ash values are utilized to establish the quality and purity of the crude drug. It implies the existence of various impurities like carbonate, oxalate and silicate. The water soluble ash is water soluble part of total ash, employed to calculate the amount of inorganic substances found in the drugs. The acid insoluble ash comprises mostly silica and indicates contamination with earthy matter. The moisture content of drugs might be at minimum level in order to suppress the growth of microorganisms like bacteria, yeast or fungi during storage. The extractive values are helpful to judge the chemical constituents present in the crude drug and also assist in the evaluation of particular constituents soluble in a specific solvent. Total ash and acid insoluble ash are essential indices to illustrate the quality and purity of the herbal medicine.

Total ash consists of physiological ash, which is derived from plant tissue itself, and non physiological ash that is usually from atmosphere contaminations includes sand and soil. Total ash content alone is not adequate to indicate the quality of herbal medicine, because the plant materials usually contain a significant level of physiological ash, calcium oxalate in particular.

Therefore, the acid insoluble ash content is another index to indicate the quality of herbal medicine<sup>20, 21, 22</sup>. The phytochemical analysis of extracts viz., petroleum ether, chloroform, methanol and water were analyzed and it indicates the presence of steroids, alkaloids, tannins, saponins, carbohydrates, glycosides, amino acids and proteins.

**CONCLUSION:** Standardization of herbal drugs is very much crucial because they are produced from heterogeneous sources which could result in variations. These kinds of variations can cause spurious results in various pharmacological and phytochemical studies. *Eranthemum nigrum* root are recognized for many therapeutical properties, therefore, the current study might be beneficial to supplement the information in respective to its identification, authentication, and standardization; no such information is available for the same till date.

**ACKNOWLEDGEMENT:** Nil

**CONFLICTS OF INTEREST:** Nil

#### REFERENCES:

1. Akbar S, Hanif U, Ali J and Ishtiaq S: Pharmacognostic studies of stem, roots and leaves of *Malva parviflora* L. Asian Pac J Trop Biomed. 2014; 4(5): 410-5.
2. Amponsah IK, Mensah AY, Otoo A, Mensah MLK and Jonathan J: Pharmacognostic standardisation of *Hillieria latifolia* (Lam.) H. Walt. (Phytolaccaceae). Asian Pac J Trop Biomed. 2014; 4(12): 941-6.
3. Randhawa GS and Mukhopadhyay A: Floriculture in India. New Delhi: Allied Publ.; 2004.
4. Thekkayam SG and Peter KV: Ornamental plants. New Delhi: New India Pub. Agency; 2009.
5. Ona NJ: Foliage: astonishing color and texture beyond flowers North Adams, MA: Storey Pub.; 2007. Available from: <http://public.ebib.com/choice/publicfullrecord.aspx?p=4540058>.
6. Samvatsar S and Diwanji V: Plants used for the treatment of different types of fevers by Bhils and its subtribes in India 2004.
7. Dhruv D, Anil T, Sanjay S and Sapna D: Antidiabetic and antioxidant properties of roots of *Eranthemum roseum* vahl r. br. Inventi Rapid: Ethnopharmacology 2010.
8. Kamble S, Patil S, Sawant P, Sawant S, Pawar S and Singh E: Studies on plants used in traditional medicine by Bhilla tribe of Maharashtra 2010.
9. Jain A, Surana S, Gokhale S, Tatiya A and Bothara R: Antimicrobial Properties of *Eranthemum roseum* (Vahl) R. Br. Iran J Pharm Res. 2010; 6(2): 131-3.
10. Milne-Redhead E: *Eranthemum* of the "Flora of Tropical Africa". Bulletin of Miscellaneous Information (Royal Botanic Gardens, Kew) 1936; (4): 255-74.
11. Patil P and Surana S: Gastroprotective effect of *Eranthemum roseum* R. Br. Linn Root extracts in albino rats. Int J Pharmacol Biol Sci. 2009; 3(1): 81-93.
12. Tatiya A, Desai D, Surana S and Patil P: Anti inflammatory and nitric oxide scavenging activity of roots of *Eranthemum roseum*. 2007; 44(11): 815-8.
13. Jhade D, Ahirwar D, Jain R, Sharma N and Gupta S: Pharmacognostic standardization, physico- and phytochemical evaluation of *Amaranthus spinosus* linn. Root. J Young Pharm. 2011; 3(3): 221-5.
14. Ghorpade P, Siddiqui A, Patil MJ and Rub RA: Pharmacognostic and phytochemical evaluation of *Celosia argentea*. Phcog J. 2012; 4(33): 7-15.

15. WHO: Quality control methods for herbal materials Geneva: World Health Organization 2011.
16. Khandelwal KR: Practical pharmacognosy: Techniques and experiments. Maharashtra: Niral Prakashan 2008.
17. Harborne A: Phytochemical methods a guide to modern techniques of plant analysis: springer science and business media 1998.
18. WHO: Quality control methods for medicinal plant materials. Geneva: World Health Organization 1998.
19. Galani VJ and Patel BG: Psychotropic activity of *Argyreia speciosa* roots in experimental animals. Ayu. 2011; 32(3): 380-4.
20. Dave R, Nagani K and Chanda S: Pharmacognostic Studies and Physicochemical Properties of the *Polyalthia longifolia* var. pendula Leaf. Phcog J. 2010; 2(13): 572-6.
21. Vaghasiya Y, Nair R and Chanda S: Antibacterial and preliminary phytochemical and physicochemical analysis of *Eucalyptus citriodora* Hk leaf. Nat Prod Res. 2008; 22(9): 754-62.
22. Prasanth D, Rao AS and Yejella RP: Assessment of Pharmacognostic, Phytochemical and Physicochemical Standards of *Aralia racemosa* (L.) root. Ind J Pharm Edu Res. 2016; 50(3): S225-S30.

**How to cite this article:**

Prasanth DSNBK and Rao AL: Pharmacognostic study of *Eranthemum nigrum* root. Int J Pharmacognosy 2018; 5(4): 242-48;.doi link: [http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5\(4\).242-48](http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5(4).242-48).

This Journal licensed under a Creative Commons Attribution-Non-commercial-Share Alike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)