



Received on 07 December, 2017; received in revised form, 05 January, 2018; accepted, 13 February, 2018; published 01 April, 2018

EVALUATION OF LAXATIVE ACTIVITY OF VARIOUS FRUIT EXTRACTS OF *ANETHUM GRAVEOLENS* L.

Suchismita Pani, Sangram K. Panda* and B. Bhagaban Patra

Jeypore College of Pharmacy, Rondapalli, Jeypore, Koraput - 764002, Odisha, India.

Keywords:

Anethum graveolens,
Laxative activity,
Agar-agar, Albino rats

Correspondence to Author: Sangram K. Panda

Jeypore College of Pharmacy,
Rondapalli, Jeypore, Koraput -
764002, Odisha, India.

Email: sangrampanda04@gmail.com

ABSTRACT: *Anethum graveolens* an important medicinal plant and one of the most widely cultivated species of the family (Umbelliferae). It is highly valued from time immemorial because of its vast medicinal properties. The present work deals with the investigation of laxative activity of various extracts of *Anethum graveolens* fruits. The laxative effect was checked in using Wister strain Albino rats. All the crude extracts such as ethanol, ethyl acetate, *n*-butanol and petroleum ether at a dose of 200 and 400 mg/kg were tested for laxative activity, where as agar-agar 300 mg/kg, p.o. was used as standard drugs. Among all the extract, the petroleum ether extract was found to possess the most effective laxative activity. Therefore *Anethum graveolens* fruits can be a substitute of synthetic laxative drugs having adverse effects.

INTRODUCTION: *Anethum graveolens* L. commonly known as (dill) fruit has been used in Ayurvedic medicines since ancient times. The tribal areas of Koraput (District) of Eastern Orissa, due to its unique varieties geographical and climatic factors has had a rich variety of medicinal plant and extensively used traditionally by the tribal people. *Anethum graveolens*, belong to the family Umbelliferae, is indigenous to Southern Europe. It is an annual herb growing in the Mediterranean region, Central and Southern Asia. Now it is cultivated widely throughout the world¹⁻⁴. It is used traditionally as a popular aromatic herb and spice that has a very long history of use going back to more than 5,000 years. It was used as a remedy for indigestion and flatulence and as milk secretion stimulant.


Moreover, it is used as an anti-convulsion, anti-emetic, anti-cramp (in children), as a wound healer and to increase the appetite and strengthen the stomach⁵⁻⁷. The major compounds of dill seed essential oils were carvone and limonene whereas dill apiole, trans - dihydrocarvone and α -phellandrene were present in appreciable amounts. Trace amounts of α -thujene, α - pinene, sabinene), myrcene, p-cymene γ -terpinene, dill ether, iso-dihydro carveol, trans-carveol and anethole were also present in essential oil⁸.

Some pharmacological effects have been reported, such as antimicrobial, antihyperlipidemic and anti-hypercholesterolemic activities⁹.

MATERIAL AND METHODS:

Collection of Plant Material and Identification:

The fruits of *Anethum graveolens* were collected from the local area of Jeypore and all so identified, confirmed and authenticated by the Biju Patnaik Medicinal Plants Garden and Research Centre, Jeypore, Koraput, Odisha. (Letter No. MJ/SS/P-607/17, dated (22.10.2017)). After authentication the fruits were collected in bulk and dried in

	<p>QUICK RESPONSE CODE</p> <p>DOI:</p> <p>10.13040/IJPSR.0975-8232.IJP.5(4).256-58</p>
	<p>Article can be accessed online on:</p> <p>www.ijpjournal.com</p>
<p>DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5(4).256-58</p>	

sunlight and made into coarse powder by grinding in mechanical grinder. Then stored in a closed air tight container for further use.

Preparation of Extracts: The coarse powder was taken in Soxhlet apparatus and extracted successively with ethanol, ethyl acetate, *n*-butanol and petroleum ether as solvent. A total amount of 500 g coarse powder was extracted with 1000 ml of each solvent. For each solvent, 10 cycles were run to obtain thick slurry. Each slurry was then concentrated under reduced pressure to obtain the crude extract. All crude extracts were kept in closed air tight containers under cool and dark place for further study^{10,11}.

Animal: Healthy adult Wister strain of Albino rats were procured and housed in animal house maintained under standard hygienic conditions and the studies conducted were approved by the institutional animal ethical committee (1906/PO/Re/S/16/CPCSEA), Jeypore College of Pharmacy, Jeypore, Odisha according to prescribed guide-lines of the committee for the purpose of control and supervision of experiments on animals (CPCSEA), Government of India.

Acute Toxicity Studies: The acute toxicity was performed according to OECD 423, 2001. The selected Albino rats were used to determine the dose. The animals were divided into five groups, six animals in each group. The animals were fasted overnight prior to the acute experimental procedure. Distilled water was used as vehicle to suspend the different fruit extracts of *Anethum graveolens* and administered orally as following doses of 100, 300, 600, 1000 and 2000 mg/kg body weight. Immediately after dosing, the animals were observed continuously for first four hours for behavioral changes and for mortality at the end of 24 h and daily for 14 days respectively¹².

Laxative Activity: The test was performed according to Capasso *et al.*, on rats of either sex weighing 200 - 220 g were kept in individual cages during one week. Any rat producing wet feces was rejected. The rats were fasted for 12 h before the experiment, but with water provided *ad libitum*. The animals were divided into eight groups of six in each. The animal groups were administered orally either with vehicle (1% Tween 80 solution in

normal saline, 25 ml/kg, reference standard drug, agar-agar (300 mg/ kg, p.o.) in saline or doses of extracts 200 and 400 mg/kg. Immediately after dosing, the animals were separately placed in cages suitable for collection of faeces. (Each cage is with a wire mesh at the bottom and a funnel to the urine; stainless-steel sieves are placed in the funnel to retain feces). After 8 h of drug administration, the faeces were collected and weighed. Thereafter, food and water were given to all rats and faecal outputs were again weighed after a period of 16 h^{13,14}.

TABLE 1: LAXATIVE ACTIVITY OF VARIOUS FRUITS EXTRACTS OF ANETHUM GRAVEOLENS

Treatment	Dose mg/kg	Fecal output (g)	
		8 h	8 - 16 h
Control	-	0.7 ± 0.1	1.1 ± 0.2
Agar-Agar (standard)	300	1.3 ± 0.1	2.3 ± 0.3
Ethanol extract	200	1.5 ± 0.1	2.5 ± 0.2
	400	1.3 ± 0.2	2.2 ± 0.2
Ethyl acetate extract	200	0.7 ± 0.2	1.6 ± 0.1
	400	0.6 ± 0.3	2.6 ± 0.2
<i>n</i> -butanol extract	200	0.7 ± 0.3	1.7 ± 0.3
	400	0.9 ± 0.1	2.7 ± 0.3
Pet. ether extract	200	0.8 ± 0.2	1.9 ± 0.2
	400	1.2 ± 0.1	2.0 ± 0.2




FIG. 1: LAXATIVE ACTIVITY OF ANETHUM GRAVEOLENS

1: Normal control, 2: Standard (Agar-agar), 3: Ethanol ext. 200 mg/kg, 4: Ethanol ext. 400 mg/kg, 5: Ethyl acetate ext. 200 mg/kg, 6: Ethyl acetate ext. 400 mg/kg, 7: *n*-butanol ext. 200 mg/kg, 8: *n*-butanol ext. 400 mg/kg, 9: Pet. ether ext. 200 mg/kg, 10: Pet. ether ext. 400 mg/kg

Statistical Analysis: The results are reported as mean ± SE. Statistical analysis was done using ANOVA (Tukey - multiple comparison Test). When probability (p) was less than 0.05 was considered as significant¹⁵.

RESULT AND DISCUSSION: In present study four extracts (ethanol, ethyl acetate, *n*-butanol and petroleum ether) of fruit part of *Anethum graveolens* were studied for laxative activity.

A preliminary acute toxicity study in mice showed that all the extracts were not toxic ($LD_{50} > 1000$ mg/kg). The pet-ether extract of fruit of *Anethum graveolens* shows significant laxative activity than other two extract in a dose of 200 mg/kg body weight as compared to standard drug agar-agar in a dose of (300 mg/kg) which were shown in **Table 1** and **Fig. 1**.

CONCLUSION: In present study four extracts ethanol, ethyl acetate, *n*-butanol and petroleum ether of fruits of *Anethum graveolens* were studied for laxative activity among all the extract pet-ether extract shows significant laxative activity than other three extract in a dose of 200 mg/kg body weight as compared to standard drug Agar-agar in a dose of 300 mg/kg. Therefore, the traditional use has been pharmacologically validated and the *Anethum graveolens* fruits can be use a substitute of synthetic laxative having adverse effect.

ACKNOWLEDGEMENT: Authors wish to thank to local people of south eastern Odisha for giving valuable information about the plant and all so wish to express their gratitude to Jeypore College of Pharmacy, Rondapalli, Jeypore, Koraput, Odisha.

CONFLICTS OF INTEREST: Nil

REFERENCES:

1. Wren RC: Potter's new cyclopedia of botanical drugs and preparations. Saffron Walden, CW Daniel 1988.

2. Leung AY and Foster S: Encyclopedia of common natural ingredients used in food, drugs and cosmetics. New York, John Wiley and Sons 1996.
3. Yazdanparast R and Bahramikia S: Evaluation of the effect of *Anethum graveolens* L. crude extracts on serum lipids and lipoproteins profiles in hypercholesterolaemic rats. DARU 2008; 16(2): 88-94.
4. African pharmacopoeia. Lagos, Organization of African Unity, Scientific Technical and Research Commission 1985; 1.
5. Zargari A: Medicinal Plants. 6th ed. Tehran University Press, Tehran 1996; 2: 531-528.
6. Ishikawa TM, Kudo M and Kitajima J: Water-soluble constituents of dill. Chem Pharm Bull 2002; 55: 501-507.
7. Kaur GJ and Arora DS: Bioactive potential of *Anethum graveolens*, *Foeniculum vulgare* and *Trachyspermum ammi* belonging to the family Umbelliferae - Current status. Journal of Medicinal Plants Research 2010; 4(2): 87-94.
8. Chahal KK, Monika, Kataria D and Singh R: Antifungal potential of dill seed essential oil and its constituents. Indian Journal of Ecology 2016; 43(S-2): 903-906.
9. Radulescu V, Popescu ML and Ilies DC: Chemical composition of the volatile oil from different plant parts of *Anethum graveolens* L. (Umbelliferae). Farmacia 2010; 58: 594-600.
10. Kokate CK, Purohit AP and Gokhale BS: Pharmacognosy, 47 editions. Pune: Nirali Prakashan 2008; 324-328.
11. Geissman A: Modern Methods of Plant Analysis, Berlin: Springer Verlag 1955; 3.
12. 423, OECD, Guideline for the Testing of Chemicals, Guidance document on acute toxic class method 2001.
13. Ogunti EO and Elujoba AA: Laxative activity of *Cassia alata*, Fitoterapia 1993; 9: 64-437.
14. Hughes S, Higgs NB and Turnberg LA: Loperamide has antisecretory activity in the human jejunum *in vivo*. Gut. 1984; 5: 25-931.
15. Bolton S: Analysis of variance. In: Pharmaceutical statistics-practical and clinical application. Marcel Dekker NY 1997; 2(1).

How to cite this article:

Pani S, Panda SK and Patra BB: Evaluation of laxative activity of various fruit extracts of *Anethum graveolens* L. Int J Pharmacognosy 2018; 5(4): 256-58. doi link: [http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5\(4\).256-58](http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5(4).256-58).

This Journal licensed under a Creative Commons Attribution-Non-commercial-Share Alike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)