



Received on 01 April 2017; received in revised form, 24 May 2017; accepted, 30 May 2017; published 01 August 2017

## NATURAL REMEDIES TARGET DIFFERENT THERAPEUTIC PATHWAYS IN ORAL MUCOSITIS INDUCED BY CANCER CHEMO OR RADIOTHERAPY

Sherin Zakaria

Department of Pharmacology and Toxicology, Faculty of Pharmacy, Damanhour University, Damanhour, Egypt.

### Keywords:

Oral mucositis, Herbal remedies, Antioxidant, Anti-inflammatory, Keratinocyte apoptosis (KGF)

### Correspondence to Author:

**Sherin Zakaria**

Department of Pharmacology and Toxicology, Faculty of Pharmacy, Damanhour University, Damanhour, Egypt.

**E-mail:** sherinzm\_2008@yahoo.com

**ABSTRACT:** Oral mucositis (OM) is an inflammatory condition affecting oral mucosa. The etiology of this type of inflammation is associated with exposure to radiotherapy or chemotherapeutic drugs. Herbal drugs have been used to induce healing of different types of gastrointestinal ulcers. The role of medicinal plant in cancer therapy-induced oral mucositis has been investigated extensively. Preclinical studies refer to healing activity of some medicinal plants such as Chamomile, royal jelly, *Calendula officinalis* and *salvadora perisca* in oral mucositis induced by either chemotherapy or radiotherapy. Some herbal drugs are proved to have potential efficacy in decreasing both incidence and severity of oral mucositis in clinical studies. The molecular targets modulated by herbal drugs are various including scavenging reactive oxygen species (ROS), inhibition of inflammatory cascade, prevention of keratinocyte apoptosis and induction of some growth factors. This review, illustrates the common herbal drugs that have been used for oral mucositis management focusing on different therapeutic pathways that are implicated in this pharmacologic activity.

**INTRODUCTION:** Oral mucositis is an inflammatory condition that affects the mucosa of the oral cavity. The etiology of this type of inflammation is associated with exposure to radiotherapy or chemotherapeutic drugs<sup>1, 2</sup>. Patients under chemotherapeutic protocols that induced bone marrow-suppression are at high risk of oral mucositis. Approximately 60-100% of those patients may encounter oral mucositis, patients who need radiotherapy directed at the oral, submandibular and pharyngeal area such as those who have squamous cell carcinoma in the head and neck region commonly have oral mucositis<sup>3</sup>.

Combination of radiotherapy and chemotherapy is associated with a risk of around 100%<sup>2</sup>. Oral mucositis is a painful condition that significantly affects patients' quality of life<sup>4</sup>. The severe case is associated with ulcerated mucosa and secondary infection which may lead to life-threatening sepsis.

Herbal drugs have been used to induce healing of different types of gastrointestinal ulcers such as gastric ulcer, aphthous stomatitis, ulcerative colitis and oral mucositis<sup>5-8</sup>. The role of medicinal plants in cancer therapy-induced oral mucositis has been investigated extensively. Preclinical studies refer to healing activity of some medicinal plants like Chamomile, royal jelly, *Calendula officinalis* and *Salvadora perisca* in oral mucositis<sup>9-12</sup>. Recently, some of medicinal plants products are formulated in various dosage forms. These preparations have been proved to be effective in the prevention of oral mucositis in patients under chemo or radiotherapy<sup>13-15</sup>.

	<p><b>QUICK RESPONSE CODE</b></p>
	<p><b>DOI:</b> 10.13040/IJPSR.0975-8232.IJP.4(8).250-56</p>
<p>The article can be accessed online on <a href="http://www.ijournal.com">www.ijournal.com</a></p>	
<p><b>DOI link:</b> <a href="http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.4(8).250-56">http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.4(8).250-56</a></p>	

This review illustrates the common herbal drugs that have been used for oral mucositis management focusing on different therapeutic pathways that are targeted by these medicinal plants.

**Pathophysiology of Oral Mucositis:** Oral mucositis induced by antineoplastic via direct epithelial cell injury starting with DNA strand breaks concurrently with production of reactive oxygen species (ROS) <sup>16</sup> especially in early stage of oral mucositis which results in consumption of large quantities of antioxidants molecules including glutathione <sup>17</sup>. Degraded DNA strand and ROS activate nuclear factor- $\kappa$ B (NF- $\kappa$ B) <sup>18</sup>. It an important factor that up-regulates expression of pro-inflammatory cytokines such as tumor necrosis factor  $\alpha$  (TNF-  $\alpha$ ), Interleukin 1 $\beta$  (IL-1 $\beta$ ), and Interleukin 6 (IL-6). Those cytokines amplify tissue damage <sup>19, 20</sup>. Tissue damage is deteriorated with activation of matrix metalloproteinase as a result of activated pro-inflammatory molecules <sup>21, 22</sup>.

All these processes led to the formation of ulcers in the mucosa. The healing phase of oral mucositis initiated by signaling pathways starting in an endothelial cell with a specific growth factor that controls the renewal of epithelial cell proliferation and differentiation. Keratinocyte growth factor (KGF) is one of the key molecules that regulate communication between endothelial and epithelial cells. It is associated with triggering epithelial cells growth and differentiation <sup>23</sup>.

### Evidence for Efficacy of Traditional Medicine in Oral Mucositis:

**Chamomile:** There are two types of chamomile, german chamomile (*Chamomilla recutita*) and roman chamomile (*Chamaemelum nobile*). The main active constituents of Chamomile flowers are terpenoids and flavonoids <sup>24</sup>.

Pavesi and his colleagues <sup>9</sup> reported that topical preparation of chamomile decreases the incidence and severity of oral mucositis induced by 5-fluorouracil in the hamster. It has been reported that Chamomile ointment decreases the severity of oral mucositis by decreasing pro-inflammatory cytokine <sup>25</sup>. These results were further evaluated in clinical studies. One of these studied use cammomile extract as an infusion. Cammomile infusion decreased the incidence and severity of

mucositis when used as adjuvant therapy with cryotherapy in a patient under 5-fluorouracil and leucovorin protocols. Moreover, the toxic profile of Cammomile infusion showed a wide margin of safety <sup>13</sup>.

Results of another randomized, controlled, phase II clinical trial pointed out that cammomile mouthwash (containing 1% *C. recutita* extract) can be associated with the reduction in incidence, intensity, and duration of mucositis in patients undergoing allogeneic hematopoietic stem cell transplantation HSCT <sup>26</sup>.

**Calendula officinalis:** *Calendula officinalis* is a member of the Asteraceae family. Extract of *C. officinalis* flowers contains active constituents that have a different pharmacological activity such as bactericidal, antiseptic, anti-inflammatory, and antioxidant <sup>11</sup>. Topical *Calendula officinalis* extract formulated as gel accelerated the healing of oral mucositis induced by 5-fluorouracil (5-FU) in hamsters <sup>11</sup>. Clinically, *C. officinalis* mouthwash ameliorated the severity of radiotherapy-induced oral mucositis in patients with head and neck cancer <sup>14</sup>.

**Salvadora persica:** Miswak (*Salvadora persica* L.) is the most widely used chewing stick for oral hygiene in middle-eastern and eastern African cultures, which is prepared from the roots or stems of *Salvadora persica* L. (*S. persica*) <sup>27</sup>. *S. persica* accelerates healing of ulceration induced by ethanol, indomethacin and cold restraint stress in rats <sup>28</sup>. Recently, *Salvadora persica* aqueous extract was found to decrease the incidence of oral mucositis induced by 5-fluorouracil in rats. The mechanism underlying this effect may be multifactorial including preservation of oral hygiene as well as increased level of KGF in mucosa tissues <sup>12</sup>.

Different herbal remedies that showed promising results in preventing or accelerate healing of oral mucositis are summarized in **Table 1**.

**Herbal Remedies Combination:** Traditional Japanese herbal medicine prescribes herbal remedies that contain different medicinal plants such as Daiokanzoto (TJ-84) and Hangeshashinto. The composition of Hangeshashinto is illustrated in **Table 2**.

The herbal combination Daiokanzoto has an anti-inflammatory and immunomodulatory effect suggesting possible benefits in treating oral mucositis. It has been reported that Daiokanzoto exerts some activity in decreasing apoptosis in gingival cells (SA3 cell line) exposed to 5-fluorouracil through inhibition of ROS release from mitochondria<sup>37</sup>. Similarly, Using Hangeshashinto

(TJ-14) enables patients under chemo-radiotherapy to complete their treatment course and preserve their nutritional status during treatment periods<sup>38</sup>. Results from the random placebo-controlled study revealed that Hangeshashinto is effective in treating mucositis in colorectal cancer patients under different chemotherapeutic protocols<sup>39</sup>.

**TABLE 1: EVIDENCE OF DIFFERENT HERBAL REMEDIES EFFICACY IN PREVENTING OR RETARDING ORAL MUCOSITIS INDUCED BY CHEMOTHERAPY**

Plant name	Oral Mucositis Model	Results	Proposed Mechanism
Korean red ginseng	Radiation-induced oral mucositis in a rat	Decrease the severity of oral mucositis in rats	Inhibition of keratinocyte apoptosis <sup>29</sup>
<i>Hippophae rhamnoides</i> extract	Oral mucositis induced in rats with methotrexate	Decrease incidence and severity of oral mucositis in rats	Antioxidant and anti-inflammatory <sup>30</sup>
Topical olive leaf extract	Oral mucositis induced by 5-fluorouracil in the golden hamster	Improve healing of oral mucositis in the golden hamster	Antioxidant <sup>8</sup>
<i>Zizyphus jujuba</i> extract	Oral mucositis induced by 5-fluorouracil in the golden hamster	Improve healing of oral mucositis in the golden hamster	Antioxidant and anti-inflammatory <sup>31</sup>
Hydroalcoholic extract of <i>Carum carvi</i> L.	Oral mucositis induced by 5-fluorouracil in the golden hamster	Improve healing of oral mucositis in the golden hamster	Antibacterial <sup>32</sup>
<i>Salvadora persica</i> extract	Oral mucositis induced by 5-fluorouracil in rats	Decrease incidence and severity of oral mucositis in rats	Oral hygiene Induction of KGF <sup>12</sup>
<i>Clandula officinalis</i>	Oral mucositis induced by 5-fluorouracil in hamster	Improve healing of oral mucositis in the golden hamster	Antioxidant <sup>11, 14</sup>
Royal jelly (3%, 10%, and 30%) ointments	Oral mucositis induced by 5-fluorouracil in hamster	Improve healing of oral mucositis in the golden hamster	Antioxidant <sup>10, 33</sup>
Honey	Clinical studies on radiochemotherapy-induced mucositis in head and neck cancer patients.	Significant reduction in grade 4 mucositis	Anti-inflammatory <sup>34, 35</sup>

**TABLE 2: COMPOSITION OF HANGESHASHINTO<sup>36</sup>**

Name of Herb	Dry Weight (g) per day
<i>Pinelliae ternatae</i> Rhizoma	5
<i>Scutellariae baicalensis</i> Radix	2.5
<i>Glycyrrhizae uralensis</i> Radix	2.5
<i>Zizyphi jujubae</i> Fructus	2.5
<i>Gin seng</i> Radix	2.5
<i>Coptidis</i> Rhizoma	1
<i>Zingiberis officinalis</i> Recens Rhizoma	2.5

### Molecular Targets Modulated by Traditional Medicine:

**Antioxidant:** Chemotherapy and radiotherapy are associated with the release of ROS which plays a pivotal role in initiating a cascade that ends in tissue damage. Moreover, ROS have direct toxicity to mucosal cells<sup>40 - 42</sup>. Hence, scavenging ROS could limit the progress of mucositis cascade and ameliorates tissue damage. In normal cells, ROS level is managed by the balance between ROS and

antioxidant enzyme, e.g., glutathione peroxidase, glutathione reductase, SOD and catalase. Scavenging ROS and activation of an antioxidant enzyme is considered as the main pharmacologic targets modulated by many herbal drugs. This may attribute to many antioxidant substances such as polyphenol and flavonoid. For example, *Calendula officinalis* decreases the intensity of radiotherapy-induced oral mucositis<sup>14</sup>. The effects of this medicinal plant are mostly related to its antioxidant compositions such as polyphenols, carotenoids and triterpenes. The main flavonoid in *C. officinalis* is quercetin this compound has potent antioxidant activities<sup>14, 43</sup>.

In the same manner, *Hippophae rhamnoides* exerts a prophylactic effect against methotrexate-induced oral mucositis<sup>30</sup>. The *H. rhamnoides* leaf extract has potent antioxidant activity due to the bioactive

phenolic constituents, such as quercetin-3-O-galactoside, quercetin-3-O-glucoside, kaempferol and isorhamnetin<sup>44</sup>. Similarly, the antioxidant polyphenol particularly oleuropein and hydroxytyrosol found in topical olive leaf extract is responsible for its healing activity of oral mucositis induced by 5-fluorouracil<sup>8,11</sup>.

*Zizyphus jujuba* is an herb that is widely distributed in Europe and South-eastern Asia. The main active constituents in this herb are cyclopeptide alkaloids, flavonoids, sterols, jujuboside A, jujuboside B, lauric acid, and triterpenoid and saponins<sup>45</sup>. Topical and systemic forms of *Z. jujuba* hydro-alcoholic extract are associated with the reduced intensity of oral mucositis of golden hamster undergoing 5-FU consumption mainly due to antioxidant effect as measured as the low level of MDA and increased activity of SOD on mucosa<sup>31</sup>.

**Anti-inflammatory:** As discussed in the pathophysiology section above, induction of pro-inflammatory cytokines such as (TNF- $\alpha$ ), Interleukin 1 $\beta$  (IL-1 $\beta$ ), and Interleukin 6 (IL-6) play an important role in the amplification of tissue injury through activation NF- $\kappa$ B and matrix metalloproteinase<sup>19, 20</sup>. Targeting these pro-inflammatory cytokines may be an important part of the pharmacological activity of some medicinal herbs such as chamomile and *Hippophae rhamnoides*. Chamomile, Royal jelly, and *Hippophae rhamnoides* reduce the tissue levels of IL-1 $\beta$  and TNF- $\alpha$ <sup>25, 30, 33</sup>. Other herbal drugs block inflammatory cascade and decrease the formation of prostaglandin. Honey has an anti-inflammatory effect; it inhibits prostaglandin level in both plasma and mucosa tissue<sup>35</sup>.

The Japanese herbal combination hangeshashinto enable patients to complete chemo-radiation therapeutic course partially due to its anti-inflammatory activity<sup>46</sup>. Kono and his colleagues explained that hangeshashinto reduce PGE2 production in human oral keratinocyte. They suggested that this anti-inflammatory action is due to the presence of an active ingredient that inhibits inflammation such as [6] - shogaol, [6] - gingerol, wogonin, baicalein, baicalin, and berberine<sup>47</sup>.

**Oral Hygiene:** Maintaining Oral hygiene is very important in patients receiving chemotherapy<sup>48</sup>. Those patients are of high risk of secondary

infection which may be life-threatening as those patients are already neutropenic due to chemotherapy<sup>49</sup>. *Salvadora persica* Linn. protect a variety of microorganism<sup>50-53</sup>. The reported antiseptic effect of *S. persica* is attributed to its phytoconstituents such as vitamin C, salvadorine, salvadouria, alkaloids, trimethylamine, cyanogenic glycosides, tannins, saponins, and salts mostly as chlorides<sup>54-57</sup>.

Moreover, it has been reported that *S. persica* aqueous extract contains potential antimicrobial anionic compound such as Cl, SO<sub>4</sub> and SCN<sup>27</sup>. The use of *S. persica* was associated decrease in incidence and severity of oral mucositis induced by 5- fluorouracil in rats. This may impart due to maintaining oral hygiene and induction of growth factor expression such as KGF<sup>12</sup>. Similarly, a topical form of *Carum carvi* Linn. (caraway) is associated with the reduced intensity of oral mucositis due to impart appropriate antibacterial activity of its terpinene contents<sup>32</sup>.

**Inhibition of Keratinocyte Apoptosis:** Radio therapy-induced apoptosis is one of the important molecular events in oral mucositis. Radiation induces activation of caspase<sup>3</sup>. The activated caspase 3 cleaves the RNA-binding protein HuR and subsequently promotes the expression of the pro-apoptotic factor Bax<sup>58</sup>. Inflammatory molecules that released during the progress of mucositis often share in apoptosis induction in mucositis<sup>59</sup>. Korean red ginseng inhibit caspase induced opotosis in the oral mucosa of irradiated rats<sup>29</sup>, *Artemisia asiatica* is an herbal drug that have an anti-apoptotic effect. *Artemisia asiatica* reduced the expression of cytochrome c, cleaved caspase-3 and nuclear factor-kappa B (NF- $\kappa$ B) induced by cisplatin. Moreover, it induces the expression of Antiapoptotic gene Bcl-2<sup>29</sup>.

**Analgesic Effect:** Some medical herbs have an analgesic effect. This effect is potentially important in oral ulcerative mucositis which is very painful. The processed ginger extract contains two analgesic compounds namely [6] - gingerol and [6] - shogaol. These constituents inhibit voltage-activated Na<sup>+</sup> currents. Moreover, they inhibit the stimulant-induced release of substance P and action potential generation in cultured rat sensory neurons<sup>60</sup>.

**Induction of Growth Factor:** Healing process is initiated by signaling pathways that target proliferation and differentiation of epithelial cells. Various types of growth factor are incorporated in different interaction to stimulate epithelial cell growth<sup>61</sup>. The most promising growth factor that regulates growth and proliferation of epithelial cell is the fibroblast growth factors (FGFs). Keratinocyte growth factor (KGF) is the most potent growth factors that induce epithelial growth and proliferation. Palifermin, the generic name of KGF, is the first compound approved by the FDA to reduce oral mucositis in patients receiving HSCT<sup>62</sup>. Limited studies investigated the role of herbal remedies in the induction of growth factor. *S. persica* enhances expression of KGF in mucosal tissues of 5- fluorouracil treated rats<sup>12</sup>. On the other hand, Watanabe and his colleagues<sup>33</sup> reported that royal jelly couldn't increase the release of KGF from HPdLFs cells.

**CONCLUSION:** Herbal remedies are effective as an alternative therapy for oral mucositis. Data collected from both animal and clinical studies suggest herbal remedies as adjuvant therapy for oral mucositis induced by chemo or radiotherapy. Herbal drugs modulate different therapeutic pathways such as scavenging ROS, inhibiting inflammatory processes, inhibiting keratinocyte apoptosis and maintaining oral hygiene in addition to the analgesic effect of some product. I think that multicenter randomized studies are needed to clarify the efficacy and safety of different herbal products as a therapeutic option for oral mucositis.

**ACKNOWLEDGEMENT:** Nil

**CONFLICT OF INTEREST:** Nil

## REFERENCES:

- Alvariño-Martín C and Sarrión-Pérez MG: Prevention and treatment of oral mucositis in patients receiving chemotherapy. *J Clin Exp Dent* 2014; 6: e74-80.
- Rohani B, Pourfar K, Pourshahidi H and Ebrahimi SH: Oral manifestation of hematologic malignancies. *Jundishapur Sci Med J* 2015; 14: 477-85.
- Khan M and Gupta N: Oral mucositis. *E J Dent* 2013; 3: 405-11.
- Kolokythas A: Long-term surgical complications in the oral cancer patient: A comprehensive review. Part II. *J Oral Maxillofac Res* 2010; 1(3): e2.
- Fakhraei N, Abdolghaffari AH, Delfan B, Abbasi A and Rahimi N: Protective effect of hydro alcoholic olive leaf extract on experimental model of colitis in rat: Involvement of nitregeric and opioidergic systems. *Phytother Res* 2014; 28(9): 1367-73.
- Moghadamnia AA, Motallebnejad M and Khanian M: The efficacy of the bioadhesive patches containing licorice extract in the management of recurrent aphthous stomatitis. *Phytother Res* 2009; 23(2): 246-50.
- Jabri MA, Aissani N, Tounsi H, Sakly M and Marzouki L: Protective effect of chamomile (*Matricaria recutita* L.) decoction extract against alcohol-induced injury in rat gastric mucosa. *Pathophysiology* 2016; 16: 30047-5.
- Showraki N, Mardani M, Emamghoreishi M, Andishe-Tadbir A and Aram A: Topical olive leaf extract improves healing of oral mucositis in golden hamsters. *J Dent (Shiraz)* 2016; 17(4): 334-342.
- Pavesi VC, Lopez TC, Martins MA, Sant' Ana Filho M and Bussadori SK: Healing action of topical chamomile on 5-fluorouracil induced oral mucositis in hamster. *Support Care Cancer* 2011; 19(5): 639-46.
- Suemaru K, Cui R, Li B, Watanabe S and Okihara K: Topical application of royal jelly has a healing effect for 5-fluorouracil-induced experimental oral mucositis in hamsters. *Methods Find Exp Clin Pharmacol* 2008; 30(2): 103-6.
- Tanideh N, Tavakoli P, Saghiri MA, Garcia-Godoy F and Amanat D: Healing acceleration in hamsters of oral mucositis induced by 5-fluorouracil with topical *Calendula officinalis*. *Oral Surg Oral Med Oral Pathol Oral Radiol* 2013; 115(3): 332-8.
- Zakaria S and Abdel-Raheem IT: *Salvadora persica* aqueous extract promotes healing of 5-fluorouracil induced oral mucositis in rats: a possible role of KGF. *Asian Journal of Phytomedicine and Clinical Research* 2016; 4(2): 69 - 79.
- Dos Reis PE, Ciol MA, de Melo NS, Figueiredo PT and Leite AF: Chamomile infusion cryotherapy to prevent oral mucositis induced by chemotherapy: a pilot study. *Support Care Cancer* 2016; 24(10): 4393-8.
- Babae N, Moslemi D, Khalilpour M, Vejdani F and Moghadamnia Y: Antioxidant capacity of *Calendula officinalis* flowers extract and prevention of radiation induced oropharyngeal mucositis in patients with head and neck cancers: a randomized controlled clinical study. *Daru* 2013; 21(1): 18.
- Yamauchi K, Kogashiwa Y, Moro Y and Kohno N: The effect of topical application of royal jelly on chemo radiotherapy-induced mucositis in head and neck cancer: a preliminary study. *Int J Otolaryngol* 2014; 974-967.
- Sonis ST: Oral mucositis in cancer therapy. *J Support Oncol* 2004; 2(6): 3-8.
- Yoshino F, Yoshida A, Nakajima A, Wada-Takahashi S and Takahashi S: Alteration of the redox state with reactive oxygen species for 5-Fluorouracil-induced oral mucositis in Hamsters. *PLoS One* 2013; 8(12): e82834
- Sonis S: The biologic role of nuclear factor-kB in disease and its potential involvement in mucosal injury associated with antineoplastic therapy. *Crit Rev Oral Biol Med* 2002; 13: 300-9.
- Sonis ST: Pathobiology of oral mucositis: novel insights and opportunities. *J Support Oncol* 2007; 5(9): 3-11.
- McCarthy GM, Awde JD, Ghandi H, Vincent M and Kocha WI: Risk factors associated with mucositis in cancer patients receiving 5-fluorouracil. *Oral Oncol* 1998; 34(6): 484-90.
- Bamba S, Andoh A, Yasui H, Araki Y and Bamba T: Matrix metalloproteinase-3 secretion from human colonic subepithelial myofibroblasts: role of interleukin-17. *J Gastroenterol* 2013; 38: 548-54.

22. Sasakica M, Kashima M, Ito T, Watanabe A and Izumiyama N: Differential regulation of metalloproteinase production, proliferation and chemotaxis of human lung fibroblasts by PDGF, interleukin-1b and TNF- $\alpha$ . *Mediators Inflamm* 2000; 9: 155-60.
23. Wearing HJ and Sherratt JA: Keratinocyte growth factor signaling: a mathematical model of dermal-epidermal interaction in epidermal wound healing. *Math Bio Sci* 2000; 165: 41-62.
24. Srivastava JK, Shankar E and Gupta S: Chamomile: A herbal medicine of the past with bright future. *Mol Med Rep* 2010; 3(6): 895-901.
25. Curra M, Martins MA, Lauxen IS, Pellicoli AC, Sant' Ana Filho M: Effect of topical chamomile on immunohistochemical levels of IL-1 $\beta$  and TNF- $\alpha$  in 5-fluorouracil-induced oral mucositis in hamsters. *Cancer Chemother Pharmacol* 2013; 71(2): 293-9.
26. Braga FT, Santos AC, Bueno PC, Silveira RC and Santos CB: *Chamomilla recutita* in the prevention and treatment of oral mucositis in patients undergoing hematopoietic stem cell transplantation: A randomized, controlled, Phase II clinical trial. *Cancer Nurs* 2015; 38(4): 322-9.
27. Darout IA, Christy AA, Skaug N and Egeberg PK: Identification and quantification of some potentially antimicrobial anionic components in miswak extract. *Indian J Pharmacol* 2000; 32: 11-14.
28. Monforte MT, Miceli N, Mondello MR, Sanogo R and Rossitto A: Antiulcer activity of *Salvadora persica* on experimental ASA-induced ulcer in rats: Ultra structural modifications. *Pharma Biol* 2001; 39: 289-92.
29. Chang JW, Choi JW, Lee BH, Park JK and Shin YS: Protective effects of Korean red ginseng on radiation-induced oral mucositis in a preclinical rat model. *Nutr Cancer* 2014; 66(3): 400-7.
30. Kuduban O, Mazlumoglu MR, Kuduban SD, Erhan E, Cetin N: The effect of *Hippophae rhamnoides* extract on oral mucositis induced in rats with methotrexate. *J Appl Oral Sci* 2016; 24(5): 423-430.
31. Koohi-Hosseiniabadi O, Andisheh-Tadbir A, Bahadori P, Sepehrimanesh M and Mardani M: Comparison of the therapeutic effects of the dietary and topical forms of *Zizyphus jujuba* extract on oral mucositis induced by 5-fluorouracil: A golden hamster model. *J Clin Exp Dent* 2015; 7(2): e304-9.
32. Mardani M, Afra SM, Tanideh N, Tadbir AA and Modarresi F: Hydroalcoholic extract of *Carum carvi* L. in oral mucositis: a clinical trial in male golden hamsters. *Oral Dis* 2016; 22(1): 39-45.
33. Watanabe S, Suemaru K, Takechi K, Kaji H and Imai K: Oral mucosal adhesive films containing royal jelly accelerate recovery from 5-fluorouracil-induced oral mucositis. *J Pharmacol Sci* 2013; 121(2): 110-8.
34. Rashad UM, Al-Gezawy SM, El-Gezawy E and Azzaz AN: Honey as topical prophylaxis against radio chemotherapy-induced mucositis in head and neck cancer. *J Laryngol Otol* 2009; 123(2): 223-8.
35. Viuda-Martos M, Ruiz-Navajas Y, Fernández-López J and Pérez-Alvarez JA: Functional properties of honey, propolis, and royal jelly. *J Food Sci* 2008; 73: R117-24.
36. Kawashima K, Fujimura Y, Makino T and Kano Y: Pharmacological properties of traditional medicine (XXXII): protective effects of hangeshashinto and the combinations of its major constituents on gastric lesions in rats. *Biol Pharm Bull* 2006; 29(9): 1973-5.
37. Yoshida K, Yoshioka M, Okamura H, Moriyama S and Kawazoe K: Preventive effect of Daiokanzoto (TJ-84) on 5-fluorouracil-induced human gingival cell death through the inhibition of reactive oxygen species production. *PLoS One* 2014; 9(11): e112689.
38. Hatakeyama H, Takahashi H, Oridate N, Kuramoto R and Fujiwara K: Hangeshashinto improves the completion rate of chemo-radiotherapy and the nutritional status in patients with head and neck cancer. *ORL J Otorhinolaryngol Relat Spec* 2015; 77(2): 100-8.
39. Matsuda C, Munemoto Y, Mishima H, Nagata N and Oshiro M: Double-blind, placebo-controlled, randomized phase II study of TJ-14 (Hangeshashinto) for infusional fluorinated-pyrimidine-based colorectal cancer chemotherapy-induced oral mucositis. *Cancer Chemo ther Pharmacol* 2015; 76(1): 97-103.
40. Conklin KA: Chemotherapy-associated oxidative stress: impact on chemotherapeutic effectiveness. *Integr Cancer Ther* 2004; 3(4): 294-300.
41. Valerie K, Yacoub A, Hagan MP, Curiel DT and Fisher PB: Radiation-induced cell signaling: inside-out and outside-in. *Mol Cancer Ther* 2007; 6(3): 789-801.
42. Glasauer A and Chandel NS: Targeting antioxidants for cancer therapy. *Biochem Pharmacol* 2014; 92(1): 90-101.
43. Zhang LJ, Yeh SF, Yu YT, Kuo LM, Kuo YH: Antioxidative flavonol glucuronides and anti-HBs Ag flavonol from *R. rotundifolia*. *J Trad Comp Med* 2011; 1(1): 57-63.
44. Andersson SC, Rumpunen K, Johansson E and Olsson ME: Tocopherols and tocotrienols in sea buckthorn (*Hippophae rhamnoides* L.) berries during ripening. *J Agric Food Chem* 2008; 56(15): 6701-6.
45. Liu J, Chen B and Yao S: Simultaneous analysis and identification of main bioactive constituents in the extract of *Zizyphus jujuba* var. *sapinosa* (*Zizyphi spinosi* semen) by high-performance liquid chromatography-photodiode array detection-electrospray mass spectrometry. *Talanta* 2007; 71: 668-75.
46. Yamashita T, Araki K, Tomifuji M, Kamide D and Tanaka Y: Traditional Japanese medicine-Hangeshashinto (TJ-14)-alleviates chemoradiation-induced mucositis and improves rates of treatment completion. *Support Care Cancer* 2015; 23(1): 29-35.
47. Kono T, Kaneko A, Matsumoto C, Miyagi C and Ohbuchi K: Multitargeted effects of hangeshashinto for treatment of chemotherapy-induced oral mucositis on inducible prostaglandin E2 production in human oral keratinocytes. *Integr Cancer Ther* 2014; 13(5): 435-45.
48. Rankin KV, Jones DJ and Redding SW: Oral health in cancer therapy. *Texas Cancer Council, Edition 2<sup>nd</sup>*, 2004; 43-52.
49. Sonis ST: Mucositis as a biological process: a new hypothesis for the development of chemotherapy-induced stomato toxicity. *Oral Oncol* 1998; 34: 39-43.
50. Bhat PK, Kumar A and Sarkar S: Assessment of immediate antimicrobial effect of miswak extract and toothbrush on cariogenic bacteria – A clinical study. *JoAOR* 2012; 3(1): 13-18.
51. Mohammed SG: Comparative study of in vitro antibacterial activity of miswak extracts and different toothpastes. *American Journal of Agricultural and Biological Sciences* 2013; 8 (1): 82-88.
52. Jelvehgaran Esfahani Z, Kadkhoda Z, Eshraghi SS and SalehiSurmaghi MH: Antibacterial effect of an herbal product persica on *Porphyromonas gingivalis* and *Aggregatibacter actinomycetemcomitans*: an in-vitro study. *J Dent (Tehran)* 2014; 11(4): 464-72.
53. Naseem S, Hashmi K, Fasih F, Sharafat S and Khanani R: In-vitro evaluation of the antimicrobial effect of miswak against common oral pathogens. *Pak J Med Sci* 2014; 30(2): 398-403.

54. Alali F and Al-Lafi T: GC-MS analysis and bioactivity testing of the volatile oil from the leaves of the toothbrush tree *S. persica* L. Nat Prod Res 2003; 17(3): 189- 94.
55. Almas K: The effect of *Salvadora persica* extract (miswak) and chlorhexidine gluconate on human dentin: A SEM study. J Contemp Dent Prac 2005; 3: 27-35.
56. Almas K, Skaug N and Ahmad I: An *in-vitro* antimicrobial comparison of miswak extract with commercially available non-alcohol mouth rinses. Int J Dent Hyg 2005; 3: 18-24.
57. Rajesh V, Suresh P, Anil B, Brijesh K and Priyanka P: *Salvadora persica* L. (Tooth Brush Tree): A review. JPR 2009; 2(12): 1809-1812.
58. Talwar S, House R, Sundaramurthy S, Balasubramanian S and Yu H: Inhibition of caspases protects mice from radiation-induced oral mucositis and abolishes the cleavage of RNA-binding protein HuR. J Biol Chem 2014; 289(6): 3487-3500.
59. Kwon Y: Mechanism-based management for mucositis: option for treating side effects without compromising the efficacy of cancer therapy. Onco Targets Ther 2016; 9: 2007-16.
60. Hitomi S, Ono K, Terawaki K, Matsumoto C and Mizuno K: [6]-gingerol and [6]-shogaol, active ingredients of the traditional Japanese medicine hangeshashinto, relief oral ulcerative mucositis-induced pain *via* action on Na<sup>+</sup> channels. Pharmacol Res 2016; 117: 288-302.
61. Redding SW: Cancer therapy-related oral mucositis. J Dent Educ 2005; 69(8): 919-29.
62. Meropol NJ, Somer RA, Gutheil J, Pelley RJ and Modiano MR: Randomized phase I trial of recombinant human keratinocyte growth factor plus chemotherapy: potential role as a mucosal protectant. J Clin Oncol 2003; 21: 1452-8.

**How to cite this article:**

Zakaria S: Natural remedies target different therapeutic pathways in oral mucositis induced by cancer chemo or radiotherapy. Int J Pharmacognosy 2017; 4(8): 250-56. doi link: [http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.4\(8\).250-56](http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.4(8).250-56).

This Journal licensed under a Creative Commons Attribution-Non-commercial-Share Alike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)