



Received on 14 October 2019; received in revised form, 28 October 2019; accepted, 29 October 2019; published 31 October 2019

## ANALYSIS OF FATTY ACID COMPOSITIONS OF APRICOT KERNEL OILS FOR COSMETIC PURPOSE WITH GAS CHROMATOGRAPHY-MASS SPECTROMETRY METHOD

Bilal Yilmaz <sup>\*1</sup> and Ismail Celik <sup>2</sup>

Department of Analytical Chemistry <sup>1</sup>, Faculty of Pharmacy, Department of Criminalistic <sup>2</sup>, Graduate School of Natural and Applied Sciences, Ataturk University, Erzurum, Turkey.

### Keywords:

Fatty acid, Apricot kernel oil, *Prunus armeniaca* (L.), Gas chromatography-mass spectrometry

### Correspondence to Author:

**Prof. Dr. Bilal YILMAZ**

Department of Analytical Chemistry,  
Faculty of Pharmacy, Ataturk  
University, Erzurum, Turkey.

**E-mail:** yilmazb@atauni.edu.tr

**ABSTRACT:** In this study, fatty acid compositions of apricot kernel oils for cosmetic purpose were analyzed. Qualitative and quantitative analysis of fatty acids in apricot kernel oils were made by using gas chromatography-mass spectrometry (GC-MS). In this analysis, apricot kernel oils -which trademarked Karden, Organix and Biovitals-supplied from a herbalist were used. The oils were converted to the methyl ester with KOH hydrolysis. Just after the derivatization process, the oils were analyzed by GC-MS. In conclusion, it was determined that these apricot kernel oils which belong to three different trademarks have a similar fatty acid profile, however, there was no specific standard between them.

**INTRODUCTION:** Apricot known as *Prunus armeniaca* (L.) **Fig. 1** which belongs to Rosaceae family is a fruit that grows in the apricot tree. Although apricot is distributed geographically to almost all parts of the world, it has mostly spread to Europe, Africa, America and Central Asia continents, and found growth areas in countries close to the Mediterranean. Apricot is one of the most traded fruits in the World <sup>1</sup>. Although, apricots are produced anywhere in Turkey, the majority of apricots are produced in the Eastern Anatolia Region of Turkey. The most important apricot production centers are Malatya, Erzincan, Aras Valley (Iğdir-Kagizman), Mersin (Mut), Elazığ, Sivas, Kahramanmaraş, Kayseri, Ankara, Hatay, Nigde, and Nevşehir <sup>2</sup>.

The apricot fruit which has different types contains vitamins A, B, C and E also has beta-carotene, high levels of potassium, iron, protein, pectin, and some organix substances. The apricot kernel has an important place in terms of nutritional value since it contains 40-50% edible fat besides the abundant protein and mineral content. Apricot kernel oil is rich in unsaturated fatty acids, and important parts of these are oleic and linoleic fatty acids <sup>3</sup>. An active ingredient named Amygdalin, one of the components of apricot kernel and one of the most effective herbal ingredients used against cancer is the main component of a cancer drug called "Laetrile" on the market.

In addition, apricot kernel oil is used in the treatment of asthma, cough, constipation, and as massage oil, it's also used in make-up products owing to its nourishing and moisturizing properties <sup>4</sup>. The general formulas of saturated fatty acids are shown as  $C_nH_{2n}O_2$  or  $CH_3(CH_2)_n-COOH$ . Generally, their chains have an even number of carbon atoms. Carbon atoms are connected to each other in a single bond. Fatty acids containing one or

	<p><b>QUICK RESPONSE CODE</b></p>
	<p>DOI: 10.13040/IJPSR.0975-8232.IJP.6(10).340-46</p>
<p>The article can be accessed online on <a href="http://www.ijpjournal.com">www.ijpjournal.com</a></p>	
<p>DOI link: <a href="http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.6(10).340-46">http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.6(10).340-46</a></p>	

more double bonds between different carbon atoms on carbon chains are referred to as unsaturated fatty acids<sup>5</sup>. Linoleic acid, linolenic acid and arachidonic acid are essential for humans; that is, they cannot be synthesized in the body, they must be taken through nutrition.



FIG. 1: *PRUNUS ARMENIACA* (L.) FRUIT

## MATERIALS AND METHODS:

**Chemicals and Reagents:** Analytical purity chemicals were used in the study. The n-hexane and methanol used in the derivatization process were purchased from Sigma-Aldrich (Sigma Aldrich Chemie GmbH, Steinheim, Germany). Karden, Organix and Biovitals brand apricot kernel oils used in the analysis were obtained from a herbal store.

**GC-MS System:** Chromatographic analyzes were carried out on an Agilent 6890N gas chromatography system equipped with 5977 series mass selective detector, 7693 series autosampler and chemstation (Agilent Technologies, Palo Alto, CA). HP-5 MS column with 0.25  $\mu\text{m}$  film thickness (30 m  $\times$  0.25 mm I.D., USA) was used for separation. The temperatures of the inlet, transfer line and detector were 50, 100, 180, 220 and 300  $^{\circ}\text{C}$ , respectively.

**GC-MS Conditions:** Different temperature programs were investigated for GC-MS method. The end of this investigation, the temperature program of the GC/MS was as follows: initial temperature was 50  $^{\circ}\text{C}$ , held for 1 min, increased to 100  $^{\circ}\text{C}$  at a rate of 20  $^{\circ}\text{C}/\text{min}$  held for 1 min, increased to 180  $^{\circ}\text{C}$  at a rate of 10  $^{\circ}\text{C}/\text{min}$  held for 1 min, increased to 220  $^{\circ}\text{C}$  at a rate of 5  $^{\circ}\text{C}/\text{min}$  held for 5 min, and finally to 300  $^{\circ}\text{C}$  at a rate of 10  $^{\circ}\text{C}/\text{min}$  and held for 5.5 min.

**Derivatization Procedure:** In the GC study, volatilization of fatty acids was carried out by conversion to methyl ester derivatives. Because in this method, samples must be volatile or turn into volatile. For this purpose, according to the following procedure, the oil samples to be analyzed were volatilized by methylation reaction. 0.1 ml oil sample was taken into a 15 ml plastic centrifuge tube and 10 ml n-hexane was added to it. Then, the centrifuge tube was shaken strongly. After that, 0.5 ml of 2 M methanol KOH solution was added to the tube. The centrifuge tube was shaken strongly again. After the sample tube was kept in the dark for 2 h, 1 ml of the organic phase in the sample tube was put in the GC vial and GC-MS analysis was performed.

**Identification of Components:** Identification was based on the molecular structure, molecular mass and calculated fragments. Interpretation on mass spectrum GC-MS was conducted using the database on National Institute of Standard and Technology (NIST) having more than 62,000 patterns. The name, molecular weight and structure of the components of the test materials were ascertained. The relative percentage amount of each component was calculated by comparing its average peak area to the total areas. The spectrum of the unknown component was compared with the spectrum of the component stored in the NIST Library Version (2005), Software, Turbo mass 5.2.

## RESULTS AND DISCUSSION:


**Method Development and Optimization:** In this study, GC-MS method was developed for the analysis of fatty acid compositions of apricot kernel oils used for cosmetic purposes and these method conditions were optimized. The capillary column coated with 5% phenyl, 95% dimethylpolysiloxane is a good choice for separation of these analytes since, they elute as symmetrical peaks at a wide range of concentrations. Different temperature programs were investigated for GC oven. The end of this investigation, the best temperature program was selected for a good separation. The temperature programs of the GC oven were as follows: initial temperature 50  $^{\circ}\text{C}$ , held for 1 min, increased to 100  $^{\circ}\text{C}$  at a rate of 200  $^{\circ}\text{C}/\text{min}$  held for 1 min, increased to 180  $^{\circ}\text{C}$  at a rate of 10  $^{\circ}\text{C}/\text{min}$  held for 1 min, increased to 220  $^{\circ}\text{C}$  at a rate of 5  $^{\circ}\text{C}/\text{min}$  held for 5 min, and finally to 300  $^{\circ}\text{C}$  at a

rate of 10 °C/min and held for 5.5 min. The split less injection mode was chosen. The injector volume was 1 µl in split less mode and the carrier gas was helium at a flow rate of 1 ml/min.

**GC-MS Analysis:** The more precise information in qualitative analysis can be obtained by GC-MS. For quantitative determination, gas chromatography with flame ionization detector (GC-FID) and GC-MS are preferred. GC-MS is one of the best techniques to identify the constituents of volatile matter, long chain, branched chain hydrocarbons, alcohols, acids, esters *etc.*


The GC-MS analysis of three different apricot kernel oils revealed the presence of various fatty acids. The identification of the fatty acids was confirmed based on the retention time, molecular formula and mass spectrum. The major fatty acid constituents as methyl esters of Organix branded

apricot kernel oil are presented as compound chromatogram in **Fig. 2**.




**FIG. 2: GC-MS CHROMATOGRAM OF APRICOT KERNEL OIL BELONGS TO ORGANIX TRADEMARK**

Biovitals and Karden branded apricot kernel oils' GC-MS chromatograms are shown in **Fig. 3** and **Fig. 4** respectively.



**FIG. 3: GC-MS CHROMATOGRAM OF APRICOT KERNEL OIL BELONGS TO BIOVITALS TRADEMARK**



**FIG. 4: GC-MS CHROMATOGRAM OF APRICOT KERNEL OIL BELONGS TO KARDEN TRADEMARK**

The choice of methods to be used in the analysis of fatty acids is made according to the environment and laboratory conditions in which the analysis is performed. After the analytical method has been developed for an analysis, the optimization conditions of the method must be determined to demonstrate that this method is applicable. GC-MS analysis revealed the presence of various fatty acid compounds as methyl esters in apricot kernel oils. Peak number, retention time, compound name and molecular formula are stated in **Table 1**. According to our analysis results, major compounds of apricot kernel oil are linoleic, oleic, palmitic and stearic fatty acids. Linoleic acid is a fatty acid. It is an 18-

carbon chain with two double bonds in cis configuration. A shorthand notation like "18:2 (n-6)" or "18:2 cis-9, 12" may be used in literature<sup>6</sup>. It typically occurs in nature as a triglyceride ester; free fatty acids, the form not combined with glycerol to form triglyceride, are typically low in foods<sup>7</sup>. It is very soluble in acetone, benzene, diethyl ether and ethanol<sup>8</sup>. It is found in the lipids of cell membranes. It is abundant in many nuts, fatty seeds (flax seeds, hemp seeds, poppy seeds, sesame seeds, *etc.*) and their derived vegetable oils; comprising over half (by weight) of poppy seed, safflower, sunflower, corn, and soybean oils<sup>9</sup>.

**TABLE 1: FATTY ACIDS COMPOUNDS OF THREE DIFFERENT APRICOT KERNEL OILS**

Peak	Retention time (min)	Fatty acid	Molecular formula	% of total		
				Organix	Biovitals	Karden
1	23.89	Methyl Tetradecanoate	C <sub>15</sub> H <sub>30</sub> O <sub>2</sub>	0.11	0.13	0.29
2	26.62	Methyl Pentadecanoate	C <sub>16</sub> H <sub>32</sub> O <sub>2</sub>	0.06	0.09	-
3	28.36	Methyl 7, 10-Hexadecadienoate	C <sub>17</sub> H <sub>30</sub> O <sub>2</sub>	0.05	0.08	-
4	28.64	Methyl Palmitoleate	C <sub>17</sub> H <sub>32</sub> O <sub>2</sub>	-	0.39	0.42
5	29.22	Methyl Palmitate	C <sub>17</sub> H <sub>34</sub> O <sub>2</sub>	-	8.57	15.91
6	31.67	Methyl Heptadecanoate	C <sub>18</sub> H <sub>36</sub> O <sub>2</sub>	-	0.09	-
7	33.22	Methyl Linoleate	C <sub>19</sub> H <sub>34</sub> O <sub>2</sub>	-	-	24.59
8	33.48	Methyl 11-Octadecenoate	C <sub>19</sub> H <sub>36</sub> O <sub>2</sub>	-	-	30.14
9	33.75	Methyl Linolenate	C <sub>19</sub> H <sub>32</sub> O <sub>2</sub>	-	-	0.75
10	33.95	Methyl Oleate	C <sub>19</sub> H <sub>36</sub> O <sub>2</sub>	71.47	51.77	-
11	34.01	Methyl Stearate	C <sub>19</sub> H <sub>38</sub> O <sub>2</sub>	4.87	5.16	2.77
12	34.67	Oleic Acid	C <sub>18</sub> H <sub>34</sub> O <sub>2</sub>	-	-	1.06
13	34.89	Trans-13-Octadecanoic Acid	C <sub>18</sub> H <sub>34</sub> O <sub>2</sub>	0.83	-	-
14	34.98	Ethyl-9, 12-Octadecadienoate	C <sub>20</sub> H <sub>36</sub> O <sub>2</sub>	0.21	-	-
15	35.09	Ethyl Oleate	C <sub>20</sub> H <sub>38</sub> O <sub>2</sub>	0.58	-	-
16	41.55	Methyl 11-Eicosenoate	C <sub>21</sub> H <sub>40</sub> O <sub>2</sub>	2.42	2.69	2.33
17	42.27	Methyl Arachisate	C <sub>21</sub> H <sub>42</sub> O <sub>2</sub>	0,96	1.18	-
18	42.92	Glycidyl Oleate	C <sub>21</sub> H <sub>38</sub> O <sub>3</sub>	2.28	0.19	1.76
19	48.71	Methyl Erucate	C <sub>23</sub> H <sub>44</sub> O <sub>2</sub>	0.79	-	-
20	49.74	Methyl Behenate	C <sub>23</sub> H <sub>46</sub> O <sub>2</sub>	0.49	-	-
21	41.55	Methyl Nervonate	C <sub>25</sub> H <sub>48</sub> O <sub>2</sub>	-	0.15	-
22	42.27	Methyl Lignocerate	C <sub>25</sub> H <sub>50</sub> O <sub>2</sub>	-	-	0.26

The consumption of linoleic acid is vital to proper health, as it is an essential fatty acid<sup>10</sup>. In rats, a diet deficient in linoleate (the salt form of the acid) has been shown to cause mild skin scaling, hair loss<sup>11</sup>, and poor wound healing<sup>12</sup>. However, chronic consumption of high levels of linoleic acid may be associated with the development of ulcerative colitis<sup>13</sup>. Linoleic acid has become increasingly popular in the beauty products industry because of its beneficial properties on the skin. Research points to linoleic acid's anti-inflammatory, acne reductive, skin-lightening and moisture retentive properties when applied topically on the skin<sup>14-17</sup>.

Since, it is among the essential fatty acids, it is worth mentioning about linolenic acid as well. Linolenic acid is divided into two groups; alpha linolenic acid and gamma linolenic acid. Alpha linolenic acid or ALA ( $\alpha$ -linolenic acid), (from Greek linon, meaning flax), is an n-3, or omega-3 essential fatty acid. ALA is found in many seeds and oils, including flaxseed, walnuts, chia, hemp, and many common vegetable oils. ALA can only be obtained by humans through their diets because the absence of the required 12- and 15-desaturase enzymes makes "de novo synthesis" from stearic acid impossible. Eicosapentaenoic acid (EPA; 20:5, n-3) and docosahexaenoic acid (DHA; 22:6, n-3) are readily available from fish and algae

oil and play a vital role in many metabolic processes. These can also be synthesized by humans from dietary  $\alpha$ -linolenic acid: ALA  $\rightarrow$  stearidonic acid  $\rightarrow$  eicosatetraenoic acid  $\rightarrow$  eicosapentaenoic acid  $\rightarrow$  docosapentaenoic acid  $\rightarrow$  9, 12, 15, 18, 21-tetracosapentaenoic acid  $\rightarrow$  6, 9, 12, 15, 18, 21-yetracosahexaenoic acid  $\rightarrow$  docosahexaenoic acid, but with an efficiency of only a few percent<sup>18</sup>. Because the efficacy of n-3 long-chain polyunsaturated fatty acid (LC-PUFA) synthesis decreases down the cascade of  $\alpha$ -linolenic acid conversion, DHA synthesis from ALA is even more restricted than that of EPA<sup>19, 20</sup>. According to one scientific review, higher ALA consumption is associated with a moderately lower risk of cardiovascular disease, but wide variation in results across multiple studies highlights the need for additional research before drawing firm conclusions<sup>21</sup>.

Gamma-Linolenic acid or GLA ( $\gamma$ -linolenic acid), (INN and USAN gamolenic acid) is a fatty acid found primarily in vegetable oils. When acting on GLA, arachidonate 5-lipoxygenase produces no leukotrienes and the conversion by the enzyme of arachidonic acid to leukotrienes is inhibited. GLA is categorized as an n-6 (also called  $\omega$ -6 or omega-6) fatty acid, meaning that the first double bond on the methyl end (designated with n or  $\omega$ ) is the sixth

bond. In physiological literature, GLA is designated as 18:3 (n-6). GLA is a carboxylic acid with an 18-carbon chain and three cis double bonds. It is an isomer of  $\alpha$ -linolenic acid, which is a polyunsaturated n-3 (omega-3) fatty acid, found in rapeseed canola oil, soybeans, walnuts, flax seed (linseed oil), perilla, chia, and hemp seed. GLA is obtained from vegetable oils such as evening primrose (*Oenothera biennis*) oil (EPO), black-currant seed oil, borage seed oil, and hemp seed oil. GLA is also found in varying amounts in edible hemp seeds, oats, barley<sup>22</sup> and spirulina. Normal safflower (*Carthamus tinctorius*) oil does not contain GLA, but genetically modified GLA safflower oil available in commercial quantities since 2011 contains 40% GLA<sup>23</sup>.

Borage oil contains 20% GLA, evening primrose oil ranges from 8% to 10% GLA, and black-currant oil contains 15-20%<sup>24</sup>. It also constitutes 12.23% of the fats from the fruit of the durian species *Durio graveolens*<sup>25</sup>. The human body produces GLA from linoleic acid (LA). This reaction is catalyzed by  $\Delta 6$ -desaturase (D6D), an enzyme that allows the creation of a double bond on the sixth carbon counting from the carboxyl terminus. LA is consumed sufficiently in most diets, from such abundant sources as cooking oils and meats. However, a lack of GLA can occur when there is a reduction of the efficiency of the D6D conversion (for instance, as people grow older or when there are specific dietary deficiencies) or in disease states wherein there is excessive consumption of GLA metabolites<sup>26</sup>.

Oleic acid is a fatty acid that occurs naturally in various animal and vegetable fats and oils. It is odourless, colourless oil, although commercial samples may be yellowish. In chemical terms, oleic acid is classified as a monounsaturated omega-9 fatty acid, abbreviated with a lipid number of 18:1 cis-9. It has the formula  $\text{CH}_3(\text{CH}_2)_7\text{CH}=\text{CH}(\text{CH}_2)_7\text{COOH}$ <sup>27</sup>. The name derives from the Latin word *oleum*, which means oil<sup>28</sup>. It is the most common fatty acid in nature. The salts and esters of oleic acid are called as oleates. Fatty acids (or their salts) do not often occur as such in biological systems. Instead fatty acids like oleic acid occur as their esters, commonly triglycerides, which are the greasy materials in many natural oils. Oleic acid is the most common monounsaturated fatty acid in

nature. It is found in fats (triglycerides), the phospholipids that make membranes, cholesterol esters, and wax esters<sup>29</sup>. Triglycerides of oleic acid comprise the majority of olive oil. It is abundantly present in many animal fats, constituting 37 to 56% of chicken and turkey fat<sup>30</sup>. Oleic acid is the most abundant fatty acid in human adipose tissue<sup>31</sup> and second in abundance in human tissues overall only to palmitic acid<sup>32</sup>. Palmitic acid, or hexadecanoic acid in IUPAC nomenclature, is the most common saturated fatty acid found in animals, plants and microorganisms<sup>33</sup>. Palmitic acid is naturally produced by a wide range of other plants and organisms, typically at low levels. It is naturally present in butter, cheese, milk, and meat, as well as cocoa butter, soybean oil and sunflower oil. Karukas also contain 44.90% palmitic acid<sup>34</sup>. According to the World Health Organization, evidence is "convincing" that consumption of palmitic acid increases the risk of developing cardiovascular disease<sup>35</sup>, based on studies indicating that it may increase LDL levels in the blood.

Stearic acid is a saturated fatty acid with an 18-carbon chain and has the IUPAC name octadecanoic acid. It is a waxy solid and its chemical formula is  $\text{C}_{17}\text{H}_{35}\text{CO}_2\text{H}$ . The salts and esters of stearic acid are called stearates. As its ester, stearic acid is one of the most common saturated fatty acids found in nature following palmitic acid<sup>36</sup>. Stearic acid is obtained from fats and oils by the saponification of the triglycerides using hot water (about 100 °C).

The resulting mixture is then distilled<sup>37</sup>. Commercial stearic acid is often a mixture of stearic and palmitic acids, although purified stearic acid is available. Fats and oils rich in stearic acid are more abundant in animal fat (up to 30%) than in vegetable fat (typically <5%). The important exceptions are the foods cocoa butter and shea butter, where the stearic acid content (as a triglyceride) is 28-45%<sup>38</sup>. When the results obtained from these analyzes are compared with the data in previous studies, it is seen that the apricot kernel oils' fatty acid profiles are generally similar. Three different apricot kernel oils we studied on have a similar fatty acid profile, but there is no specific standard between them.

**CONCLUSION:** In this study, fatty acid constituents of apricot kernel oils used for cosmetic purposes were analyzed qualitatively and quantitatively by GC-MS method. In the literature, the analysis of fatty acid constituents of cosmetic oils is rare. It has been observed that the fatty acid constituents of cooking oils are mostly examined. Karden, Organix and Biovitals branded apricot kernel oils were used for the analysis. Fatty acids were derivatized by converting them into volatile methyl esters by KOH hydrolysis procedure. GC-MS method was preferred for analysis. According to our analysis results, major compounds of apricot kernel oil are oleic, linoleic, palmitic and stearic fatty acids. While Organix and Biovitals branded apricot kernel oils are rich in oleic acid, Karden is rich in linoleic acid. However, it has been found that there is no standard between these three products in terms of the proportion of fatty acid components.

**ACKNOWLEDGEMENT:** Nil

**CONFLICTS OF INTEREST:** Nil

## REFERENCES:

- Orhan I, Koca U, Aslan S, Kartal M and Kusmenoglu S: Fatty acid analysis of some Turkish apricot kernel oils by GC and GC-MS techniques. *Turkish Journal of Pharmaceutical Sciences* 2008; 5(1): 29-34.
- Fidan I: Economic analysis of apricot production in Iğdir province. MS Thesis, Atatürk University, Erzurum, Turkey 2009.
- Demir NK: Effect of enzyme on apricot seed oil extraction: optimization of extraction conditions. MS Thesis, İstanbul Technical University, İstanbul, Turkey 2011.
- Soylu SM: The effect of amygdalin on electrocardiogram and blood nitric oxide levels in guinea pigs. PhD Thesis, Ankara University, Ankara, Turkey 2006.
- Basoglu F: *Cooking Oil Technologies*, ed 1<sup>st</sup> Nobel Publisher, Ankara 2006; 43-97.
- Eldor J: Linoleic acid: The code of life? *Journal of Health Science and Development* 2018; 1 (2): 18-32.
- Mattes RD: Is there a fatty acid taste? *Annual Review of Nutrition* 2009; 29: 305-27.
- Haynes WM: *CRC Handbook of Chemistry and Physics*, ed. 97<sup>th</sup> CRC Press, Boca Raton 2016; 3-338.
- USDA national nutrient database for standard reference, release 20. U.S. Department of Agriculture, Agricultural Research Service 2007.
- Whelan J and Fritsche K: Linoleic acid. *Advances in Nutrition* 2013; 4 (3): 311-312.
- Cunnane S and Anderson M: Pure linoleate deficiency in the rat: influence on growth, accumulation of n-6 polyunsaturates and (1-14C) linoleate oxidation. *The Journal of Lipid Research* 1997; 38 (4): 805-12.
- Ruthig DJ and Meckling-Gill KA: Both (n-3) and (n-6) fatty acids stimulate wound healing in the rat intestinal epithelial cell line, IEC-6. *Journal of Nutrition* 1999; 129(10): 1791-98.
- Gastrointestinal Society, Canadian Society of Intestinal Research: Role of fats in ulcerative colitis. Canada 2010.
- Diezel WE, Schulz E, Skanks M and Heise H: Plant oils: Topical application and anti-inflammatory effects (croton oil test). *Dermatologische Monatsschrift* 1993; 179: 173.
- Letawe C, Boone M and Pierard GE: Digital image analysis of the effect of topically applied linoleic acid on acne microcomedones. *Clinical and Experimental Dermatology* 1998; 23 (2): 56-58.
- Ando H, Ryu A, Hashimoto A, Oka M and Ichihashi M: Linoleic acid and  $\alpha$ -linolenic acid lightens ultraviolet-induced hyperpigmentation of the skin. *Archives of Dermatological Research* 1998; 290(7): 375-81.
- Darmstadt GL, Mao-Qiang M, Chi E, Saha SK, Ziboh VA, Black RE, Santosham M and Elias PM: Impact of tropical oils on the skin barrier: possible implications for neonatal health in developing countries. *Acta Paediatrica* 2002; 91 (5): 546-54.
- Breanne MA and David WL Ma: Are all n-3 polyunsaturated fatty acids created equal. *Lipids in Health and Disease* 2009; 8 (33): 33.
- Innis SM: Fatty acids and early human development. *Early Human Development* 2007; 83(12): 761-66.
- Burdge GC and Calder PC: Conversion of alpha-linolenic acid to longer-chain polyunsaturated fatty acids in human adults. *Reproduction Nutrition Development* 2005; 45(5): 581-97.
- Pan A, Chen M, Chowdhury R, Wu JH, Sun Q, Campos H, Mozaffarian D and Hu FB:  $\alpha$ -Linolenic acid and risk of cardiovascular disease: a systematic review and meta-analysis. *The American Journal of Clinical Nutrition* 2012; 96 (6): 1262-73.
- Qureshi AA, Schnoes HK, Din ZZ and Peterson DM: Determination of the structure of cholesterol inhibitor II isolated from high-protein barley flour (HPBF). *Federation Proceedings* 1984; 43(7): 2626.
- Nykiforuk C and Shewmaker C: High level accumulation of gamma linolenic acid in transgenic safflower (*Carthamus tinctorius*) seeds. *Transgenic Research* 2011; 21(2): 367-381.
- Flider FJ: GLA: Uses and new sources. *Inform* 2005; 16 (5): 279-82.
- Nasaruddin MH, Noor NQIM and Mamat H: Proximate and fatty acid composition of Sabah Yellow Durian (*Durio graveolens*) Sains Malaysiana 2013; 42(9): 1283-88.
- Horrobin DF: Fatty acid metabolism in health and disease: the role of delta-6-desaturase. *The American Journal of Clinical Nutrition* 1993; 57 (5): 732-37.
- Thomas A: Fats and fatty oils. *Ullmann's Encyclopedia of Industrial Chemistry* 2000.
- Bailey and Bailey, Dorothy and Kenneth: An etymological dictionary of chemistry and mineralogy. *Nature* 1929; 124 (3134): 789-90.
- Ntambi, James M, Miyazaki and Makoto: Recent insights into stearoyl-CoA desaturase-1. *Current Opinion in Lipidology* 2003; 14(3): 255-61.
- Nutter MK, Lockhart EE and Harris RS: The chemical composition of depot fats in chickens and turkeys. *Oil and Soap* 1943; 20(11): 231-34.
- Kokatnur MG, Oalman MC, Johnson WD, Malcom GT and Strong JP: Fatty acid composition of human adipose tissue from two anatomical sites in a biracial community *The American Journal of Clinical Nutrition* 1979; 32(11): 2198-205.

32. Oliveira AF, Chunhab DA, Ladriere L, Igoillo-Esteve M, Bugliani M, Marchetti P and Cnop M: *In-vitro* use of free fatty acids bound to albumin: A comparison of protocols. *Bio Feedback. Bio Techniques* 2015; 58(5): 228-33.
33. Gunstone FD, John LH and Albert JD: *The Lipid Handbook*, ed. 3<sup>rd</sup> CRC Press, Boca Raton 2007.
34. Purwanto Y and Munawaroh E: Ethnobotany types of pandanaceae as foodstuffs in Indonesia. *Berkala Penelitian Hayati* 2010; 5(A): 97-08.
35. World Health Organization: Diet, nutrition and the prevention of chronic diseases. WHO Technical Report Series 916, Report of a Joint WHO/FAO Expert Consultation, Geneva 2003; 88.
36. Gunstone FD, John LH and Albert JD: *The Lipid Handbook*, ed. 3<sup>rd</sup> CRC Press, Boca Raton 2007.
37. David JA, Sabine B, Ralf C, Georg F, Udo S and Alfred W: "Fatty Acids" in *Ullmann's Encyclopedia of Industrial Chemistry*. Wiley-VCH, Weinheim 2006.
38. Beare-Rogers J, Dieffenbacher A and Holm JV: *Lexicon of lipid nutrition (IUPAC Technical Report)*. *Pure and Applied Chemistry* 2001; 73(4): 685-44.

**How to cite this article:**

Yilmaz B and Celik I: Analysis of fatty acid compositions of apricot kernel oils for cosmetic purpose with gas chromatography-mass spectrometry method. *Int J Pharmacognosy* 2019; 6(10): 340-46. doi link: [http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.6\(10\).340-46](http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.6(10).340-46).

This Journal licensed under a Creative Commons Attribution-Non-commercial-Share Alike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)