



Received on 30 October 2019; received in revised form, 25 November 2019; accepted, 27 November 2019; published 30 November 2019

PHYTOCHEMICAL SCREENING, ANALYTICAL METHOD DEVELOPMENT AND VALIDATION FOR THE DETERMINATION OF CATECHINS IN *B. CILIATA* BY RP-HPLC

N. D. V. R. Saradhi^{*}, N. Madan Gopal, C. Madhusudhna Chetty and K. Sushmitha

Santhiram College of Pharmacy, NH - 40, Nerawada, Kurnool - 518112, Andhra Pradesh, India.

Keywords:

B. ciliata, RP- HPLC, Catechins, Phytochemical screening etc.

Correspondence to Author:

Dr. N. D. V. R. Saradhi

Associate Professor,
Department of Pharmaceutical
Analysis Santhiram College of
Pharmacy, NH - 40, Nerawada,
Kurnool -518112, Andhra Pradesh,
India.

E-mail: saradhi2u@gmail.com

ABSTRACT: Ayurveda an ancient system of medicines detailing a number of medicinal plants and their activities in humans or animals. The present research work aimed to develop an analytical procedure for the determination of catechins in the *B. ciliata*. It is famously known as a stone flower/stone breaker having biological activities like anti urolithiasis, antiviral, antidiabetic antitumor, and cardioprotective activity. The methanolic extract of the plant is isolated, and an analytical method is developed by using RP HPLC for the determination of catechins in the crude plant extract using a C₁₈ column (200 × 4.6 mm, 5 μ) and detected at 241 nm. The method is validated for its system suitability, linearity, accuracy, precision, robustness, and sensitivity as per the ICH guidelines Q2 (R1) to meet the analytical procedure in academic and industrial usage.

INTRODUCTION: *B. ciliata* is one of the medicinal plants discussed in the Ayurveda system of medicine. The plant even though old but having many activities of it, the usage of the plant leaves and roots is more. *B. ciliata* belongs to Saxifragaceae is a perennial herb. The catechins are the phytochemicals having antioxidants, antimicrobial activity, and used in kidney ailments. The earlier work through the light that only a few methods have been enlisted in the method development of catechins in various plants. There is no report on the study for the estimation of catechins in *B. ciliata* with the selected stationary and mobile phases. The fresh leaves and rhizomes were collected from the Nandyal, Andhra Pradesh, and its surroundings, which were authenticated from the senior botanist Dr. V. J. Sailaja Rani.

The plant authentication voucher number is KVSC/A/2019/P09. The plant parts were shade dried and extracted with methanol from the pure extract the RP HPLC method is developed for the estimation of catechins. The plant leaves and rhizomes are shown in **Fig. 1**.



FIG. 1: (A) LEAVES OF *B. CILIATA* (B) DRIED ROOTS OF *B. CILIATA*

MATERIALS AND METHODS: The chemicals used in the method were HPLC and AR grades, their percentage of purity between 99.98 - 100.02% (Chloroform, methanol, ethanol, acetone, distilled water). The pure form of the catechin was a gifted sample from the Hi Q Laboratory, Hyderabad.

QUICK RESPONSE CODE 	DOI: 10.13040/IJPSR.0975-8232.IJP.6(11).360-64
	The article can be accessed online on www.ijpjournal.com
DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.6(11).360-64	

The Soxhlet apparatus, volumetric flasks, beakers, round bottom flasks used for extraction and other procedures activities is borosil grade. The equipments are pre-calibrated electronic balance - Shimadzu (ATY 240), Ultrasonicator, Double Beam spectrophotometer – Shimadzu (UV- 1800), HPLC - Analytical technologies (2230) powered with N 2000 software, column – waters (B. No: 080606) – 200 × 4.6 mm, 5 μ particle size stationary Phase were used for the experiment.

Plant Extraction: The shade dried leaves were grounded to a fine size and sieved through no: 44, the powder was packed in a muslin cloth and adjusted to a position of $\frac{3}{4}$ the level of Soxhlet apparatus. In a 250 ml of the round bottom flask was added about 200 ml of methanol and made hot percolation continued for 3 h, cooled the flask and make the crude extract was concentrated into solid on a rotary evaporator. The flakes of the crude drug were collected and stored in a SS bottle.

Mobile Phase / Diluent, Standard, and Sample Solutions: An equal portion of the HPLC Grade methanol and Chloroform (50:50 v/v) into a reservoir bottle, standard is prepared by weighing and diluting the standard catechin which is filtered through 0.25 μ membrane filter and the resultant concentration of catechin becomes 3 ppm level, and approximate level of the sample also prepared.

Phytochemical Screening: General tests were performed for both standard and sample solutions for the identification of various phytochemicals. The test performed for carbohydrates, fats, proteins, alkaloids, glycosides, resins, and tannins. Both show positive results for tannins and further identification tests conducted for identification of tannins by bromine water test and match stick test, which showed and confirmed the presence of catechins in the plant extract. A TLC method is used for the separation of catechins from the crude plant extract. A pre-coated TLC silica gel 60 F 254 aluminum sheet is used as a stationary phase and methanol, water, chloroform (35:10:65 v/v) is a mobile phase. The sample and standard applied on the plate and eluted for 20 min.

The plate was dried and sprayed with 0.55% vanillin in a 4% hydrochloric acid. The spots were scrapped and dissolved in the methanol. Now, this

solution is measured for the absorbance to quantify the catechin present in the sample solution.

Measurement of Lambda Max: A spectrum plotted using diluent as blank to the sample, and standard solutions both are shown their lambda max at 241 nm. **Fig. 2** represented as:




FIG. 2A: STANDARD CATECHIN




FIG. 2B: EXTRACT OF *B. CILIATA* OVER STANDARD CATECHIN

HPLC Method: Various trails have been conducted by changing the column conditions, mobile phase concentrations, and UV detection wavelengths. The final optimized method developed by using waters C18 (200 × 4.6 mm, 5 μ) column with methanol, chloroform in 1:1 ratios, the flow rate maintained at 1ml/min. The chromatograms of the placebo, sample, and standard were shown in **Fig. 3, 4, 5**.




FIG. 3: BLANK / PLACEBO AT 241 nm




FIG. 4: STANDARD CATECHINS SHOWING PEAK AT 241 nm




FIG. 5: PLANT EXTRACT SHOWING PEAK AT 241 nm

Validation of the Method:

System Suitability: This procedure validates that the selected procedure for the estimation of the catechins in *B. ciliata* by using the HPLC conditions. A replicate injection of the same concentration of the catechin standard is injected into the system and measure the peak area, mean, SD % RSD was calculated. The results were shown in **Table 1**.

Linearity: Linearity or range is a measure of analytical procedure that the selected method is suitable for the least to maximum concentrations. The selected procedure is followed for beer lambert law.

From the standard primary stock solution, 1-5 ml was taken into an individual volumetric flask and diluted with diluent to get 1-5 ppm. The peak response was measured for the individual concentration. The correlation between the individual concentrations was shown in **Fig. 6**.




FIG. 6: LINEARITY

Precision: It is the agreement between individual test results with the procedure applied. The precision is completed in 3 steps repeatability, intermediate precision, and reproducibility. The procedure involves the replicate injection of the same concentration of the sample injected to the HPLC, the peak area was noted between the days, analysts, instruments, and laboratories, *etc.* the results were shown in **Table 1**.

Accuracy: It is a validated protocol for the measure the exactness of the experimental procedure. There are two ways for determining the accuracy compared to a reference method, recovery of the analyte spiked into a blank matrix and standard addition of the analyte. This can be done by taking 3 different concentrations of the sample and injected to the HPLC system, measure the peak area of each concentration, and calculate the percentage of recovery from each level of concentration. The results were shown in **Table 2**.

Robustness: it is a measure of the capacity of the developed method. The procedure involves the

change in the system conditions like column flow rate, temperature, detection wavelength, etc. the results were **Table 3**.

Sensitivity: The sensitivity of the experiment can be calculated by the LOD/LOQ of the validation parameters. The LOD is the lowest limit of quantity to be determined using the developed procedure. LOQ is the highest minimum limit of quantity to be determined by using the procedure.

They were theoretically calculated using linearity graph for its signal to noise ratio, and the results are shown in **Table 5**.

RESULT:

TABLE 1: PRECISION

Injection no.	Conc. in PPM	System suitability	Precision			
			Repeatability	Intermediate precision	Analyst variation	Instrument variation
1	3	61587	61587	61597	61483	60823
2	3	61498	67570	61591	61532	59878
3	3	61481	61581	61599	61345	61233
4	3	61594	61594	61590	61563	60384
5	3	61595	61595	61596	61423	61482
6	3	61497	61597	61587	61082	62383
Mean		61542	61587	61595	61404	61030
Standard Deviation		55.172	10.36	3.76	176.31	879.0
% RSD		0.089	0.01	0.006	0.287	1.440
ICH Criteria		>2	>2	>2	>2	>2
Test Result		Passes	Passes	Passes	Passes	Passes

TABLE 2: ACCURACY

S. no.	Accuracy level	Area	Spike added in mg	Recovered in mg	% Recovery
1	50%	31216	1.5	1.52	101.35
2		30682	1.5	1.49	99.62
3		30429	1.5	1.48	98.80
4	100%	60394	3.0	2.93	98.04
5		60297	3.0	2.96	97.88
6		62199	3.0	3.09	100.97
7	150%	91899	4.5	4.01	99.46
8		92385	4.5	4.52	99.99
9		92899	4.5	4.51	99.99

TABLE 3: ROBUSTNESS

Parameter	Mean retention time	Mean area	% RSD
Change in detection wave length (nm)			
237	2.09	61488.6	0.0095
241	2.09	61586.6	0.0080
245	2.09	61738.6	0.0195
Change in flow rate (ml/min)			
0.9	2.0736	55435	0.0075
1.0	2.0946	61594	0.0048
1.1	2.1254	67755	0.0055

TABLE 4: LOD AND LOQ:

Name of the drug	Slope	SD	LOD	LOQ
Catechins	20591	55.172	0.0088 µg/ml	0.0267 µg/ml
Acceptability			< 3.0	< 10

CONCLUSION: The method initiated with methanol extract, isolation, and phytochemical screening and thereby HPLC method development. The instruments and chemicals used for the purpose are calibrated and fine graded suitable for the analysis through HPLC.

The method developed is validated as per the ICH guidelines of Q2R1 for its system suitability, linearity, precision, accuracy, sensitivity, and robustness to meet the academic and industrial needs.

ACKNOWLEDGEMENT: Special thanks to Santhiram College of Pharmacy for providing all facilities in the laboratories for the new analytical method development for the estimation of catechins in *B. ciliata* by RP-HPLC method.

AUTHORS CONTRIBUTION: I am grateful to Dr. C. Madhusudhan Chetty in encountering the problems in relation to this work, and my supporter Mr. N. Madan Gopal played and continued the experiment at late hours of his duty. Students K. Sushmita has great in collecting the literature and extraction of the plant materials successfully.

CONFLICTS OF INTEREST: Nil

REFERENCES:

- Gurav SS: A comprehensive review: *Berginia ligulata* wall-A controversial clinical candidate. International Journal of Pharmaceutical Science and Research 2014; 5(5): 1630-42.
- Ruby KM, Dwivedi J and Chauhan R: *Berginia ciliata* mine of medicinal properties; a review. International Journal of Pharmaceutical Sciences 2012; 15(2): 20-23.
- Hagerman ANNE: Extraction of tannins from fresh and preserved leaves. J of Chemical Ecology 1998; 14(2).
- Gangwal A: Extraction and estimation and TLC of tannins: a review. International Journal of Pharmaceutical Sciences 2013; 2(3): 1585-88.
- Durgawale TP, Durgawale PP and Khanwelkar CC: Quantitative estimation of tannins by HPLC. Scholars Research Library 2016; 8(3): 123-26.
- Chavan U D and Amarowicz R: Effect of various solvent systems on extraction of phenolics tannins and sugars from beach pea. International Food Research Journal 2013; 20(3): 1139-44.
- Zohara SF, Meriem B, Samira S and Muneer MSA: Phytochemical screening and identification of some compounds from Mallow 2012; 2(4): 512-16.
- Srivastava N, Srivastava A and Srivastava S: Simultaneous quantification of berginin, epicatechin, (+) - catechin, and gallic in *Berginia ciliata* using HPLC. Journal of Liquid Chromatography and Related Data 2015; 38: 1207-12.
- Rawat AKS, Singh DP, Srivastava SK and Govindarajan R: HPLC determination of bergenin in different *Berginia* species. ACTA Chromatographia 2007; 19: 246-52.
- Jingyuanwen, Danhuili, Nataly and Marizlini: Development of an isocratic HPLC method for catechin quantification and its application to formulation studies, Elsevier 2012; 83: 1267-7.
- Felinger A, Boros B and Jakabova S: Validated HPLC method for simultaneously quantification of bergenin, arbutin, and gallic acid in leaves of different *Berginia* species. Springer Plus 2014; 77(17-18): 1129-35.
- Shankhan M, Khan W and Singh WAM: Berginin determination in different extracts by HPLC densitometry. Journal of Pharmacy and Bioallied Sciences 2015; 7(4): 272-74.
- Gupta SP and Gang G: Quantitative estimation of gallic acid and tannic acid in *Bhuvnesvara vati* by RP-HPLC. Scholars Research Library 2014; 6(2): 31-36.
- Madhavi GP, Vishal P and Rakesh KP: Development and Validation of Improved RP-HPLC method for identification and estimation of ellagic acid and gallic acid in triphala churna. International Journal of Chem Tech Research 2010; 2(3): 1486-93.
- Ganjage PJ, Tapkir AS and Nipanikar SU: Pawar, Development and validation of a RP-HPLC method for the determination of ellagic acid in *Terminalia bellirica* extract and single herb capsule of *Terminalia bellirica*. International Journal of Pharmaceutical Education and Research 2018; 52(4): S56-S62.
- Sanghavi N, Bhosale SD and Malode Y: RP-HPLC method development and validation of quercetin isolated from the plant *Tridax procumbence* L. Journal of Scientific and Innovative Research 2014; 3(6): 594-97.
- Klein T, Longhini R and Mello JCPD: Development of an analytical method using RP-HPLC-PDA for a semi purified extract of *Paullinia cupana* var. *Sorbilis*. Elsevier 2012; 502-06.

How to cite this article:

Saradhi NDVR, Gopal NM, Chetty CM and Sushmitha K: Phytochemical screening, analytical method development and validation for the determination of Catechins in *B. ciliata* by RP-HPLC. Int J Pharmacognosy 2019; 6(11): 360-64. doi link: [http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.6\(11\).360-64](http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.6(11).360-64).

This Journal licensed under a Creative Commons Attribution-Non-commercial-Share Alike 3.0 Unported License.

This article can be downloaded to **Android OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Play store)