



Received on 27 September 2021; received in revised form, 14 December 2021; accepted, 15 December 2021; published 31 December 2021

QUALITATIVE ASSAY OF ESSENTIAL OILS IN COMMERCIAL LAVENDER OIL BY GC-MS METHOD

Bilal Yilmaz^{*}, Burak Bayrak and Yucel Kadioglu

Department of Analytical Chemistry, Faculty of Pharmacy, Ataturk University, Erzurum, Turkey.

Keywords:

Lavender, Essential oils, GC-MS, Quality control

Correspondence to Author:

Dr. Bilal Yilmaz

Professor,
Department of Analytical Chemistry,
Faculty of Pharmacy, Ataturk
University, Erzurum, Turkey.

E-mail: yilmazb@atauni.edu.tr

ABSTRACT: Essential oils are largely employed for their therapeutic properties, being marketed extensively in pharmaceutical and cosmetic industry. Lavender oil is known for its excellent aroma. Therefore, it is extensively used in the cosmetic, perfumery, and flavour industries. The lavender oil is known to possess sedative, carminative, antidepressive, and anti-inflammatory properties. The aim of our study was to assess the purity and quality of lavender oils, available on the market from various commercial producers. Therefore, essential oil compositions of lavender oil were analyzed using gas chromatography-mass spectrometry (GC-MS). Chromatographic analyses were carried out on HP-5 MS column. The injector volume was 1 µl in split mode (40:1) and the carrier gas was helium at a flow rate of 0.8 ml/min. 33 essential chemical constituents were identified based on GC-MS in lavender oil supplied from an herbalist. The major components of the lavender oil were found linalyl acetate (37.82 %), linalool (33.07 %), eucalyptol (4.88 %), camphor (4.07 %), lavandulol acetate (1.53 %) and caryophyllene (1.43 %).

INTRODUCTION: Nowadays, essential oils as alternative therapies have gained worldwide concern, owing to their various biological activities. Considerable attention has been devoted to peppermint oil, which is widely used for its important properties, including antimicrobial, anti-inflammatory, antispasmodic, cytoprotective, hepatoprotective with strong antioxidant actions. It is also used for its cooling effect to enhance the dermal penetration of pharmaceuticals. Then, it is a standout amongst the most vital seasoning added substances in the World^{1, 2}. Lavender genus is a crucial member of the Lamiaceae family.

Lavandula species are generally appropriated in the Mediterranean district and developed in France, Italy and Spain. Also, lavender oil **Fig. 1** is widely used for its medicinal actions, including antispastic, anti-inflammatory, sedative, antimicrobial and general tonic action. Externally, it can be used for wounds and superficial burns. It contains as major compounds linalool (20 - 45%), linalyl acetate (25 - 46%), monoterpenes, alcohols, and esters³⁻⁵.



FIG. 1: LAVENDER OIL

QUICK RESPONSE CODE



DOI:

10.13040/IJPSR.0975-8232.IJP.8(12).502-05

Article can be accessed online on:
www.ijournal.com

DOI link: [http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.8\(12\).502-05](http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.8(12).502-05)

Even if essential oils are marketed extensively in the pharmaceutical and cosmetic industry, not all products available to use are properly controlled in terms of the quality of their composition. The analytical methods applied for the analysis of essential oils are numerous and include methods exploiting volatility or their optical activity, along with the IR spectral methods,

Refractometry and Thin Layer Chromatography⁶⁻⁸: Several methods have been reported to determine the main constituents of essential oils by GC-MS method⁹⁻¹⁵. Other techniques, inexpensive, remain a fast alternative, although there are limitations in terms of specificity and sensitivity. But, GC-MS is the elected technique for analyzing. To date, no GC-MS method is accounted for till date for assurance of essential oils by GC-MS in commercial lavender oil in Turkey.

Therefore, in this study, 33 essential chemical constituents were identified based on GC-MS in lavender oil supplied from an herbalist. The components were identified by comparing linear Kovats retention index, their retention times, and mass spectra with those obtained from the MS library.

MATERIALS AND METHODS:

Chemicals and Reagents: Linalyl acetate, linalool, eucalyptol, camphor, lavandulol acetate, caryophyllene, and different synthetic compounds were acquired from Sigma-Aldrich (St. Louis, MO, USA). Lavender oil was purchased by a local herbalist. It was labeled as a 100% natural product.

GC-MS System: GC-MS investigations were completed on an Agilent 7820A gas chromatography system equipped with 7673 series autosampler chemstation and 5977 series mass selective detector. HP-5 MS segment with 0.25 μm film thickness (30 m \times 0.25 mm I.D., USA) was utilized for separation. The temperatures of the inlet and transfer lines were 250 and 300 $^{\circ}\text{C}$, respectively.

GC-MS Conditions: Different temperature programs were researched for the technique. At the finish of this examination, the GC broiler temperature was at 60 $^{\circ}\text{C}$ for 10 min and afterward was customized to 220 $^{\circ}\text{C}$ at a rate of 4 $^{\circ}\text{C}/\text{min}$ and

kept in this temperature for 10 min. The stove temperature was at long last modified to 240 $^{\circ}\text{C}$ at a rate of 1 $^{\circ}\text{C}/\text{min}$ with the last hold time of 80 min. The split proportion was 40:1.

Identification of Components: The range of the obscure segment was contrasted, and the range of the part was put away in the National Institute of Standards and Technology Library Version (2005), Software, Turbomass 5.2. The parts were distinguished by contrasting direct Kovats maintenance list and mass spectra with those obtained from the MS library. Understanding on mass range GC-MS was directed utilizing the database on National Institute Standard and Technology having more than 62,000 examples. The relative rate measure of every part was determined by contrasting its normal pinnacle region with the complete areas. The name, atomic weight, and structure of the segments of the test materials were discovered.

RESULTS AND DISCUSSIONS:

Method Development and Optimization: The strategy advancement for the examination of phytocomponents depended on their synthetic properties. In this investigation, the fine segment covered with 5% phenyl and 95% dimethylpolysiloxane is a decent decision to partition these analytes since they elute as symmetrical tops at a wide scope of focuses. Distinctive temperature programs were explored for GC broiler.

At the finish of this examination, the best temperature program was chosen for good detachment. The GC broiler's temperature projects were as follows: beginning temperature of 60 $^{\circ}\text{C}$, held for 1. min, expanded to 220 $^{\circ}\text{C}$ at a rate of 4 $^{\circ}\text{C}/\text{min}$ held for 10 min. The broiler temperature was at long last modified to 240 $^{\circ}\text{C}$ at a rate of 1 $^{\circ}\text{C}/\text{min}$ with the last hold time of 80 min. Mass spectra were taken at 70 eV, and the mass range was from m/z 35 to 450.

GC-MS Analysis: GC-MS is a standout amongst the best procedures to distinguish the constituents of unstable issues, long-chain, stretched chain hydrocarbons, alcohols, amines, esters, ketones, heterocycles, and terpenes. GC-MS examination uncovered the nearness of 33 mixes in oil of lavender oil. Pinnacle number, maintenance time,

and compound name are expressed in **Table 1**. The substance 33 basic constituents were recognized dependent on GC-MS in lavender oil provided by the botanist. The significant parts of the lavender

oil were linalyl acetic acid derivation (37.82 %), linalool (33.07 %), eucalyptol (4.88 %), camphor (4.07 %), lavandulol acetic acid derivation (1.53 %) and caryophyllene (1.43 %).

TABLE 1: CHEMICAL COMPOSITION OF THE ESSENTIAL CONSTITUENTS OF LAVENDER OIL

Number	Retention time (min.)	Retention index	Name	% Ratio
1	7.67	937	α -pinene	0.18
2	8.47	952	camphene	0.22
3	10.06	974	sabinen	0.34
4	10.22	986	β -pinen	0.27
5	11.34	992	β -myrcene	0.24
6	12.05	994	amylethylcarbinol	0.17
7	13.64	1032	eucalyptol	0.32
8	14.89	1049	trans- β -acimene	1.98
9	16.23	1074	linalool oxide	0.11
10	16.91	1088	terpinolene	0.17
11	18.22	1099	linalool	33.07
12	19.71	1142	(-) camphor	4.07
13	21.02	1167	Endo borneol	2.08
14	21.43	1177	4-terpineol	0.11
15	21.70	1183	crypton	0.09
16	22.20	1190	L- α -terpineol	0.99
17	23.28	1226	bornyl formate	0.12
18	24.63	1257	linalyl acetate	37.82
19	25.57	1285	bornyl acetate	0.08
20	25.81	1306	lavandulol acetate	1.53
21	28.48	1364	nerol acetate	0.36
22	29.16	1382	geranyl acetate	0.78
23	29.32	1391	17-epi-sesquit hujene	0.12
24	30,39	1419	caryophyllene	1.43
25	30.86	1435	α -bergamotene	0.04
26	31.14	1464	sesquisabinene	0.03
27	31.57	1477	t- β -farnesene	0.97
28	32.45	1481	D-germacrene	0.38
29	33.22	1508	α -farnesene	0.34
30	33.54	1577	γ -muurolene	0.04
31	35.75	1581	caryophyllene oxide	0.11
32	37.58	1640	α -epicadinol	0.05
33	38.77	1684	α -bisabolol	0.39

On examination of the present outcomes with those detailed from different parts of India, it is very clear that the groupings of 1,8 cineole, camphor, β -caryophyllene, and caryophyllene oxide were somewhat higher, though the centralizations of α -terpineol, linalyl acetic acid derivation, geranyl acetic acid derivation, neryl acetic acid derivation and lavandulyl acetic acid derivation were generally lower in the Kashmir oil than in the present oil¹⁶. However, the lavender oil announced from the Kashmir valley contained a lot of limonene (11.0 %), citronellol (10.0 %) and α -terpineol (7.6 %) and low substance of linalool (10.0 %). Moreover, the centralizations of (E)- β -ocimene, 1-octen-3-yl acetic acid derivation, α -terpineol, and β -caryophyllene were marginally

higher in the oil delivered in Kodaikanal when contrasted with the present oil. These varieties could be because of contrasts in area, rise, hereditary makeup of the plant, or a versatile procedure to specific natural conditions. Lawrence additionally watched a wide variety in the quantitative organization of lavender oil contingent upon plant genotype and development zone, and the arrangement of the oil from lavenders were perceived to shift essentially as per height, microclimate, and area.

CONCLUSION: This investigation recommends that GC-MS is the viable technique for examining the fundamental oils. Applying the proposed system, a sum of 33 segments was distinguished for

the lavender fundamental oil. These parts represent 89.0% of the all-out relative substance of the basic lavender oil. This examination likewise exhibited that the basic lavender oil is rich in linalyl acetic acid derivation and linalool and could be a decent hotspot for this compound. The lavender fundamental oil may have immediate or roundabout mitigating or antinociceptive exercises. The effects of linalyl acetate and linalool in the future can be investigated.

ACKNOWLEDGEMENT: Nil

CONFLICT OF INTEREST: Nil

REFERENCES:

- Gutiérrez GLS, Reyes RC and Jaime HB: Medicinal plants for the treatment of nervios, anxiety and depression in mexican traditional medicine. *Revista Brasileira De Farmacognosia* 2014; 24: 591-608.
- Singh R, Shushni MAM and Belkheir A: Antibacterial and antioxidant activities of mentha piperita L. *Arabian Journal of Chemistry* 2015; 8: 322-328.
- Bajalan I and Pirbalouti AG: Variation in chemical composition of essential oil of populations of lavandula intermedia collected from Western Iran. *Industrial Crops and Product* 2015; 69: 344-347.
- Masoumeh BN, Seyed AS, Attieh N, Fatemeh E and Fahimeh-Sadat GK: The effects of aromatherapy with lavender essential oil on fatigue levels in haemodialysis patients: a randomized clinical trial. *Complementary Therapies in Clinical Practice* 2016; 22: 33-37.
- Silva GL, Luft C, Lunardelli A, Amaral RH, Melo DA, Donadio MV, Nunes FB, De Azambuja MS, Santana JC, Moraes CM, Mello RO, Cassel E, Pereira MA, De Oliveira JR: Antioxidant, analgesic and anti-inflammatory effects of lavender essential oil. *Anais Da Academia Brasileira De Ciencias* 2015; 87: 1397-1408.
- Huck WC: Advances of infrared spectroscopy in natural product research. *Phytochemis Letters* 2015; 11: 384-393.
- Lafhal S, Vanloot P, Bombarda I, Kister J and Dupuy N: Chemometric analysis of french lavender and lavandin essential oils by near infrared spectroscopy. *Industrial Crops and Products* 2016; 80: 156-164.
- Wozniak KS and Walasek M: Preparative separation of menthol and pulegone from peppermint oil (*Mentha Piperita* L.) by high-performance counter-current chromatography. *Phytochemistry Letters* 2014; 10: XCIV–XCVIII.
- Cioanca O, Mircea C, Hritcu L, Trifan A, Mihasan M, Aprotosoia AC, Robu S, Gille E and Hancianu M: In vitro-in vivo correlation of the antioxidant capacity of salviae aetheroleum essential oil. *Farmacia* 2015; (63-1): 34-39.
- Gaoa Q, Zhaoa X, Yina L, Zhanga Y, Wanga B, Wua X, Zhanga X, Fua X and Sund W: The essential oil of artemisia capillaris protects against CCl4. *Farmacognosia* 2016; 26: 369-374.
- Gavris I, Bodoki E, Verite P and Oprean R: Assessment of perfume ingredients with aphrodisiac potential by gas chromatography-mass spectrometry. *Farmacia* 2015; (63-5): 760-764.
- Li B, Zhang C, Peng L, Liang Z, Yan X, Zhu Y and Liu Y: Comparison of essential oil composition and phenolic acid content of selected salvia species measured by GC-MS and HPLC methods. *Industrial Crops and Products* 2015; 69: 329-334.
- Koch WK, Woźniak KS, Wska ES, Widelski J, Urjin O and Głowniak P: Comparative study on the volatiles composition of iris dichotoma pall. rhizome extracts. *Farmacia* 2014; (62-5): 874-881.
- Mehdi JH, Roudabeh SMP and Hassan S: Elimination of chromatographic and mass spectrometric problems in GC-MS analysis of lavender essential oil by multivariate curve resolution techniques: improving the peak purity assessment by variable size moving window-evolving factor analysis. *Journal of Chromatography B* 2015; 983: 83-89.
- Tyagi AK and Malik A: Antimicrobial potential and chemical composition of mentha piperita oil in liquid and vapour phase against food spoiling microorganisms. *Food Control* 2011; 22: 1707-1714.
- Tajuddin Shawl AS, Kumar S, Nigam MC and Hussain A: Production of lavender oil in Kashmir Valley. *Indian Perfumer* 1983; 27: 56-59.

How to cite this article:

Yilmaz B, Bayrak B and Kadioglu Y: Qualitative assay of essential oils in commercial lavender oil by GC-MS method. *Int J Pharmacognosy* 2021; 8(11): 502-05. doi link: [http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.8\(12\).502-05](http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.8(12).502-05).

This Journal licensed under a Creative Commons Attribution-Non-commercial-Share Alike 3.0 Unported License.

This article can be downloaded to **Android OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)