



Received on 11 July 2016; received in revised form, 18 August 2016; accepted, 27 August 2016; published 31 August 2016

IN-VITRO ANTHELMINTIC ACTIVITY OF METHANOLIC LEAF EXTRACT OF *ABRUS PRECATORIUS* LINN. (FABACEAE)

S. Selvadurai *, M. D. Dhana Raju, A. Venkateswara Rao and T. Shri Vijaya Kirubha

Department of Pharmacognosy, GIET School of Pharmacy, Rajahmundry - 533296, Andhra Pradesh, India.

Keywords:

Abrus precatorius Linn.
Anthelmintic activity, Piperazine citrate, methanolic extract, *Pheretima posthum*, *Tubifex tubifex*

Correspondence to Author:

S. Selvadurai

Assistant Professor,
Department of Pharmacognosy,
GIET School of Pharmacy,
Rajahmundry - 533296, Andhra Pradesh, India.

E-mail: sasdurai85@gmail.com

ABSTRACT: The present study was aimed to the investigation of the anthelmintic potential of crude methanolic extract of the leaves of *Abrus precatorius* Linn. on Indian earth-worm "*Pheretima posthum*" and "*Tubifex tubifex*." Three concentrations (5, 10, 15 mg/ml) of methanolic extract were studied in activity which involved the determination of the time of paralysis (vermifuge) and time of death (vermicidal) of the worms. Piperazine citrate (10 mg/ml) was used as reference standard drug whereas distilled water as the control. Determination of paralysis time and death time of the worms were recorded. The result shows that methanolic extract possesses vermifugal activity and found to be effective as an anthelmintic. Therefore, the anthelmintic activity of the methanolic extract of *Abrus precatorius* Linn. Leaves have been reported.

INTRODUCTION: Parasitic infection including Helminthiasis is a critical serious problem in the tropical regions including the Asian countries which affects more than two billions of people worldwide. Helminths produce the serious problem in human and other animals around the world specifically to the third world countries¹. Different type of helminths infects the human and animals out of which intestinal roundworms (*Ascaridia* sp.) are most common. The helminths parasites mainly subsist in the human body in the intestinal tract, but they are also found in tissue, as their larvae migrate towards them². Despite this prevalence of parasitic infections, the research on the anthelmintic drug is sparse.

According to the WHO, only a few drugs are used in the treatment of helminths in humans. Anthelmintics from natural sources could play a key role in the treatment of these parasite infections. Given this, attempts have been made to study the anthelmintic activity of traditional medicinal plants. For our studies, we used *Abrus precatorius* Linn. that belong to the Fabaceae family *Abrus precatorius* Linn., a highly medicinal plant, and commonly known as Indian Liquorice in English, *Gunchi*, *Ratti* in Hindi; Kannada-*Gulganji*; Malayalam-*Kumi*; Sanskrit- *Rektika*, *Gunja* and in Tamil as *Kuntumani*. on open lands.

Its aerial parts are used for treating certain health problems like leucorrhoea, gonorrhoea, diarrhea, and dysentery. The roots, leaves, and seeds are used for medicinal purpose. The roots and leaves are astringent, sweet, emetic, diuretic, anthelmintic and exoteric. They are useful in cough pharyngodynia, pectoralgia, inflammation, strangury and vitiated condition of vata.

	<p>DOI: 10.13040/IJPSR.0975-8232.IJP.3(8).371-74</p>
	<p>Article can be accessed online on: www.ijpjournal.com</p>
<p>DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.3(8).371-74</p>	

The seeds are acrid, bitter, astringent, purgative, toxic abortifacient, aphrodisiac, and trichogenous; seeds paste applied locally for sciatica, the stiffness of shoulder joints, paralysis, leucoderma, ulcers, skin disease, wounds, alopecia, asthma, tubercular glands, stomatitis, hyperdipsia and fever^{3,4}.

Scientific Classification:

Division : Angiospermae

Class : Magnoliopsidia

Order : Fabales

Family : Fabaceae

Genus : *Abrus*

Species : *Precatorius*

Macroscopy:

Synonym : Rosary pea, john crow bead

Biological Source : *Abrus precatorius*

Family : Fabaceae

Vernacular Names:

English : Crab's eye, Jequirity

Hindi : Rati, Gaunchi

Kannadam : Gulaganji

Malayalam : Kunni, Gundumani

Tamil : Gundumani, Kunthamani

Telugu : Guruvinda



FIG. 1: MACROSCOPY

MATERIALS AND METHODS:

Collection of Plant Material: The plant material *Abrus precatorius* Linn leaves were collected from a local area of Rajahmundry, East Godavari DT, AP. in April 2015 and authenticated by the renowned botanist Prof. J. Suneetha, Department of Botany, Government Arts College, Rajahmundry. A voucher specimen (GCR/01/2015) was deposited in the college herbarium.

Drugs and Chemicals: Piperazine citrate (Glaxo SmithKline), methanol, Distilled water were used during the experimental protocol. All the chemicals used are a laboratory and analytical grade.

Preparation of Plant Extract:⁵ The fresh leaves were dried under shade. The dried leaves were cut into small pieces and powdered in a hand mill. Then 300 gm of coarse powder was extracted with

900 ml of extra pure Methanol successively in a Soxhlet extractor repeatedly for 48 h. The extract was dried by solvent evaporation in a thermostat water bath at 50-60 °C temperature, and the extract was kept in desiccator for the experiment.

Preliminary Phytochemical Investigation ⁵: The percentage yield of Methanol extract obtained was calculated, and the extracts were subjected to a number of proximate qualitative phytochemical analysis. These procedures are already reported by a number of workers and used without any modification.

Selection of Experimental Model ⁶: Indian adult earthworms *Pheretima posthuma* and *Tubifex tubifex* were used to experiment. The earthworms were collected from the local supplier. Worms were washed with normal saline to remove all fecal matter. The earthworms of 7-12 cm in length and 0.4-0.5 cm in width were used for all the experimental protocol. Ready availability, anatomical and physiological resemblance of *Pheretima posthuma* and *Tubifex tubifex* made it be used initially for *in-vitro* evaluation of the anthelmintic activity.

Experimental Design ⁷⁻¹⁰: The anthelmintic activity was carried out as described by Ajaiyeoba EO. *et. al.*, 2001, with minor modifications. The Indian earthworm. The assay was performed on adult Indian earthworm *Pheretima posthuma*, due to its anatomical and physiological resemblance with the intestinal roundworm parasites of human beings.

Pheretima posthuma worms are easily available and used as a suitable model for screening of anthelmintic drug. The assay was also performed

on the aquarium worm, *Tubifex tubifex* because they belong to the same group of Annelida.

Briefly, 20 ml formulations containing three different concentrations, of crude methanolic extract of a leaf (5, 10 and 15 mg/ml in double distilled water) were prepared and six earthworms (same size) and approximately 5g of *Tubifex* worms were placed in it. Both the test solution and standard drug solution were freshly prepared and 'time for paralysis' was noted when no movement of any sort could be observed except when the worms were vigorously shaken. The 'time for death' of worms was recorded after ascertaining that the worms neither moved when shaken vigorously nor when dipped in warm water at 50°. A maximum time period of 120 min was ascertained for the paralyzing as well as death time of *Pheretima posthuma* and *Tubifex tubifex* worms. Piperazine citrate (10 mg/ml) was used as a reference standard with distilled water as the vehicle control. All experiments were repeated thrice.

RESULTS AND DISCUSSION: Preliminary phytochemical screening of crude extract of the leaves of *Abrus precatorius* L., revealed the presence of tannins, steroids, flavonoids, saponin, terpenoids etc. The presence of phytoconstituents is reported in **Table 1**. From the observations made, a dose-dependent paralytic effect much earlier and the time of death was observed **Table 2**. Although, methanolic extract showed anthelmintic activity in a dose-dependent manner, but the methanolic extract appeared to be more effective for both types of worms. Evaluation of anthelmintic activity was compared with reference standard piperazine citrate.

TABLE 1: DATA SHOWING THE PRELIMINARY PHYTOCHEMICAL SCREENING OF ABRUS PRECATORIUS L

Phytoconstituents	Petroleum ether extract	Chloroform extract	Acetone extract	Methanolic extract	Aqueous extract
Alkaloids	(±)	(±)	(±)	(±)	(-)
Carbohydrates	(-)	(±)	(±)	(±)	(±)
Glycosides	(-)	(-)	(-)	(±)	(±)
Flavonoids	(-)	(-)	(-)	(±)	(±)
Phytosterols	(-)	(-)	(-)	(-)	(-)
Fixed oils and Fats	(-)	(-)	(-)	(-)	(-)
Saponins	(-)	(-)	(-)	(-)	(±)
Phenolic compounds and Tannins	(±)	(-)	(±)	(±)	(±)
Lignins	(-)	(-)	(-)	(-)	(-)
Proteins and Free Amino acids	(-)	(-)	(-)	(-)	(±)
Gums and Mucilage	(-)	(-)	(-)	(-)	(-)

TABLE 2: ANTHELMINTIC ACTIVITY OF METHANOLIC EXTRACT OF *ABRUS PRECATORIUS* L.

Groups	Conc. mg/ml	<i>Pheretima posthuma</i>		<i>Tubifex tubifex</i>	
		Paralyzing time (min)	Death time (min)	Paralyzing time (min)	Death time (min)
Distilled water	-	-	-	-	-
Leaf extract (Methanolic)	5	60.66±0.67*	80.66 ± 0.67*	63.00±2.08*	75.33±1.45*
	10	36.32±0.88*	66.33±0.88*	31.33±0.67*	39.33±0.88*
	15	22.33±0.88**	45.00±0.58*	14.66±0.88*	20.66±1.33*
Piperazine citrate	10	30.6±0.06*	51±0.006*	25.00±1.16*	64±0.88*
	15	20.13±0.98*	40.19±0.15*	19.01±1.11*	50.07±1.19*

Results are expressed as mean±sem N=3; *P<0.05 as compared to standards.

The methanolic extract of the leaves of *abrus precatorius* L. caused paralysis at 22.33 min. and time of death at 45.00 min. for *Pheretima posthuma* and 14.66 min as paralysis time and 20.66 min as death time for *Tubifex tubifex* worms respectively. The reference drug piperazine citrate showed the time of disease and time of death as 25.00 and 64.00 min, respectively. Considering the methanolic extract of leaf showed comparable activity, it would be important to identify the key phytoconstituents. The mean and SEM were analyzed statistically by ANOVA followed by Dunnett's test, $P<0.05$ is considered as significant.

CONCLUSION: The *Abrus precatorius* L. leaf extracts have shown significant anthelmintic activity at all the tested doses when compared to control as vermifuge and vermucidal while highest activity exhibited by the higher conc. (15 mg/ml) which assures the ethnomedicinal claim. Hence, we can think about this herb as an alternate source of anthelmintic drugs and also can generate a new active lead for the suitable anthelmintic drug.

ACKNOWLEDGEMENT: Nil

CONFLICT OF INTEREST: Nil

How to cite this article:

Selvadurai S, Dhana Raju MD, Rao AV and Kirubha TSV: *In-vitro* anthelmintic activity of methanolic leaf extract of *Abrus precatorius* Linn. (Fabaceae). Int J Pharmacognosy 2016; 3(8): 371-74. doi: 10.13040/IJPSR.0975-8232.3(8).371-74.

This Journal licensed under a Creative Commons Attribution-Non-commercial-Share Alike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)

REFERENCES:

- Hossain E, Chandra G, Nandy AP, Mandal SC and Gupta JK: Anthelmintic effect of a methanol extract of leaves of *Dregea volubilis* on *Paramphistomum explanatum*. Parasitol Res 2012; 110: 809-814.
- Tripathi KD. Essentials of Medical Pharmacology. 6th ed. New Delhi: Jaypee Brothers Medical Publishers (P) Ltd. 2008: 808-810.
- Fernando R: National Poisons Information Centre, Written a review about *Abrus precatorius*, Faculty of Medicine, Kynsey Road, Colombo 8, Sri Lanka.
- Karitikaran KR and Basu BD: Indian medicinal plants, I, 711.
- Kokate CK, Purohit AP and Gokhale SB: Text Book of Pharmacognosy. 41st ed. Pune: Nirali Prakashan; 2008.
- Chatterjee KD: Parasitology, Protozoology and Helminthology. Calcutta: Guha Ray Sree Saraswaty Pres Ltd 1967: 168-169.
- Ajaiyeoba EO, Onocha PA and Olarenwaju OT: *In-vitro* anthelmintic properties of *Buchholzia coriacea* and *Gynandropsis gynandra* extract. Pharm Bio 2001; 39: 217-220.
- Kumar T, Alexander A, Dewangan D and Nagori K.: Anthelmintic activity of the whole plant of *Bauhinia purpurea* (linn.). Asian J Pharm Clin Res 2011; 4(3): 110-111.
- Kosalge SB and Fursule RA: Investigation of *in-vitro* Anthelmintic activity of *Thespesia Lampas* (Cav.). Asian J Pharm Clin Res 2009; 2(2): 69-71.
- Chandan HS, Tapas AR and Sakarkar DM: Anthelmintic activity of extracts of *Coriandrum sativum* Linn. in Indian earthworm. Int J Phytomedicine 2011; 3: 36-40.