



Received on 05 November 2013; received in revised form, 06 December 2013; accepted, 27 December 2013; published 01 January 2014

CHEMICAL COMPOSITION AND ANTI-INFLAMMATORY ACTIVITY OF ESSENTIAL OILS FROM THE LEAVES OF *OCIMUM BASILICUM* L. AND *OCIMUM GRATISSIMUM* L. (LAMIACEAE)

B. C. Okoye-Festus^{* 1}, O. Obonga Willfred², A. Onyegbule Felix¹, I. Okoro Ogheneogaga², Okokpedi Peace², I. Nwosu Ngozi², Ajibo Jane² and O. Ndu Okechukwu³

Department of Pharmaceutical and Medicinal Chemistry¹, Faculty of Pharmaceutical Sciences, Nnamdi Azikiwe University, Awka, Anambra State, Nigeria.

Department of Pharmaceutical and Medicinal Chemistry², Faculty of Pharmaceutical Sciences, University of Nigeria, Nsukka, 410001, Enugu state, Nigeria.

Department of Pharmacology and Toxicology³, Faculty of Pharmaceutical Sciences, University of Nigeria, Nsukka, 410001, Enugu state, Nigeria.

Keywords:

Ocimum basilicum,
Ocimum gratissimum, Essential oils,
Topical anti-inflammatory,
Xylene-induced ear edema,
GC-MS

Correspondence to Author: Okoye Festus BC

Department of Pharmaceutical and Medicinal Chemistry, Faculty of Pharmaceutical Sciences, Nnamdi Azikiwe University, Awka, Anambra State, Nigeria.

E-mail: basdenc@yahoo.com

ABSTRACT: *Ocimum basilicum* L. and *Ocimum gratissimum* L. (Lamiaceae) are widely distributed aromatic herbs used in ethnomedicinal management of a range of inflammatory disorders. In the present work, we evaluated the topical anti-inflammatory effects of the volatile constituents extracted from the fresh leaves of these plant species. Fresh leaves of the plants were subjected to hydro distillation to obtain the volatile oils OBV and OGV from *O. basilicum* and *O. gratissimum* respectively. The fresh leaves were also extracted with n-hexane to obtain OBHE and OGHE respectively. OBV, OGV, OBHE, and OGHE were screened for anti-inflammatory effect using xylene-induced ear edema as a model of inflammation. Their chemical constituents were also analyzed using GC/MS apparatus. At 50 µg/ear OBV, OGV, OBHE and OGHE exhibited significant (P<0.05) topical anti-inflammatory effect with edema inhibitions of 50.0, 63.3, 62.7 and 80 % respectively. The effects were comparable (P<0.05) with that of 100 µg/ear hydrocortisone (% edema inhibition of 54.8). 11 of the compounds from OGV are monoterpenes while 4 are sesquiterpenes. 8 of the compounds identified in OGHE are oxygenated monoterpene derivatives, 5 are sesquiterpenes, and the others are long chain carboxylic acid, eugenol and phthalate derivatives. OGV and OGHE contain linalool, 1-terpinen-4-ol, alpha-caryophyllene and trans-longipinocarveol in common. Compounds identified in OBV include 2 monoterpenes, 7 oxygenated monoterpene derivatives, 2 sesquiterpenes, a long chain monocarboxylic acid and a triterpene, alpha-amyrin. OBV and OBHE contain eugenol acetate in common. Some of these identified volatile constituents may be contributing to the observed anti-inflammatory effects.

INTRODUCTION: *Ocimum basilicum* L. and *Ocimum gratissimum* L. (Lamiaceae) are widely distributed aromatic herbs¹.

Ocimum basilicum (Sweet basil) is an erect herb that grows to a height of 90 cm. The stem is purplish, and the leaves are lanceolate, glossy, and fragrant. The flowers are tubular, bilabiate, purplish, and packed in whorled racemes. *Ocimum gratissimum* (Wild basil) is also a perennial herb with erect stem and can grow up to 1-3 m tall. The stem is round glandular, much branched, glabrous or pubescent and woody at the base. The leaves are opposite, slender and the left blade elliptical to



ovate². The leaves of *Ocimum basilicum* have been used in ethnomedicine for a variety of ailments ranging from respiratory disorders³, fever⁴, as a remedy for gonorrhoea, catarrh conditions, cough, constipation, dysentery, ringworm, carminative and hypertension^{4,5}. The whole plant and the essential oil of *Ocimum gratissimum* have many applications in traditional medicine, especially in Africa and India. Preparations from the whole plant are used as stomachic and in treating sunstroke, headache, and influenza. The seeds have laxative properties and are prescribed against gonorrhoea. The essential oil is applied against fever, inflammations of the throat, ears or eyes, stomach pain, diarrhea and skin diseases^{1,2,6,7} and previous studies have confirmed its hypoglycaemic effect⁸.

Although there are reported studies on the anti-inflammatory effect of some species of *Ocimum*⁹⁻¹¹, there is to date no known scientific study validating the ethnomedicinal use of *Ocimum basilicum* and *Ocimum gratissimum* in inflammatory conditions. Inflammatory mycoses, a clinical condition associated with inflammation, fungal and bacterial infection is commonly treated with a combination of the anti-inflammatory, antifungal and antibacterial agent in one formulation¹²⁻¹⁵.

There are several reported studies on the antimicrobial potentials of *Ocimum* species¹⁶⁻²⁷. Our aim in the present study is thus to evaluate the topical anti-inflammatory potentials of the two *Ocimum* spp with the view of ascertaining their suitability for use in topical herbal formulations for inflammatory mycoses. We have also adopted two methods of essential oil extraction namely hydrodistillation and solvent extraction.

MATERIALS AND METHODS

Plant Material: Fresh leaves of *Ocimum basilicum* and *Ocimum gratissimum* were purchased from a local market in Nsukka, Enugu State Nigeria. The leaves were chopped into smaller pieces and used immediately for essential oil extraction.

Extraction of the Essential Oils: About 600 g fresh leaves of *Ocimum basilicum* and *Ocimum gratissimum* were coarsely milled and the volatile constituents isolated by hydro-distillation for 4 h using Clevenger apparatus. The volatile oil

fractions OBV and OGV respectively were recovered. About 200 g each of coarsely milled fresh leaves of *Ocimum basilicum* and *Ocimum gratissimum* were extracted in 500 mL hexane for 48 h with constant checking at room temperature. The hexane extracts were concentrated *in vacuo* to yield the volatile oil fractions OBHE and OGHE respectively. OBV, OGV, OBHE, and OGHE were stored in the refrigerator (0 °C) before topical anti-inflammatory tests and analysis of the constituents with GC/MS apparatus.

Gas Chromatography-Mass Spectrometry:

OBV, OGV, OBHE, and OGHE were analyzed by GC/MS (GCMS-QP2010 PLUS, Shimadzu Japan). Operating conditions were as follows: carrier gas, helium with a flow rate of 2 ml/min; column temperature, 60-280 °C at 5 °C/min; injector and detector temperatures, 250 °C; injected volume 2 µl; split ratio, 1:50. The MS operating parameters were as follows: ionization potential, 70 eV; ionization current, 1A; ion source temperature, 200 °C and resolution of 1000.

Identification of Compounds: Identification of components in OBV, OGV, OBHE, and OGHE were based on a comparison of the retention indices and computer matching of MS fragments with the NIST05.LIB.

Topical Anti-inflammatory Tests: The effect of OBV, OGV, OBHE, and OGHE on acute topical edema was evaluated by a modification of previously reported methods^{28, 29}. Adult albino mice (20 ± 5g) of either sex were divided into groups of 5 animals. The treatment groups received the extracts dissolved in xylene at doses of 50, 100, 200 and 400 µg/ear applied on the anterior surface of the right ear. Control animals received either equivalent volume of the phlogistic agent (xylene) or hydrocortisone dissolved in xylene (100 µg/ear). Two hours after application, the mice were sacrificed and both ears removed.

Circular sections (5mm) of both the right (treated) and left (untreated) ears were punched out using a cork borer and weighed. Edema was quantified as weight differences between the two earplugs. The anti-inflammatory activity was evaluated as percent edema inhibition in the treated animals relative to the control animals^{28, 29} using the relation:

$$\% \text{ Edema Reduction} = \left[1 - \frac{R_t - L_t}{R_c - L_c} \right] \times 100$$

Where R_t = mean weight of right earplug of treated animals; L_t = mean weight of left earplug of treated animals; R_c = mean weight of the right earplug of control (vehicle treated) animals; L_c = mean weight of the left earplug of control (vehicle treated) animals.

Statistical Analysis: Results of the anti-inflammatory effect obtained were analyzed by SPSS version 11 using one way ANOVA and subjected to Fischer LSD post hoc tests and expressed as mean \pm SEM. Differences between means were considered significant at $P < 0.05$.

RESULTS:

Anti-inflammatory Activity: The result of the topical anti-inflammatory screening of the volatile oils is shown in **Table 1** and **2**. At the dose of 50 $\mu\text{g}/\text{ear}$, all the tested oils exhibited significant ($p < 0.05$) inhibition of edema induced by topical application of hexane. OGHE showed better anti-inflammatory effect than OGV at the tested doses, while OBV showed better anti-inflammatory effect than OBHE. When compared with hydrocortisone, OGHE and OBV exhibited the better anti-inflammatory effect. The anti-inflammatory effect exhibited by the oils is however not dose-dependent.

TABLE 1: EFFECT OF THE VOLATILE OILS FROM *OCIMUM GRATISSIMUM* LEAVES ON XYLENE-INDUCED EAR EDEMA IN MICE

Test Material	Dose ($\mu\text{g}/\text{ear}$)	Mean ear edema (Mean \pm SEM) at 2 h	Inhibition (%) at 2 h	Inhibition vs. Hydrocortisone (%)
OGV	50	2.68 \pm 0.81*	52.57	126.40
	100	3.68 \pm 0.56	34.87	83.84
	200	3.64 \pm 1.10*	35.57	85.52
	400	3.66 \pm 0.34*	35.22	84.68
OGHE	50	0.80 \pm 0.37**	85.84	206.40
	100	2.40 \pm 0.37*	57.52	138.30
	200	1.40 \pm 0.40*	75.22	180.86
	400	2.60 \pm 0.68*	53.98	129.79
Hydroc	100	3.30 \pm 0.00*	41.59	100
Xylene	0.005 ml	5.65 \pm 0.91	-	-

OGV = Volatile oil from *Ocimum gratissimum* leaves obtained by hydrodistillation

OGHE = Volatile oil from *Ocimum gratissimum* leaves obtained by solvent (hexane) extraction.

The test materials were applied topically at the stated doses. Treatment animals were compared to control animals which had received vehicle only or prednisolone;

* $P < 0.05$, ** $P < 0.01$, $n = 5$.

TABLE 2: EFFECT OF THE VOLATILE OILS FROM *OCIMUM BASILICUM* LEAVES ON XYLENE-INDUCED EAR EDEMA IN MICE

Test Material	Dose ($\mu\text{g}/\text{ear}$)	Mean ear edema (Mean \pm SEM) at 2 h	Inhibition (%) at 2 h	Inhibition vs. Hydrocortisone (%)
OBV	50	2.00 \pm 0.45*	64.60	155.32
	100	2.40 \pm 0.75*	57.52	138.30
	200	2.40 \pm 0.51*	57.52	138.30
	400	1.60 \pm 0.40**	71.68	172.35
OBHE	50	2.72 \pm 0.60**	51.86	124.69
	100	4.72 \pm 0.81	16.46	39.58
	200	4.72 \pm 0.81*	16.46	39.58
	400	3.40 \pm 0.00*	39.82	95.74
Hydroc	100	3.30 \pm 0.00*	41.59	100
Xylene	0.005 ml	5.65 \pm 0.91	-	-

OBV = Volatile oil from *Ocimum basilicum* leaves obtained by hydrodistillation

OBHE = Volatile oil from *Ocimum basilicum* leaves obtained by solvent (hexane) extraction

The test materials were applied topically at the stated doses. Treatment animals were compared to control animals which had received vehicle only or prednisolone.

* $P < 0.05$, ** $P < 0.01$, $n = 5$

Volatile Oil Compositions of the Leaves: The results of the chemical composition of the volatile oils extracted from the two species of *Ocimum* by solvent extraction and hydrodistillation are shown in **Table 3** and **4**. A total of 15 compounds were identified in OGV as against 18 compounds identified in OGHE. 11 of the compounds from OGV are monoterpenes while 4 are sesquiterpenes. 8 of the compounds isolated from OGHE are oxygenated monoterpene derivatives, 5 are sesquiterpenes, and the others are long chain carboxylic acid, eugenol and phthalate derivatives. OGV and OGHE contain linalool, 1-Terpinen-4-ol, alpha-caryophyllene and trans longipinocarveol in common. A total of 14 and 11 compounds were identified in OBV and OBHE respectively. Compounds identified in OBV include 2 monoterpenes, 7 oxygenated monoterpene derivatives, 2 sesquiterpenes, a long chain monocarboxylic acid and a triterpene, alpha-amyrin. OBV and OBHE contain eugenol acetate in common.

TABLE 3: CHEMICAL COMPOSITION OF THE VOLATILE OILS EXTRACTED FROM *OCIMUM GRATISSIMUM* AND LEAVES

S. no.	A compound identified in OGV	A compound identified in OGHE
1	Alpha Thujene	Linalool
2	Camphene	Hexyl butanoate
3	Beta-Myrlene	Borneol (1,8 cineole)
4	Thujanol	1-Terpinen-4-ol
5	D-Fenchone	Thymoquinone
6	Linalool	O-Tertbutylphenol
7	gamma-Terpene	Durenol
8	Thujone	Verbenone
9	Beta Thujene	Caryophyllene
10		Methyl 9-oxo nonanoate
11	1-Terpinen-4-ol	Alpha caryophyllene
12		Eudesma-4(14), 11-diene
13	Thujene-2-one	(-) Alpha Pangasinan
14	Alpha caryophyllene	Alpha Cardinol
15	(-) Alpha Panasinsen	Trans Longipinocarveol
16	naphthalene	n-Hexadecanoic acid
17	azulene	Dehydrodihydrodiisoeugenol
18		Di-n-octyl phthalate

OGV = Volatile oil from *Ocimum gratissimum* leaves obtained by hydrodistillation

OGHE = Volatile oil from *Ocimum gratissimum* leaves obtained by solvent (hexane) extraction

TABLE 4: CHEMICAL COMPOSITION OF THE ESSENTIAL OIL ISOLATED FROM *OCIMUM BASILICUM* LEAVES

S. no.	A compound identified in OBV	A compound identified in OBHE
1	Eucalyptol	2-Methyl benzyl alcohol
2	cis-beta-Ocemene	Thymol
3	gamma-Terpene	Eugenol acetate
4	D-Fenchone	Eugenol methyl ether
5	Linalool	Cubanol
6	D-2-Bornanone	Hexadecanoic acid
7	L-Terpinen-4-ol	13-Hexyloxacyclotridec-10-en-2-one
8	Eugenol acetate	8-(2-octylcyclopropyl) octanal
9	Borneol (1,8 cineole) acetate	Linoleic acid
10	Alpha Bergamot	Di-n-octyl phthalate
11	Germacrene	2-methyl Benzaldehyde
12	Dotracotane	
13	Hexadecanoic acid	
14	Alpha-Amyrin	

OBV = Volatile oil from *Ocimum basilicum* leaves obtained by hydrodistillation

OBHE = Volatile oil from *Ocimum basilicum* leaves obtained by solvent (hexane) extraction

DISCUSSION: The whole leaves and essential oil of various species of *Ocimum* have several applications in traditional medicines, especially in Africa and India. Previous studies have validated the hypoglycaemic effect⁸, antibacterial²¹⁻²⁷, antinociceptive and anti-inflammatory^{10, 11} properties of some *Ocimum* species. Our present studies have shown that the essential oils from *Ocimum gratissimum* and *Ocimum basilicum* possess significant topical anti-inflammatory effect in xylene-induced mouse ear edema model. The anti-inflammatory activity was found to vary based on the method of essential oil extraction, which has a direct bearing on the chemical compositions. In general, the activity increases with increase in the proportion of some essential oils like eugenol, linalool, D-fenchone, 1-terpene-4-ol, thymol, alpha-caryophyllene and the presence of diterpenes and triterpenes, which have earlier been reported to possess anti-inflammatory activity³⁰⁻³⁵.

Essential oils obtained from *Ocimum gatissimum* by solvent (n-hexane) extraction (OGHE) showed

better topical anti-inflammatory activity than the oils obtained from the same plant species by hydro-distillation (OGV). Conversely, the essential oils obtained from *Ocimum basilicum* by hydro-distillation (OBV) showed better anti-inflammatory activity than the oils obtained from the same plant species by solvent (n-hexane) extraction (OBHE). These observations can be correlated with the chemical compositions of the essential oils from both plant species isolated by the two reported methods **Table 3** and **4**. OBV and OBHE both contained eugenol derivatives previously reported to possess significant anti-inflammatory activity^{36, 37}. OBV, however, also contained several other compounds like eucalyptol, linalool, borneol acetate, alpha-bergamot, germacrene and a triterpenoid alpha-amyrin **Table 4**. These compounds have been reported to exhibit anti-inflammatory activity³⁸⁻⁴⁰ or detected in essential oils reported exhibiting anti-inflammatory activity^{41, 42}. This might explain the significantly higher anti-inflammatory activity of OBV compared to OBHE. OGV and OGHE both contained linalool, alpha-caryophyllene, (-) alpha Pangasinan and 1-terpinene-4-ol along with other volatile constituents like alpha-thujone, camphene, borneol, verbenone, D-Fenchone, alpha cardinal, etc. **Table 3**. These volatile constituents have been detected in some essential oil isolated from plants with anti-inflammatory activity^{41, 42}.

The presence of a eugenol derivative (Dehydrodihydrodiisoeugenol), a diterpene (Eudesma-4(14), 11-diene) and Di-n-octyl phthalate in OGHE might be the plausible explanation for the observed better anti-inflammatory activity that OGV. It is also an interesting observation that OGHE and OBV (at 50 µg/ear) exhibited close to 2 times topical anti-inflammatory activity compared to indomethacin (100 µg/ear) **Table 3** and **4**. This observation thus supports the use of volatile oils obtained by solvent (n-hexane) extraction from *Ocimum gratissimum* and the oils obtained by hydro distillation from *Ocimum basilicum* for topical application in inflammatory conditions.

Previous studies have indicated the chemical variability in the composition of the essential oil extracted from *Ocimum* species. At least six chemotypes namely eugenol, thymol, citral, ethyl

cinnamate, geraniol, and linalool have been identified, with eugenol being the most important economically². In a recent study, phenylpropanoids were shown to be the major constituents of *Ocimum gratissimum* and *Ocimum basilicum*⁴³.

Specifically, in the study, methyl chavicol and linalool were shown to be the major constituent of *Ocimum basilicum* along with bicyclogermacrene and alpha-terpineol. Also, in the study the major constituents in *Ocimum gratissimum* were eugenol,^{1, 8} cineole, germacrene D and beta-caryophyllene. Our current results show somewhat a slightly different variability. Linalool, germacrene, alpha-terpineol were also shown present in *Ocimum basilicum* along with eugenol derivatives, while caryophyllene and^{1, 8} cineole were shown present in *Ocimum gratissimum* along with linalool and alpha-terpineol.

Our results so far have shown that the volatile oils isolated from *Ocimum basilicum* and *Ocimum gratissimum* show varying degrees of topical anti-inflammatory activity. This supports the ethnomedicinal uses of these plant materials in the management of inflammatory disease states. These plant materials have also been reported to possess significant antifungal and antibacterial activities. Our work thus supports their use as topical agents in the management of inflammatory mycoses.

ACKNOWLEDGEMENT: The authors are grateful to National Research Institute for Chemical Technology, Zaria, Kaduna State, Nigeria for the measurement of GC/MS spectra.

CONFLICT OF INTEREST: Nil

REFERENCES:

1. Sulistiarini D: *Ocimum gratissimum* L. In: Plant Resources of South-East Asia. No. 19: Essential-oils plants. (Oyen, LPA & Nguyen X D, Eds). Prosea Foundation, Bogor, Indonesia 1999: 140-142.
2. Orwa C, Mutua A, Kindt R, Jamnadass R and Anthony S: Agroforestry Database: a tree reference and selection guide version 4.0. World Agroforestry Centre, Kenya 2009. <http://www.worldagroforestry.org/sites/treedbs/treedatabases.asp> (accessed Dec 2011).
3. Okujagu TF, Etatuvi SO, Ifeyinwa E, Jimoh B and Nwokeke. Book of abstract of published Research finding on Nigerian Medicinal plant and traditional medicine practice 2005; 1: 90.

4. Idu M, Osawanu M and Orhue E: Medicinal plants in some local markets in Benin City, Nigeria. *Ethnobotany* 2005; 17: 118-122.
5. Odugbemi T and Akinsulire O: Medicinal plants by species names. In: *Outlines and pictures of medicinal plants from Nigeria* (Odugbemi T, ed). University of Lagos Press, Lagos 2006: 73- 116.
6. Rabelo M, Souza EP and Soares PM: Antinociceptive properties of the essential oil of *Ocimum gratissimum* L. (Labiatae) in mice. *Braz J Med Biol Res* 2003; 36: 521–524.
7. Orafidiya LO, Agbani EO and Iwalewa EO: Studies on the acute and sub-chronic toxicity of the essential oil of *Ocimum gratissimum* L. leaf. *Phytomedicine* 2004; 11: 71-76.
8. Aguiyi JC, Obi CI, Gang SS and Igwe AC: Hypoglycaemic activity of *Ocimum gratissimum* in rats. *Fitoterapia* 2000; 71(4): 444-446.
9. Singh S, Majumdar DK and Rehan HMS: Evaluation of anti-inflammatory potential of fixed oil of *Ocimum sanctum* (Holy basil) and its possible mechanism of action. *J Ethnopharmacol* 1996; 54: 19-26.
10. Lino CS, Gomes PB and Lucetti DL: Evaluation of antinociceptive and antiinflammatory activities of the essential oil (EO) of *Ocimum micranthum* Willd. from Northeastern Brazil. *Phytother Res* 2005; 19: 708-712.
11. Thakur K and Pitre KS: Anti-inflammatory activity of extracted eugenol from *Ocimum sanctum* leaves. *Rasayan J Chem* 2009; 2(2): 472-474
12. Hailu T, Endris M, Kaleab A and Tsige GM: Antimicrobial activities of some selected traditional ethiopian medicinal plants used in the treatment of skin disorders. *J Ethnopharmacol* 2005; 100 (1-2): 168-175.
13. Reddy NBB, Reddy VNRK, Torgalkar A and Murugan NR: Herbal composition for the regression of chronic inflammatory skin disorders. U.S. Patent 7,666,450. February 02, 2010.
14. Solanki R: Treatment of skin diseases through medicinal plants in different regions of the world. *Int J Biomed Res* 2011; 2(1): 73-80
15. Meisner and Lorrane F: Topical composition for the treatment of psoriasis and related skin disorders. U.S. Patents 7,670,620. May 02, 2010.
16. Koba K, Poutouli PW, Christine RC and Sanda K: Antifungal activity of essential oils from *Ocimum gratissimum* L. grown in Togo. *J Sci Res* 2009; 1: 164-171.
17. Suppakul P, Miltz J and Sonneveld K: Antimicrobial properties of basil and its possible application in food packaging. *J Agric Food Chem* 2003; 51: 3197–3207.
18. Nakamura CV, Nakamura T, Bando E, Melo AFN, Cortez DAG and Filho BPD: Antibacterial activities of *Ocimum gratissimum* L. essential oil. *Mem Inst Oswaldo Cruz* 1999; 94(5): 675-678.
19. Nakamura CV, Ishid K and Faccin LC: *In-vitro* activities of essential oils from *Ocimum gratissimum* against four *Candida* species. *Research in Microbiology* 2004; 155(7): 579-586.
20. Anadi AK, Monan M, Haider SZ and Sharma A: Essential oil composition and antimicrobial activity of three *Ocimum* species from Uttarakhand (India). *Int J Pharm Sci* 2011; 3(S 3): 223-225
21. Silva MRM, Oliveira JG and Fernandes OFL: Antifungal activities of *Ocimum gratissimum* towards dermatophytes. *J Ethnopharmacol* 2005; 48(1): 72-86.
22. Lemos JA, Passos XS and Fernandes OF: Antifungal activity of *Ocimum gratissimum* towards *Cryptococcus neoformans*. *J Ethnopharmacol* 2005; 20: 4-13
23. Lopez P, Sanchez K, Battle R and Nerin C: Solid and vapour phase anti-microbial activities of six essential oils susceptibility of selected foodborne bacterial and fungal strains. *J Agric Food Chem* 2005; 53(17): 6939-6946.
24. Avancini C, Wiest JM and Dall'agnol R: Antimicrobial activity of plants used in the prevention and control of bovine mastitis in Southern Brazil. *Lat Am J Pharm* 2008; 27(6): 894-9.
25. Oboh FOJ, Masodje HI and Enaabilele SA: Nutritional and antimicrobial properties of *Ocimum gratissimum* leaves. *J Bio Sci* 2009; 9(4): 377-380
26. Akunjobi CO, Ogbulie JN and Njoku HO: The extracts of *Ocimum gratissimum* on the dehydrogenase activity to clinical isolates of *Escherichia coli* and *Staphylococcus aureus*. *J Agric Tech* 2010; 6: 57-65
27. Abonkhai I, Ba A, Edogun O and Mu U: Antimicrobial activity of the volatile oils of *Ocimum basilicum* L. and *Ocimum gratissimum* L. (Lamiaceae) against some aerobic dental isolates. *Park J Pharm Sci* 2009; 22: 405-409
28. Abonkhai I, Ba A, Edogun O and Mu U: Antimicrobial activity of the volatile oils of *Ocimum basilicum* L. and *Ocimum gratissimum* L. (Lamiaceae) against some aerobic dental isolates. *Park J Pharm Sci* 2009; 22: 405-409
29. Tubaro A, Dri P, Delbello G, Zilli C and Della LR: The croton oil ear test revisited. *Agents Act* 1985; 17: 347-349.
30. Okoye FBC, Osadebe PO, Proksch P, Edrada-Ebel RA, Nworu CS and Esimone CO: Anti-inflammatory and membrane-stabilizing stigmastane steroids from *Alchornea floribunda* leaves. *Planta Medica* 2010; 76(2): 172-177.
31. Jadhav B, Khandelwal K, Ketkar A and Pisal S: Formulation and evaluation of mucoadhesive tablets containing eugenol for the treatment of periodontal diseases. *Drug Dev Ind Pharm* 2004; 30: 195-203.
32. Kavimani S, Vetrichelvan T, Ilango R and Jaykar B: Anti-inflammatory activity of the volatile oil of *Toddalia asiatica*. *Ind J Pharm Sci* 1996; 58(2): 67-70.
33. Kavimani S, Karpagam RI, Jaykar B and Ilango R: Anti-inflammatory activity of volatile oil of *Psidium guajava*. *Ind J Pharm Sci* 1997; 59(3): 142-144.
34. Mariea DEP, Dejanb B and Quetin-Leclercq J: GC-MS Analysis of the Leaf Essential Oil of *Ipomea pes-caprae*, a Traditional Herbal Medicine in Mauritius. *Nat Prod Comm* 2007; 2(12): 1225-1228.
35. Shimizu M, Shogawa H and Matsuzawa T: Anti-inflammatory constituents of topically applied crude drugs. IV Constituents and anti-inflammatory effect of Paraguayan crude drug “Alhucema (*Lavandula latifolia* - Vill). *Chem Pharm Bull* 1990; 38: 2283-2284.

36. Souza M, Siani A, Ramos, M, Menezes-de-Lima O and Henriques M: Evaluation of anti-inflammatory activity of essential oils from two Asteraceae species. *Pharmazie* 2003; 58: 582-586.
37. Sen P: Therapeutic potentials of Tulsi: from experience to facts. *Drug News and Views* 1993; 1(2): 15-21.
38. Prakash P and Gupta N: Therapeutic uses of *Ocimum sanctum* Linn (Tulsi) with a note on eugenol and its pharmacological actions: a short review. *Indian J Physiol Pharmacol* 2005; 49(2): 125-131.

How to cite this article:

Festus-Okoye BC, Willfred OO, Felix A, Ogheneogaga IO, Peace O, Ngozi IN, Jane A and Ndu-Okechukwu O: Chemical composition and anti-inflammatory activity of essential oils from the leaves of *ocimum basilicum l.* and *ocimum gratissimum l. (lamiaceae)*. *Int J Pharmacognosy* 2014; 1(1): 59-65. doi: 10.13040/IJPSR.0975-8232.1(1).59-65.

This Journal licensed under a Creative Commons Attribution-Non-commercial-Share Alike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)