



Received on 10 July 2018; received in revised form, 10 August 2018; accepted, 19 August 2018; published 01 September 2018

## STUDY OF ANTHELMINTIC AND INSECTICIDAL ACTIVITY OF *BACCAUREA RAMIFLORA* PLANT IN DIFFERENT EXTRACTS

Kazi Nuruddin Al-Masud<sup>\*</sup>, Zarif Morshed, Md. Nasiful Islam, Mahboob Hossain, Maliha Tasnim Deeba and Rezowana Islam

Department of Pharmacy, BRAC University, Dhaka, Bangladesh.

### Keywords:

*Baccaurea ramiflora*,  
Anthelmintic, Insecticidal,  
*Pheretima posthuma*,  
*Sitophilus oryzae*

### Correspondence to Author: Kazi Nuruddin Al-Masud

Department of Pharmacy,  
BRAC University, Dhaka,  
Bangladesh.

E-mail: nuruddinmasud123@gmail.com

**ABSTRACT:** *Baccaurea ramiflora* Lour., syn. *Baccaurea sapida* (Roxb.) Muell. Arg. known as Burmese grapes belongs to the family Euphorbiaceae and is native to Southeast Asia. The observance on *Baccaurea ramiflora* leads that this plant is utilized as an antichloristic and anodyne against rheumatoid arthritis, abscesses, cellulitis and treat injuries. This plant also has anti-diarrheal, analgesic, anti-inflammatory, neuro-pharmacological, antioxidant and cytotoxic activities. Therefore, the present study was conducted to evaluate anthelmintic and insecticidal activities of different extract of *Baccaurea ramiflora*. For anthelmintic activity, *Pheretima posthuma* was selected as test animal while 25, 50, 100 mg/ml concentrations of samples were tested in the bioassay, from which time of paralysis and time of death of worms were estimated. Evaluation of insecticidal activity was performed against *Sitophilus oryzae* to calculate the mortality rate. Extracts of MEE, EE, ACE, CHE, PEE, and NHE were for the activity determination. In the anthelmintic study, extract exhibited its activity in a dose-dependent manner showing higher the concentration, faster the effect. Extract of EE, PEE, CHE and MEE in case of 100 mg/ml concentration exhibited its paralytic effect followed by death within a short period among which ACE extract gave the best result which only took approx. 22.33 min to show paralytic effect and 33 min for death occurrence. In the insecticidal activity, all extracts showed potent activity with 100% mortality of rice insects *Sitophilus oryzae* at 80 mg/ml concentration in 24 h, proving the activity also followed a dose-dependent pattern. 80% mortality rate observed in case of 80 mg/ml concentration of ACE extract in 12 h whereas MEE, CHE, and NHE extracts showed 60% and PEE showed 70% mortality rate in treated rice insects. The present study shows that all the extracts of *Baccaurea ramiflora* are found to possess anthelmintic and insecticidal activities.

**INTRODUCTION:** Plants have been a source of medicinal substances for thousands of years.

Plants and phyto products continue to play a vital role in the treatment of various diseases. Drug discovery from plants is a multi-disciplinary approach which combines various botanical, ethnobotanicals, phytochemical and biological and chemical separation techniques.

However, despite these observations, about 80% of the world's inhabitants' problems are treated by medicinal herbal drug for their primary health care

	<p><b>DOI:</b> 10.13040/IJPSR.0975-8232.IJP.5(9).616-21</p>
	<p>The article can be accessed online on <a href="http://www.ijournal.com">www.ijournal.com</a></p>
<p>DOI link: <a href="http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5(9).616-21">http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5(9).616-21</a></p>	

<sup>1</sup>. Nowadays, the occurrences of parasitic infections are increasing extensively. Despite having numerous antibiotics available to encounter infections, antibiotics resistance has limited its use.

As a result, an alternative option can be phytotherapy to beat nematodes. At the same time, to invade different pathogenic organism's medicinal plants can be a potential choice to defeat infectious diseases <sup>2</sup>. *Baccaurea ramiflora* (Lour. family of Euphorbiaceae) is a resourceful plant which has the number of uses. The familiar names include Bhubi or Latkan (Bengali), Mafai (Thai), Leteku (Hindi) and Burmese grape. The slow-growing evergreen tree of *Baccaurea ramiflora* (*B. ramiflora*) has fruit (1-2" around), and the fruit is yellow to red. This fruit tree is native to the Southeast Asian region and found growing wild in South China, Indo-China, India, Nepal, Myanmar, the Andaman Islands, Thailand and Peninsular Malaysia <sup>3, 4</sup>. In Chinese Dai medicine, the whole plant of *Baccaurea ramiflora* is utilized as an antiphlogistic and anodyne against rheumatoid arthritis, cellulitis, abscesses and to treat injuries <sup>5</sup>.

The plant is also used as medicine by hill tribes in Northern Thailand <sup>6</sup>. Young leaves of *Baccaurea ramiflora* are used as a vegetable, flavoring agent with curries and minced meat in Bangladesh <sup>7</sup>. *B. ramiflora* (Lour) is such an underexploited fruit crop grown mainly in backyard plantation and as a forest plant. Research on *B. ramiflora* has been reported for its ethnobotanical uses, seed biology, and its isolated chemical constituents of essential oil. Three novel and four recognized compounds were isolated from the *B. ramiflora* stems <sup>1</sup>.

The two new phenols, 6'- O-vanilloylisotachioside and 6'- O-vanilloyltachioside, together with nine known compounds, were isolated from the leaves of *B. ramiflora* (Euphorbiaceae) <sup>6</sup>. The ethanol extractives of the leaves of *Baccaurea ramiflora* possess cytotoxic activity <sup>8</sup>.

A study on this plant established that the leaves of *Baccaurea ramiflora* possess hypoglycemic, hypolipidemic <sup>9</sup>. Another study supports the mechanism involved in the anti-inflammatory and antioxidant activity of *Baccaurea ramiflora* <sup>10</sup>. So, the extract of *Baccaurea ramiflora* was used to determine the insecticidal and anthelmintic property of the plant.

## MATERIALS AND METHODS:

**Collection of the Plant:** The plant *Baccaurea ramiflora* was collected from the local area of Comilla, Bangladesh in August 2017. The plant was freed from materials like dust, dirt, pollen. Then the plant was identified by Bushra Khan, Principal Scientific Officer, Bangladesh National Herbarium, Mirpur, Dhaka, and a voucher specimen has been deposited (DACB:46,964) for further reference.

**Extraction of Plant Material:** The plant parts were dried under the sun for a few days, and finally oven dried to remove all the moisture content. Then the plant parts were crushed to a coarse consistency. The coarse plant parts were extracted in a decreasing polarity order. The coarse plant material (900 g) was taken and soaked with 1500 ml of methanol for 3 consecutive days at 25 °C. The extract was filtered, and the filtrate was kept for further extraction. In the same manner, the filtrate was soaked in different solvents by polarity decreasing order.

Methanol > Ethanol > Acetone > Chloroform > Petroleum ether > *n*-hexane

For every case, the extract was preserved, and solvent evaporation was done by using rotary evaporator. Finally, all the extracts of *Baccaurea ramiflora* were kept under laminar airflow for protecting it from any type of contamination.

**Drugs and Chemicals:** Albendazole was assorted from Eskayef Bangladesh Ltd., 0.9% sodium chloride solution (normal saline) was purchased from IBN SINA Pharmaceutical Industry Ltd., Bangladesh, and other reagents were of analytical grade.

**Earthworms and Insects:** *Pheretima posthuma* (*P. posthuma*), a species of adult earthworm has a pretty high level of semblance with the human intestinal roundworm parasite from the anatomical and physiological point of view. Which lead to choosing this particular earthworm for the evaluation of the *in-vitro* anthelmintic activity. Mohakhali area of Dhaka, Bangladesh were chosen for the collection of the adult earthworms and moist soil was preferred as a collection site. All kinds of fecal matters were removed using saline water. The earthworms were of 3-5 cm in length and 0.1-0.2

cm in width as per in size. A very known and communal rice weevil, *Sitophilus oryzae* (*S. oryzae*), was collected from a local rice godown Bangladesh for detecting the insecticidal activity of *Baccaurea ramiflora*. For the insect collection, old rice was preferred.

**Anthelmintic Activity:** Development of anthelmintic resistance in helminths reported in several countries gives a clear indication that control programs based exclusively on their use are not sustainable<sup>11, 12, 13, 14, 15</sup>. The development of integrated programs to control helminths is vital, but such control programs require viable alternatives to the use of anthelmintics<sup>16</sup>. *Pheretima posthuma*, an adult earthworm, was chosen for the determination of anthelmintic activity. For every particular extract of *Baccaurea ramiflora* sample solution was prepared at 25, 50 and 100 mg/ml concentration in normal saline water and for each concentration approximately six similar sizes of earthworms (*Pheretima posthuma*) were placed in each beaker containing 50 ml of above test solutions of extract. As a reference standard, Albendazole (10 mg/ml) was used, and normal saline water as a control. Solutions of each concentration were freshly prepared just before conduction the experiment. The time required for the paralysis was noted when there was no movement from the worms despite these being shaken vigorously<sup>17</sup>. Time for the death of worms was recorded after ascertaining that worms neither moved when shaken vigorously nor when dipped in warm water (50 °C).

**Insecticidal Activity:** Insecticidal activity of many plants against several insect pests has been demonstrated<sup>18, 19, 20</sup>. Yang and Tang (1988) reviewed the plants used for pest insect control and found that there is a strong connection between medicinal and pesticidal plants<sup>21</sup>. For the beginning, the surface film activity test Petri dishes of 60 mm were taken for the extract and their replication. For every particular extract of *Baccaurea ramiflora* sample solution was prepared at (2.5, 5, 10, 20, 40, 50, 60, 70, 80 mg/ml). Now in every respective petri dish, the prepared sample solutions were poured into the lower part of it and kept for a few moments for it to dry out. Then insects were released in each of the treated petri dishes. A control petri dish was also kept similarly.

Soon performing all the steps mentioned above, the Petri dishes were kept in a sealed environment for the observance of the insecticidal activity. The mortality was observed first at 30 min and then after 12 and 24 h of exposure. After 24 h, the data was recorded. For the verification of the movement and death of insects, a simple microscope was used. In some cases, the hot needle was taken closer to the insect bodies (without movement) for the confirmation of death. Recovery of the insects if occurred, was also taken into consideration. The mortality rates of *Sitophilus oryzae* adults were corrected by Abbott's formula<sup>22</sup>.

$$\% \text{ of Mortality} = [(Ca - Ta) / Ca] \times 100$$

Where, Ca = No. of live control insects after treatment, Ta = No. of live test insects after treatment.

**Statistical Analysis:** All assays were performed in triplicate under strict aseptic conditions to ensure consistency of all findings. For each extract triplicate data was taken, and the final data was taken by the triplicate data's mean  $\pm$  SD (Standard Deviation), which was analyzed by Microsoft Excel. The bar diagram was designed using Prism software.

## RESULTS AND DISCUSSION:

**Anthelmintic Activity:** Although there has been a remarkable advancement in the field of human medicines observed, still several infectious diseases triggered by viruses, bacteria, fungi, parasites, insects, rodents and helminths are becoming a threat to communities' well-being. To be more specific, developing countries are at higher risk for their persistence. Exclusively, children and teenagers are susceptible to infectious diseases. The comparative anthelmintic drug resistance incidence has led to the increasing demand of the natural anthelmintic as a potential substitute for diseases management<sup>23, 24</sup>.

The present study demonstrates the anthelmintic activity of the plant extract of *Baccaurea ramiflora* using adult earthworms because of the high level of semblance with the human intestinal roundworm parasite. When the earthworms (*Pheretima posthuma*) were exposed to the *Baccaurea ramiflora* extract, they started gradually losing their motility by providing stimulant effect primarily.

Ultimately, dose-dependent paralysis was observed followed by the occurrence of death in each crude extract containing 25, 50, and 100 mg/ml. **Table 1** represents the effects of *Baccaurea ramiflora* in different extracts. The extract denoted anthelmintic activity in a dose-dependent manner, describing the higher concentration of each extract exhibited a paralytic effect much earlier and resulted in death within a short period.

As per the data denoted in **Table 1**, it reflects that at 25 mg/ml concentration in all extracts (MEE, EE, ACE, CHE, PEE, and NHE) the time for paralysis and ultimately the time required for the death worms took more time comparing to 50 and 100 mg/ml concentration. The standard reference, Albendazole, showed strong anthelmintic action. As a whole, different extracts of *Baccaurea*


*ramiflora* showed anthelmintic activity in a dose-dependent manner<sup>24</sup>. Previous phytochemical screening reported different extraction extracts of *B. ramiflora* to contain carbohydrates, tannins, flavonoids, proteins, alkaloids and<sup>25</sup>.

Tannins being polyphenolic compounds are known to have anthelmintic activities. The reports on the anthelmintic effect of tannins are due to binding of tannins to free proteins in the gastrointestinal tract of host animal or glycoprotein on the cuticle of the parasite and may be responsible for death<sup>26</sup>. Secondary metabolites like flavonoids also possess anthelmintic activity, but their exact mechanism is yet to be known<sup>27</sup>. So the presence of tannins and secondary metabolites like flavonoids present in the extract of *Baccaurea ramiflora* produced similar anthelmintic effects.

**TABLE 1: EFFECT OF DIFFERENT EXTRACTS OF *BACCAUREA RAMIFLORA* ON *PHERETIMA POSTHUMA***

Group	Dose (mg/ml)	No. of worms	Time taken for paralysis	Time taken for death
Control	20ml	6	-	-
Standard	10	6	28.33 ± 1.15	35 ± 1.73
MEE	100	6	27 ± 2.64	43.33 ± 2.52
	50	6	32.66 ± 1.52	55 ± 2.65
	25	6	51.33 ± 1.52	72.66 ± 1.15
	100	6	23.66 ± 3.05	34.33 ± 2.52
EE	50	6	34.66 ± 4.04	47.66 ± 1.53
	25	6	47.66 ± 0.57	62 ± 2.65
	100	6	22.33 ± 2.08	33 ± 2
ACE	50	6	43 ± 2.64	63.66 ± 3.51
	25	6	46.66 ± 3.21	75.66 ± 4.50
	100	6	31.33 ± 2.08	40.33 ± 1.53
CHE	50	6	37.66 ± 1.15	55.66 ± 2.52
	25	6	44 ± 1	62.33 ± 1.15
	100	6	23.66 ± 4.72	32 ± 2
PEE	50	6	35.66 ± 3.05	47.33 ± 1.53
	25	6	44.33 ± 1.15	55.33 ± 2.52
	100	6	31 ± 2	46.33 ± 2.30
NHE	50	6	39.33 ± 1.52	52.33 ± 3.05
	25	6	50.66 ± 2.51	70.66 ± 2.52
	100	6	50.66 ± 2.51	70.66 ± 2.52

Control group received saline water 20 ml, the standard group received Albendazole 10 mg/ml and test groups MEE, EE, ACE, CHE, PEE, and NHE were treated with 25, 50 and 100 mg/ml of the extracts respectively. MEE=Methanol extract, EE=Ethanol extract ACE= Acetone extract, CHE=Chloroform extract, PEE= Petroleum ether extract, NHE=*n*-hexane extract.



**FIG. 1: GRAPHICAL REPRESENTATION OF OCCURRENCE OF DEATH (IN MIN)**

**Insecticidal Activity:** A major interference in the use of insecticides is resistance developed by insects to fight against diseases due to its fewer efficacies. So, a potent substitution can be natural insecticides prepared from plants since they contain several bioactive chemicals<sup>28</sup>. The present study demonstrates the insecticidal activity of the plant extract of *Baccaurea ramiflora* against rice induced insect named *Sitophilus oryzae*.

For every particular extract of *Baccaurea ramiflora* sample solution was prepared at (2.5, 5, 10, 20, 40, 50, 60, 70, 80 mg/ml) and for evaluation was observed for 12 and 24 h. Among all the extracts, ACE showed the highest mortality (80%) whereas MEE, CHE, NHE extracts provided % percentage of mortality of (60%) and PEE (70%) at 80 mg/ml in 12 h **Table 3**. After 24 h, at 80 mg/ml in all extracts exhibited 100% **Table 3**.

The mortality of the insect was found to be concentration dependent. It is observed that the carbohydrates, saponins, phytosterols, phenols, flavonoids, and tannins are having mosquito larvicidal activity<sup>29</sup>. So the presence of tannins and secondary metabolites like flavonoids present in the extract of *Baccaurea ramiflora* plant may describe the toxic effects on the insects.

**TABLE 2: INSECTICIDAL ACTIVITY OF *BACCAUREA RAMIFLORA* ON *S. ORYZAE* (DEATH OF INSECTS)**

Sample Conc. mg/ml	No. of insect taken	No. of insects dead											
		MEE		EE		ACE		CHE		PEE		NHE	
		12h	24h	12h	24h	12h	24h	12h	24h	12h	24h	12h	24h
2.5	10	0	1	0	0	0	1	1	3	0	2	0	0
5	10	1	3	0	0	0	2	1	4	1	2	0	2
10	10	1	4	1	3	2	4	2	4	1	2	1	3
20	10	2	5	1	3	3	4	3	4	2	4	2	4
40	10	2	5	2	6	5	6	4	6	3	5	3	5
50	10	3	5	3	6	5	7	5	7	5	7	3	5
60	10	4	6	3	7	7	9	5	7	6	7	4	7
70	10	5	7	4	9	8	10	5	8	6	8	5	7
80	10	6	10	5	10	8	10	6	10	7	10	6	10

**TABLE 3: INSECTICIDAL ACTIVITY OF *BACCAUREA RAMIFLORA* ON *S. ORYZAE* (% OF MORTALITY)**

Sample Conc. mg/ml	% of Mortality											
	MEE		EE		ACE		CHE		PEE		NHE	
	12h	24h	12h	24h	12h	24h	12h	24h	12h	24h	12h	24h
2.5	0	10	0	0	0	10	10	30	0	20	0	0
5	10	30	0	0	0	20	10	40	10	20	0	20
10	10	40	10	30	20	40	20	40	10	20	10	30
20	20	50	10	30	30	40	30	40	10	40	20	40
40	20	50	20	60	50	60	40	60	20	50	30	50
50	30	50	30	60	50	70	50	70	30	70	30	50
60	40	60	30	70	70	90	50	70	60	70	40	70
70	50	70	40	90	80	100	50	80	60	80	50	70
80	60	100	50	100	80	100	60	100	70	100	60	100

**CONCLUSION:** The present study indicates the different plant extract of *Baccaurea ramiflora* possess a varying degree of anthelmintic activity and toxic principles with the presence of insecticidal activity. The anthelmintic study shows the acetone extract provides the fastest paralysis effect (22.33 min) followed by the fastest death time (33 min) in 100 mg/ml concentration. Whereas chloroform and *n*-hexane extract provides the slowest paralysis time and death occurrence time 100 mg/ml concentration.

The insecticidal study demonstrates that the acetone extract provides the highest % percentage of mortality of 80% in 12 h at 80 mg/ml concentration comparing with the other extract. Whereas MEE, CHE, NHE extracts provided % percentage of mortality of 60% and PEE 70% in 12

h at 80 mg/ml concentration. All the extract provided 100% mortality in 24 h at 80 mg/ml concentration. Therefore, the observed biological activities can play a great role for further detailed investigation of other particular activities of different plant extracts to signify the research on medicinal plants.

**ACKNOWLEDGEMENT:** Nil

**CONFLICT OF INTEREST:** Nil

**REFERENCES:**

1. Ali MR, Hossain M, Runa JF and Hasanuzzaman M: Preliminary cytotoxic activity of different extracts of *Averrhoa bilimbi* (fruits). International Current Pharmaceutical Journal 2013; 2(3): 83-84.
2. Dash PR, Mou KM, Erina IN, Ripa FA, Masud KNA and Ali MS: Study of anthelmintic and insecticidal activities of

- different extracts of *Kaempferia galanga*. International Jour of Pharmaceutical Science and Research 2017; 8(2).
3. Khan B: Encyclopedia of Flora and Fauna of Bangladesh. Asia Society of Bangladesh, Dhaka, Edition 1, Vol. 7, 2008: 392-3.
  4. Hoang SV, Pieter B and Keler PJA: Uses and conservation of plant species in a national park- a case study of Ben En, Vietnam. Eco Botany 2008; 62(4): 574-93.
  5. Lin YF, Yi Z and Zhao YH: Chinese Dai Medicine Colorful Illustrations. Yunnan Nationality Press, Edition 1<sup>st</sup>, 2003.
  6. Yang XW, Wang JS, Ma YL, Xiao HT, Zuo YQ, Lin H, He HP, Li L and Hao XJ: Bioactive Phenols from the Leaves of *Baccaurea ramiflora*. Planta Medica 2007; 73: 1415-17.
  7. Hasan SMR, Hossain MM, Akter R, Jamila M, Mazumder MEH and Rahman S: DPPH free radical scavenging activity of some Bangladeshi medicinal plants. Journal of Medicinal Plants Research 2009; 3(11): 875-79.
  8. Howlader A, Sultana S, Ghosh CK and Bachar SC: Oral hypoglycemic effect of *Baccaura ramiflora* Linn. (Phyllanthaceae) bark methanolic extract on mice. Journal of Southeast University Sci and Engineering 2009; 3: 89-92.
  9. Obayedullah M, Urmi KF, Howlader MA, Hossain MK, Ahmed MT and Hamid K: Hypoglycemic, hypolipidemic and antioxidant effects of leaves methanolic extract of *B. ramiflora*. Int. J. Pharm Pharm. Sci. 2012; 4(3), 266-69.
  10. Usha T, Middha SK, Bhattacharya M, Lokesh P and Goyal AK: Rosmarinic acid, a new polyphenol from *Baccaurea ramiflora* Lour. leaf: A probable compound for its anti-inflammatory activity. Antioxidants 2014; 3(4): 830-42.
  11. Jackson F: Anthelmintic resistance-the state of play. British Vet. J. 1993; 149: 123-38.
  12. Sanyal PK: Gastrointestinal parasites and small ruminant production in India. Sustainable parasite control in small ruminants. Proc. Workshop Bogor, Indonesia, ACIAR Proceedings no. 74, 1996; 22-25, 109-12.
  13. Rolfe PF: Anthelmintic resistance in Australia, its development and management. Proc. 4<sup>th</sup> Intl. Cong. Sheep Vet., Armidale, Australia 1997: 51-8.
  14. VanWyk JA, Malan FS and Randles JL: How long before resistance makes it impossible to control some field strains of *Haemonchus contortus* in South Africa with any of the modern anthelmintics? Vet. Parasitol. 1997; 70: 111-22.
  15. Waller PJ: The global perspective of anthelmintic resistance in nematode parasites of sheep-excluding Australasia. Proc. 4<sup>th</sup> Intl. Cong. Sheep Vet., Armidale, Australia 1997: 59-63.
  16. Waller PJ: International approaches to the concept of integrated control of nematode parasites of livestock. Int J Parasitol. 1999; 29: 155-64.
  17. Kumanan R, Sridhar C, Jayaveera KN, Sudha S, Rubesh Kumar S and Duganath N: Comparative study of anthelmintic activity of different leaf extracts of *Tecoma stans* (L.) on adult Indian earthworms. International Journal of Pharmaceutical and Clinical Res 2015; 2(2): 63-65.
  18. Jilani G and Su HCF: Laboratory studies on several plant materials as insect repellent for protection of cereal grains. J Econ Entomol 1983; 76: 154-57.
  19. Isman MB: Plant essential oils for pest diseases management. Crop Prot 2000; 19: 603-08.
  20. Carlini CR and Grossi-de Sá MF: Plant toxic proteins with insecticidal properties: A review on their potentialities as bioinsecticides. Toxicon 2002; 40: 1515-1539.
  21. Yang RZ and Tangs CS: Plants used for pest control in China: a literature review. Econ Bot 1988; 42: 376-06.
  22. Khatun MH, Nesa ML, Hosen AH, Bashar S, Sarker S and Islam MR: Evaluation of the antibacterial, cytotoxic and insecticidal activities of *Hibiscus sabderiffa* barks. International Journal of Pharmacy and Research 2015; 5(3): 170-75.
  23. Kumar MBS, Kumar MCR, Bharath AC, Kumar HRV, Kekuda TRP, Nandini KC, Rakshitha MN and Raghavendra HL: Screening of selected biological activities of *Artocarpus lakoocha* roxb (Moraceae) fruit pericarp. Journal of Basic and Clinical Pharmacy 2010; 1(4): 239-45.
  24. Riaz M, Rahman N and Haq MZU: Anthelmintic and insecticidal activity of *Verbascum thapsus* L. Pakistan Journal of Zoology 2013; 45(6): 1593-98.
  25. Goyal AK, Mishra T and Sen A: Antioxidant profiling of Latkan (*Baccaurea ramiflora* Lour.) wine. Indian Journal of Biotechnology 2013; 12(1): 137-39.
  26. Mulla WA, Thorat VS, Patil RV and Burade KB: Anthelmintic activity of leaves of *Alocasia indica* Linn. International Jou of Pharm Tech Res 2010; 2(1): 26-30.
  27. Kerboeuf D, Riou M and Guegnard F: Flavonoids and related compounds in parasitic disease control. Bentham Science Publisher 2008; 8: 116-28.
  28. Kiran R, Kekuda TRP, Kumar HGP, Hosetti BB and Krishnaswamy K: Biological activities of *Sarcanthus pauciflorus*. Journal of Applied Pharmaceutical Science 2013; 3(7): 105-10.
  29. Khanna VG and Kannabiran K: Larvicidal effect of *Hemidesmus indicus*, *Gymnema sylvestre* and *Elipta prostrata* against *Cules quinquefasciatus* mosquito larvae. African Journal of Biotechnology 2007; 6(3): 307-11.

**How to cite this article:**

Al-Masud KN, Morshed Z, Islam MN, Hossain M, Deeba MT and Islam R: Study of anthelmintic and insecticidal activity of *Baccaurea ramiflora* plant in different extracts. Int J Pharmacognosy 2018; 5(9): 616-21. doi link: [http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5\(9\).616-21](http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.5(9).616-21).

This Journal licensed under a Creative Commons Attribution-Non-commercial-Share Alike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)