



Received on 28 February 2020; received in revised form, 24 April 2020; accepted, 28 April 2020; published 30 April 2020

PHARMACOGNOSTICAL EVALUATION AND STANDARDIZATION OF AERIAL PARTS OF *SPHAERANTHUS INDICUS* LINN. (ASTERACEAE)

Ankit Sharma* and Siddhraj S. Sisodia

Bhupal Nobles' College of Pharmacy, Bhupal Nobles' University, Udaipur - 313001, Rajasthan, India.

Keywords:

Aqueous extract, Total flavonoids, Total phenolics, Ayurvedic system of medicine, *Sphaeranthus indicus*

Correspondence to Author:

Ankit Sharma

Senior Research Fellow,
M. Pharma (Pharmacology),
Regional Ayurvedic Research
Institute for Drug Development,
Aamkho, Gwalior - 474009, Madhya
Pradesh, India.

E-mail: rasthehearts@gmail.com

ABSTRACT: *Sphaeranthus indicus* Linn (Asteraceae) is a popular and very useful plant used in Indian system of medicine. It is usually found in damp areas, plains and also as a weed in the rice fields. In the Ayurvedic system of medicine, the plant as a whole plant or its different parts like leaf, stem, bark, root, flower and seed are widely used for curing many diseases. The plant is enriched with various types of secondary metabolites such as eudesmanolides, sesquiterpenoids, sesquiterpene lactones, sesquiterpene acids, flavone glycosides, flavonoid C-glycosides, isoflavone glycoside, sterols, sterol glycoside, alkaloid, peptide alkaloids, amino acids and sugars. Essential oil has been isolated from flowers and whole plant. In this present study, various methods are carried out to perform the evaluation of pharmacognostical parameters of crude drug as well as aqueous extract of aerial part of the plant which may forms the basis for the further use of the plant and also for the evaluation of its various pharmacological activity. The physicochemical evaluation confirms the quality and purity of the crude drug. Aqueous extract of *Sphaeranthus indicus* (AESI) was found to be rich in presence of alkaloids, proteins, amino acids, saponins, flavonoids and other phenolic compounds while glycosides, sterols, carbohydrates and volatile oils were found to be absent. The quantitative estimation of phytoconstituents viz. total flavonoids and total phenolics and saponins revealed that AESI was found rich in total flavonoids while less amount of total phenolic content while total saponin was present in fair amount. This present study can be a basis of further evaluation of various pharmacological activity of the aqueous extract of *Sphaeranthus indicus*.

INTRODUCTION: *Sphaeranthus indicus* Linn. (Asteraceae) is an aromatic annual spreading herb, found throughout India, at an altitude of 1500 m msl¹. It is known by different names in different parts of India such as Gorakhmundi in Hindi, East Indian globe thistle in English and Mahamundi, Mundi, Shrivani in Sanskrit².

In Ayurvedic system of medicine, at different dosages and as different preparations it is used in treatment of tuberculosis, lung disorders such as bronchitis and asthma, indigestion, spleen disorders, elephantiasis, leukoderma, vomiting, urinary discharge, hemicranis, epileptic convulsions, blood purifier, as a poultice in rheumatic pain, aphrodisiac, retaining of pregnancy and looseness of the breasts³.

It also has been pharmacologically or traditionally found to be an effective drug for treatment of various disorders, as an hepatoprotective agent⁴, antihelmintic agent⁵, antigout activity⁶, antitussive

	QUICK RESPONSE CODE DOI: 10.13040/IJPSR.0975-8232.IJP.7(4).116-20
	The article can be accessed online on www.ijpjournal.com
DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.7(4).116-20	

agent ⁷, antioxidant agent ⁸, as an renal protective agent ⁹, as immunostimulant agent ¹⁰ and as antidiabetic agent ¹¹.

MATERIALS AND METHODS:

Collection of Plant: The aerial parts of *Sphaeranthus indicus* Linn. (Asteraceae) were procured in the month of December 2018 from the village area nearby Vidisha (M.P.). This plant was identified and authenticated by Dr G. P. Sinha, Scientist-E and Head of Office, Botanical Survey of India, CRC, Allahabad, Uttar Pradesh - 211002. A voucher specimen (Accession No. 104180) of *Sphaeranthus indicus* was deposited in the herbarium of the Institute for future reference.

Preparation of Aqueous Extracts of *Sphaeranthus indicus* (AESI): The powder drug (30 g) was soaked in 1 liter purified water and kept in dark and dry place for 48 h at room temperature. Chloroform was added in quantity of 1% total mixture to avoid the microbial growth. After 48 h, solutions were filtered by Whatman filter paper no. 1. The filtered extracts were dried in a rotary evaporator and percentage yield was calculated.

Standardization of Crude Drug: ¹²

Determination of Ash Value: The determination of ash values is meant for detecting low-grade products, exhausted drugs and sandy or earthy matter. It can also be utilized as a mean of detecting the chemical constituents by making use of total ash and acid insoluble ash.

Total Ash: The air-dried powder drug (4 g) was weighed accurately in a previously ignited and tarred silica dish. The material was spread evenly and ignited in a muffle furnace at 600 °C until it is white, indicating the absence of carbon. Then the dish was cooled in a desiccator and weighed. If carbon-free ash cannot be obtained by this manner, the residue on the cool dish was moistened with about 2 ml of water or a saturated solution of ammonium nitrate, dried on a water-bath, and then ignited in the muffle furnace up to constant weight. The percentage of total ash was calculated using the following formula.

Percentage of total ash = Weight of ash / Weight of sample taken × 100

Acid Insoluble Ash: Total ash of the sample, as described above, was determined out first. To the

dish containing the total ash, 45 ml of 1:5 hydrochloric acid in three portions of 15 ml was added each time, boiled gently for 5 min and filtered. The insoluble matter was collected on an ashless filter paper (Whatman no. 41) and washed with distilled water until the residue is free from acid. The filter paper containing the insoluble matter was transferred to the original dish, dried and ignited to constant weight. After cooling the dish in desiccators, the weight was taken. The percentage of acid-insoluble ash was calculated using the following formula.

Percentage of acid-insoluble ash = Weight of acid-insoluble residue / Weight of sample taken × 100

Determination of Extractive Value: Plant materials are composed of water, mineral, and organic compound primary metabolites and secondary metabolites. These compounds are extracted in different solvents in various proportions. Extractive value is the amount of active constituents extracted with solvents from a given amount of medicinal plant material. Alcohol soluble extractive the air-dried powder drug 3 g was weighed accurately in a glass stopper flask. To this 100 ml of distilled ethanol, approx. 95% was added and shaken occasionally for 6 h. After keeping for 18 h, it was filtered (rapidly taking care not to lose any solvent). Filtrate (25 ml) was pipette out in a pre-weighed 100 ml dish and evaporated to dryness on a water bath, after which it was kept in a hot air oven at 105 °C for 6 h. After cooling the flask in a desiccator, weight was taken. The percentage of alcohol-soluble extractive was calculated using the following formula.

Alcohol soluble extractive % = Weight of the extract / 25 × Weight of sample taken × 100 × 100

Water-Soluble Extractive: The air-dried powder drug 3 g was weighed accurately in a glass stoppered flask. To this, 100 ml of distilled water was added and shaken occasionally for 6 h. After keeping for 18 h it was filtered. Filtrate 25 ml was pipette out in a pre-weighed 100 ml beaker and evaporated to dryness on a water bath, after which it was kept it in a hot air oven at 105 °C for 6 h. After cooling the flask in desiccators, weight was taken. The percentage of the water-soluble extractive matter was calculated using the following formula.

Water-soluble extractive % = $\frac{\text{Weight of the extract}}{\text{Weight of sample taken}} \times 25 \times 100$

Determination of Loss on Drying at 105 °C: Loss on drying is the loss in weight in % w/w resulting from water and volatile matter of any kind that can be driven off under specified conditions. The air-dried powder drug 2 g was weighed accurately in a previously weighed dish. Then it was heated in an oven at 105 °C for 5 h. After cooling the dish in desiccators, the weight was taken. The procedure was repeated until a constant weight is obtained. Loss on drying was calculated using the following formula.

Percentage of loss on drying at 105 °C = $\frac{\text{Loss in weight of the sample}}{\text{Weight of sample taken}} \times 100$

Determination of pH Value: The pH value of an aqueous liquid may be defined as the common logarithm of reciprocal of the hydrogen ion concentration expressed in gram per liter. The pH value of liquid is determined potentially by means of the glass electrode and a suitable pH meter. One gram of coarsely powdered drug was added in 100 ml of water, and the solution was filtered in a beaker, and pH was checked at 25 °C.

Thin Layer Chromatography (TLC) Analysis:

^{13, 11} One gram of powder of *Sphaeranthus indicus* aerial part was subjected to extraction in 25 ml methanol for 18 h at room temperature. The extract was filtered and used for TLC. Ten µl of the filtrate of *Sphaeranthus indicus* and geraniol was applied on Merck aluminum TLC plate and the plate was developed to a distance of 8 cm using toluene: ethyl acetate (7:3) as mobile phase. After development, the plate was allowed to dry in air and examined under ultraviolet light (254 and 366 nm) for the presence of spots. After this, the plate was sprayed with anisaldehyde sulphuric acid reagent, followed by heating at 105 °C for about 10 min. Spots were examined under visible light. TLC photos were captured at visible light, UV 254 nm, and UV 366 nm and after derivatization. The number of spots, colors of spots, and their R_f values were calculated.

$R_f = \frac{\text{Distance travel by solute}}{\text{Distance travel by solvent}}$

Preliminary Phytochemical Screening: ^{15, 16} The aqueous extract of the *Sphaeranthus indicus* (AESI) was tested for the presence of carbohydrates,

proteins, alkaloids, flavonoids, glycosides, saponins, tannins and essential oils using standard procedures.

Quantitative Estimation of Phytoconstituents:

Total Phenolic Content: ¹⁷ Total phenolic content of the extract was determined spectrometrically. One ml of Folin-Ciocalteu's reagent, previously diluted (1:20), was added to 1 ml of sample (AESI) (1000 µg/ml), tannic acid (10-100 µg/ml) mixed thoroughly. To the mixture, add 4 ml of sodium carbonate (75 g/l), and the total volume was made up to 10 ml with distilled water in a volumetric flask and mix well. The mixture was allowed to stand for 2 h at room temperature. Contents were then centrifuged at 2000 rpm for 5 min, and the absorbance of the supernatant was read at 760 nm. A standard curve was obtained using various concentrations of tannic acid. Results were expressed as mg of tannic acid equivalents (TAE) per gram of extract.

Total Flavonoids Content: ¹⁸ Total flavonoid content was measured by aluminum chloride colorimetric assay. One milliliter of the sample (AESI) (1000 µg/ml) or standard solution of quercetin (25-1000 µg/ml) was added to 10 ml volumetric flask containing 4 ml of distilled water. To the above mixture, 0.3 ml of 5% NaNO₂ was added. After 5 min, 0.3 ml of 10% AlCl₃ was added. At 6th min, 2 ml of 1 M NaOH was added, and the total volume was made up to 10 ml with distilled water. The solution was mixed well, and the absorbance was measured against prepared reagent blank at 510 nm. The total flavonoid content of the extracts was expressed as milligrams of quercetin equivalents per gram of extract.

Total Saponins Content: ¹⁹ The samples were ground and 3 g of each were put into a conical flask and 50 ml of 20% aqueous ethanol were added. The samples were heated over a hot water bath for 4 h with continuous stirring at about 55 °C. The mixture was filtered and the residue re-extracted with another 100 ml 20% ethanol. The concentrate was combined extract were reduced to 20 ml over water bath at about 90 °C. The concentrate was transferred into a 250 ml separating funnel then 10 ml ether was added and shaken well. Ether layer was discarded. The purification process was repeated. Thirty milliliter of n-butanol was added

and shaken well. n-butanol layer was collected. The combined n-butanol extracts were washed twice with 5 ml of 5% aqueous sodium chloride. The remaining solution was heated in a water bath. After evaporation the samples were dried in the oven to a constant weight. The content was calculated as mg saponins per 100 g of powdered material.

Total saponins content = Weight of extract / Weight of sample \times 100

RESULTS AND DISCUSSION: Results of the physico-chemical studies viz. ash content, extractive value, moisture content, and pH were as per the pharmacopoeial standard²⁰ and previous reports. This indicated that the crude drug is having the prescribed limits of all the stated parameters and suggested the appropriateness of the drug quality **Table 1**. TLC of methanol extract of the aerial parts of the *Sphaeranthus indicus* was run and compared with standard geraniol (active constituent of *Sphaeranthus indicus*). The TLC chromatogram revealed the presence of geraniol in the extract and thus authenticated the plant **Table 2**. The phytochemical screening of the plant extracts revealed the presence of alkaloids, proteins, amino acids, saponins, flavonoids and other phenolic compounds while glycosides, sterols, carbohydrates and volatile oils were found to be absent. The results were in accordance with the several previous reports mentioned earlier^{1, 2, 5, 21, 22, 23}. The quantitative estimation of phytoconstituents viz. total flavonoids and total phenolics and saponins revealed that AESI was found rich in total flavonoids (187.9 ± 10.32 quercetin equivalents mg/g of AESI) while less amount of total phenolic content (37.23 ± 0.70 tannic acid equivalents mg/g of AESI) while total saponin (3.33 g/100g powder drug) was present in fair amount. *S. indicus* Linn. is widely distributed throughout India.

The plant appears to have a broad spectrum of activity on several ailments. Traditionally, various parts of the plant have been found effective for anxiolytic activity, neuroleptic activity, immunomodulatory activity, anti-inflammatory activity, mast cell stabilizing action, antihyperglycemic activity, hepatoprotective activity, larvicidal action, bronchodilatory effect, antihyperlipidemic activity, renoprotective effect and many other miscellaneous activities. It is

reported to contain eudesmanoids, eudesmanolides, sesquiterpene lactone, sterol glycoside, flavanoids, and essential oil.

TABLE 1: PHYSICO-CHEMICAL EVALUATION OF SPHAERANTHUS INDICUS AERIAL PARTS

1	Ash analysis	
	ash content (Total ash)	9.407 ± 0.07623
	acid insoluble ash	6.692 ± 0.2866
2	Extractive value (Maceration process)	
	alcohol soluble	17.08 ± 0.4543
	water-soluble	26.46 ± 0.9702
3	Moisture content (Loss on drying)	11.89 ± 0.1158
4	pH (1% aqueous solution)	7.830 ± 0.06028

Values are expressed as Mean \pm SEM; n = 3

TABLE 2: TLC PROFILE OF SPHAERANTHUS INDICUS

S. no.	366 nm	
	Colour	R _f
1	Light pink	0.15
2	Light pink	0.19
3	Pink	0.27
4	Light pink	0.54
5	Pink	0.58
6	Light pink	0.62
7	Light pink	0.68
8	Light blue (geraniol)	0.73
9	Light pink	0.85
10	Pink	0.93

CONCLUSION: The pharmacological studies reported in this study confirms the therapeutic value of *Sphaeranthus indicus* Linn. However, less information is available regarding the clinical, toxicity, and phytoanalytical properties of this plant. Several phytochemical studies have been reported but still it needs to progress. With the availability of primary information, further studies can be carried out like clinical evaluation, phytoanalytical studies and toxicity evaluation. The plant is preclinically evaluated to some extent; if these claims are scientifically evaluated clinically, then it can provide good remedies and help the mankind in various ailments.

ACKNOWLEDGEMENT: Authors are grateful to Bhupal Nobles College of pharmacy, Udaipur and RARIDD, Gwalior for providing facilities for the fulfillment of this research work.

CONFLICTS OF INTEREST: None

REFERENCES:

- Galani VJ, Patel BG and Rana DG: *Sphaeranthus indicus* Linn. a phytopharmacological review. International Journal of Ayurveda Research 2010; 1: 247-53.

2. Ramachandran S: Review on *Sphaeranthus indicus* Linn. (Kottaiikkarantai). Pharmacogn Review 2013; 7: 157-69.
3. Mathew JE, Srinivasan KK, Dinakaran V and Joseph A: Mast cell stabilizing effects of *Sphaeranthus indicus*. J Ethnopharmacol 2009; 1(22): 394-6.
4. Sundari K, Govindaraju G and Bharathi B: Hepato-protective effect of ethanolic extracts of *Sphaeranthus indicus* (Linn.) on paracetamol induced liver toxicity in rats. Int J of App Bio and Pharma Tech 2011; 2: 315-21.
5. Sharma S, Jalalpure SS, Semwal B, Tandon S and Agarwal N: Anthelmintic activity of the whole plant of *Sphaeranthus indicus* Linn. International Journal of Ayurvedic and Herbal Medicine 2011; 1: 18-23.
6. Sharma A, Sisodia S and Narvariya V: Review on natural plant products used in treatment of gout. EJPMR 2019; 6(7): 629-33.
7. Nayak SS, Maity TK and Maiti BC: Antitussive and synergistic effects of sleeping time induced by standard sedatives of *Sphaeranthus indicus* plant extracts. Journal of Advanced Pharmaceutical Research 2010; 1: 123-32.
8. Tiwari BK and Khosa RL: Evaluation of hepatoprotective activity of *Sphaeranthus indicus* flower heads extract. Journal of Natural Remedies 2008; 8: 173-8.
9. Srinivasan VM, Jessy KK and Alex AE: Effect of *Sphaeranthus indicus* Linn. on gentamicin induced acute renal failure in rats. Indian J Pharmacol 2008; 40: 71.
10. Shekhani MS, Shah PM, Yasmin A, Siddiqui R, Perveen S and Khan KM: An immunostimulant sesquiterpene glycoside from *Sphaeranthus indicus*. Phytochemistry 1990; 29: 2573-6.
11. Pareek A, Suthar M, Bansal VK, Chauhan RK and Batra N: *In-vitro* study of *Sphaeranthus indicus* on glucose uptake by isolated rat hemidiaphragm. Journal of Pharmaceutical Research 2010; 3: 2284-5.
12. Anonymous: The Ayurvedic Pharmacopoeia of India Part-I. Ministry of Health and Family Welfare Department of Ayush Govt of India New Delhi 1996; 1: 213-4.
13. Sethi PD: High performance thin layer chromatography. CBS Publishers and Distributors Delhi 1996; 10.
14. Stahl I: Thin layer chromatography. a laboratory hand book. Student Edition Springer- Verlag Berlin 1969; 5-29.
15. Khandelwal KR: Practical Pharmacognosy. Edition 15th Nirali Prakashan Pune 2006; 149-56.
16. Khare CP: Indian medicinal plants, an illustrated dictionary. Springer New York 2007; 620-1.
17. Singleton VL and Rossi JRJA: Colorimetry of total phenolics with phosphomolybdic phosphotungstic acid reagents. Am J of Enolo and Viticulture 1965; 16: 144-58.
18. Marinova D, Ribarova F and Atanasova M: Total phenolics and flavonoids in Bulgarian fruits and vegetables. J of Univ Chem Tech and Met 2005; 40(3): 255-60.
19. Oboni BO and Ochuko PO: Phytochemical studies and comparative efficacy of the crud extracts of some Homostatic plants in Edo and Delta States of Nigeria. Global J of Pure and Applied Science 2002; 8: 203-8.
20. Anonymous: The Ayurvedic Pharmacopoeia of India Part I. Department of ISM and H Ministry of Health and Family Welfare Gove of India New Delhi 2001; 4: 78-80.
21. Shirwaikar A, Prabhu KS and Punitha IS: *In-vitro* antioxidant studies of *Sphaeranthus indicus* Linn. Indian Journal of Experimental Biology 2006; 44: 993-6.
22. Vikani KV, Dangar RD, Kapadia NS and Shah MB: A pharmacognostic study on *Sphaeranthus indicus*. Journal of Natural Remedies 2008; 8: 61-7.
23. Ambavade SD, Mhetre NA, Tate VD and Bodhankar SL: Pharmacological evaluation of the extracts of *Sphaeranthus indicus* flowers on anxiolytic activity in mice. Indian J Pharmacol 2006; 38: 254-9.

How to cite this article:

Sharma A and Sisodia S: Pharmacognostical evaluation and standardization of aerial parts of *Sphaeranthus indicus* Linn. (Asteraceae). Int J Pharmacognosy 2019; 7(4): 116-20. doi link: [http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.7\(4\).116-20](http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.7(4).116-20).

This Journal licensed under a Creative Commons Attribution-Non-commercial-Share Alike 3.0 Unported License.

This article can be downloaded to **Android OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Play store)